

CHAPTER 17

Culturally Adapted Preventive Interventions for Children and Adolescents

NANCY A. GONZALES, ANNA S. LAU, VELMA M. MURRY, ARMANDO A. PINA, and MANUEL BARRERA, JR.

WHY CULTURAL ADAPTATION IN DEVELOPMENTAL PSYCHOPATHOLOGY	874
DEFINITION OF CULTURE	875
HISTORICAL CONTEXT OF CULTURAL ADAPTATION RESEARCH	876
THE RATIONALE FOR CULTURAL ADAPTATION: GUIDELINES TOWARD A MORE SYSTEMATIC AND TARGETED APPROACH	878
THE CONTENT OF CULTURAL ADAPTATION: SURFACE STRUCTURE, DEEP STRUCTURE, AND CORE COMPONENTS	879
PROCESS MODELS OF CULTURAL ADAPTATION: SYSTEMATIC APPLICATION OF BEST PRACTICES	880
PARENT TRAINING INTERVENTIONS	884
Cultural Adaptations to Enhance Engagement in PT	885
Augmenting Intervention Content to Improve Outcomes for Ethnic Minority Families	886
Cultural Adaptations of Evidence-Based Parent Training Interventions	886
Culturally Grounded Parent Training Interventions	893
Discussion	894
Enhancing Engagement: Remaining Questions	894
Augmented Intervention Content: Paucity of Data	895
Additional Directions for PT Adaptation	895
YOUTH RISK PREVENTION PROGRAMS	896
Cultural Adaptations of Youth Risk Preventive Interventions	898
Culturally Grounded Interventions	901
Discussion	908
Remaining Gaps and Questions	909
PREVENTION OF ANXIETY AND MOOD DISORDERS	909
Cultural Adaptation and the Prevention of Anxiety and Depression	911
Discussion: Limited Evidence	912
HEALTH-FOCUSED INTERVENTIONS	913
The Cases of Obesity and Physical Activity	913
Exemplary Research Programs Integrating Culture and Children's Health	914
CONCLUSIONS AND FUTURE RESEARCH DIRECTIONS	918
Adaptation Effectiveness	918
Cultural Adaptation Process	920
Reach and Engagement	920
Understanding Cultural Components Added to Culturally Adapted Interventions	920
Accommodating Within-Group Diversity	921
REFERENCES	922

WHY CULTURAL ADAPTATION IN DEVELOPMENTAL PSYCHOPATHOLOGY

This third edition of *Developmental Psychopathology* is the first to include a chapter focused on the cultural adaptation of evidence-based interventions (EBIs), a topic that represents a point of convergence for many of the core values and goals of the field. Whereas the overarching goal of developmental psychopathology is to elucidate the interplay among the biological, psychological, and social-contextual aspects of normal and abnormal development, the ultimate goal is to inform intervention strategies to promote resilience and prevent or ameliorate dysfunctional outcomes and disorder across the life span (Cicchetti & Hinshaw, 2002; Cicchetti

& Toth, 1992). Infusion of cultural factors into EBIs is consistent with the fundamental tenet of developmental psychopathology that maturational, experiential, and cultural aspects of development are inseparably coalesced in ontogeny (Cicchetti, 1990). To the extent this produces differences in the existence, patterning, or impact of risk and protective processes for distinct subcultural groups, modifications to intervention programs and protocols may be needed to provide the best means of reducing psychopathology for these groups. Cultural adaptation has been defined as “the systematic modification of an evidence-based intervention or intervention protocol to consider language, culture, and context in such a way that it is compatible with the client’s cultural patterns, meanings, and values”

(Bernal, Jimenez-Chafey, & Domenech Rodríguez, 2009, p. 362). Introduction of this topic in the current volume is well timed with respect to several disciplinary advances in developmental, cultural, clinical, public health, and prevention science and specifically with respect to research on cultural adaptation that has evolved over the past three decades to become a systematic and rigorous field of study in its own right (Barrera, Castro, & Holleran Steiker, 2011; Bernal & Domenech-Rodríguez, 2012; Castro, Barrera, & Holleran Steiker, 2010; Lau, 2006).

This chapter reviews the historical context and challenges that have given rise to this field, the methods and models used to address these challenges, and exemplars of culturally adapted interventions for children and adolescents. Although interventions to address the needs of vulnerable youth exist on a continuum from those that seek primarily to promote health and positive development for whole populations to those targeted at individuals with severe disorders (Weisz, Sandler, Durlak, & Anton, 2005), our presentation of exemplars broadly encompasses the preventive end of the spectrum that addresses risk before it has evolved into debilitating forms of psychopathology, including *universal* prevention approaches that address risk factors in general populations, *selective* preventive interventions that target those segments of the population with greater than normal risk of developing a disorder, and *indicated* approaches that focus on subgroups exhibiting early signs or symptoms of developing a disorder (Mrazek & Haggerty, 1994). The specific goals of the chapter are to: (1) articulate the issues and challenges that have shaped current approaches to cultural adaptation of EBIs; (2) review approaches and models for designing cultural adaptations of EBIs, including frameworks for deciding when, what, and how adaptations should be made; (3) present exemplars of cultural adaptations of EBIs targeting a range of social, emotional, behavioral, and health outcomes; and (4) summarize strengths, limitations, and future directions for the field.

DEFINITION OF CULTURE

Culture is a complex construct that describes the characteristics of a particular group of people, defined by everything from language, religion, cuisine, social habits, music, and arts. A distinct group of people (a tribe, an ethnic group, professional organization, a nation) can be described as “having a culture,” meaning that its members share a collective system of values, beliefs, expectations, and norms, including traditions and customs, as well

as sharing established social networks and standards of conduct that define them as a cultural group (Betancourt & Lopez, 1993). Although culture can be conceptualized at multiple levels to apply to more universal characteristics of the human experience to narrower conceptualization of the lifeways and worldviews of specific cultural groups and subgroups (Muñoz & Mendelson, 2005), the majority of prior research on cultural adaptation equates culture with ethnicity and race; consistent with this trend, our review of exemplars will focus on adaptations for distinct ethnocultural groups. However, as noted by many scholars (e.g., Castro et al., 2010; B. D. M. Wilson & Miller, 2003), the tendency to equate cultural adaptation with adaptation for specific ethnocultural groups may be misguided in some cases because there are numerous other potential sources of mismatch between an EBI and potential consumers that may be more relevant (Reyno & McGrath, 2006). For example, among the many adaptations of parent training (PT) interventions that have been conducted for distinct ethnocultural groups, a few researchers have adapted PT interventions in ways that transcend ethnicity and race to address issues of low socioeconomic status (SES) (e.g., Brotman, Gouley, Klein, Castellanos, & Pine, 2003).

A focus on ethnicity per se is further complicated by considerable heterogeneity that exists within racial and ethnic groups. In the United States, the four major ethnic groups (Latinos/Hispanics, African American, Asian Americans, Native Americans) collectively comprise approximately 37% of the total population, and they range in size from about 1 million for Native Americans to over 52 million for Hispanics/Latinos (U.S. Census Bureau, 2010). Within each of these broad groups, there exist many homogenous subcultural groups based on factors such as nationality, SES, religious background, immigration history, and generational status (Castro, Barrera, & Martinez, 2004). Asian Americans as an ethnic minority group, for example, are quite heterogeneous with over 20 subgroups (e.g., Chinese, Japanese, Filipino, Korean, Vietnamese, Laotian, Cambodian, etc.), each made unique by linguistic, cultural, and sociodemographic backgrounds and immigration histories in the United States (Sue & Morishima, 1982). Diversity among these Asian subgroups has been associated with a variety of differences in mental health service need, utilization, and outcome. Many scholars advocate population segmentation (Balcazar, Castro, & Krull, 1995), the identification and targeting of subgroups based on the intersection of these cultural dimensions, to avoid ethnic gloss (Trimble, 1995) in planning an intervention. However, opposing arguments counter that the adaptation of interventions for increasingly smaller segments

of the population is not feasible given the number of treatments, disorders, developmental stages, and a host of other factors that would generate a matrix impossible to manage (Kazdin, 2008). Moreover, many interventions ultimately will be delivered in multicultural settings and for populations that are changing on cultural dimensions. Alternative models proposed to address this issue have included adaptive approaches (Collins, Murphy, & Bierman, 2004) that provide opportunities for dynamic sizing (Sue, 2006) or tailoring (Zayas, Bellamy, & Proctor, 2012) of culture-specific program elements on a case-by-case basis (McCabe, Yeh, Garland, Lau, & Chavez, 2005; Pina, Villalta, & Zerr, 2009). In most cases, however, clarity regarding one's conceptualization of "culture," including the theoretical rationale or empirical evidence for expecting cultural differences among subcultural groups or between individuals within a distinct subgroup, is often lacking to support such nuanced models.

HISTORICAL CONTEXT OF CULTURAL ADAPTATION RESEARCH

Cultural adaptation takes the position that, to be effective, interventions must be responsive to the cultural practices and worldviews of the subcultural groups for whom these interventions are intended (Resnicow, Soler, Braithwaite, Ahluwalia, & Butler, 2000). Although this view has gained significant traction over the years as the diversity of the U. S. population has expanded exponentially, this point continues to be a matter of significant debate with strong arguments still voiced about the wisdom, necessity, and feasibility of this position (Elliot & Mihalic, 2004; Kazdin, 2008; Ortiz & Del Vecchio, 2013). Objections notwithstanding, cultural adaptation has emerged as an active research area in response to several key developments across disciplines grappling with the role of culture in the provision of behavioral, psychological, and health interventions for diverse populations (Barrera, Castro, Strycker, & Toobert, 2013; Bernal & Domenech Rodríguez, 2012).

Historically, the broader debate about the cultural relevance of psychology to ethnically diverse populations has been a fundamental driving force. The cultural movement in psychology, including the emergence of cross-cultural, multicultural, ethnicity, culture, and race studies over the past three decades, challenged the universality of contemporary theories and raised awareness regarding the lack of representation of ethnic minority populations and the lack of attention given to culturally relevant processes in both basic and intervention research (Betancourt &

Lopez, 1993; Markus & Kitayama, 1991; Sue, 1998). In developmental science, seminal papers highlighted the importance of testing integrative, ecologically, and culturally valid models that could better account for unique cultural experiences of ethnic minority children and families and the role of culture in shaping or altering normal and abnormal development (García Coll, et al., 1996; Szapocznik & Kurtines, 1993). Several scholars in intervention sciences (e.g., McGoldrick & Giordano, 1996; Rogler, Malgady, Constantino, & Blumenthal, 1987; Sue, 1998) argued that the majority of science-based interventions, while characterized as universal, are often validated on and influenced by the values and interests of urban or suburban, White, middle-class samples. Consequently, EBIs may not be relevant for addressing issues confronting differing subgroups, including racial-ethnic minority and rural White populations. Even more serious assertions have been expressed that interventions developed without respect for diversity might be used, intentionally or not, to inculcate values, beliefs, and norms of the dominant culture with the result of devaluing, dismissing, replacing, or even eliminating the culture and language of origin (Comas Díaz, 2006; Gone, 2004). Short of such colonialistic concerns, yet still troubling, was mounting evidence of massive health disparities between European American and American ethnic minorities (U.S. Department of Health and Human Services, 2001) coupled with data showing minority populations lack access, enroll at lower rates, and are more likely to drop out of interventions that might reduce these disparities (Smedley, Stith, & Nelson, 2003). Beginning in the 1990s, these developments and other awareness-raising efforts paved the way to policy changes regarding inclusion of cultural minorities in funded research (National Institutes of Health, 1994) and a growing imperative to provide culturally relevant interventions and services for these populations (U. S. Department of Health and Human Services, 2010). Although ethnic minority groups remain underrepresented in clinical trials and published research (Ortiz & Del Vecchio, 2013; Wendler et al., 2006), particularly in top-tier journals (Hartmann, Kim, Kim, et al., 2014), the number and quality of culturally informed developmental studies and interventions trials has increased steadily since the early 1990s.

A second major influence, beginning in the 1990s, was the emphasis on empirically validated interventions as the gold standard for professional practice (Chambless & Hollon, 1998) and a prerequisite for broad diffusion of prevention and health promotion programs (Flay et al., 2005). On the basis of established criteria for EBIs to receive the label of "efficacious" or "effective," special initiatives

and governing bodies across public health service sectors were charged with the task of creating registries to identify and catalog EBIs ready for public dissemination (e.g., the National Registry of Evidence-Based Prevention Programs and Practices; the Collaborative for Academic, Social, and Emotional Learning, the Blueprints Violence Prevention Initiative, the What Works Clearinghouse). By prioritizing programs with established evidence, the majority of which had been validated initially without representation or attention to cultural diversity, the cultural adaptation agenda was solidified. During the past decade and a half, increasing attention has been given to adapting EBIs for use with ethnic, cultural, and racial minority groups who were not part of the original intervention development process. The ascendancy of EBIs and standards of evidence also resulted in a high premium being placed on fidelity of implementation, often as detailed in program manuals (Addis & Cardemil, 2006), and the belief that program effects are best attained under strict adherence to the intervention's procedures as designed and validated (Elliot & Mihalic, 2004). Thus, the fundamental fidelity-adaptation dilemma, which consists of a "dialectic involving arguments favoring fidelity in the delivery of an EBI as designed versus arguments favoring the need for adaptations to reconcile intervention-consumer mismatches in accord with the needs and preferences of a subcultural group" (Castro et al., 2004, p. 221), became the core challenge for this emerging field.

Significant developments in prevention science over the past three decades have also been important in shaping the cultural adaptation agenda. Because the goal of prevention, even more so than treatment, is to maximize public health access and achieve population-level impact (Spoth et al., 2013), ensuring relevance to diverse populations has long been a central concern. A 2009 report on prevention of child social, emotional, and behavioral disorders summarized an impressive body of evidence demonstrating that, when carefully implemented, prevention and health promotion programs can prevent a wide range of health problems, promote positive development, and achieve economic benefits (National Research Council and Institute of Medicine, 2009). As summarized in this report, numerous randomized trials have shown that these types of interventions have had significant, long-term effects in reducing unhealthy eating, physical inactivity, alcohol, tobacco and other drug abuse, teen pregnancy, school failure, delinquent behavior, and violence as well as other mental, emotional, behavioral, and physical health problems. However, this same report also summarized studies showing that a relatively small percentage of these interventions

actually are implemented by community-based program delivery systems (e.g., public schools, health care facilities, social service agencies). This report thus underscored a significant problem and an important shift in emphasis, already under way in the field, toward greater focus on effectiveness and implementation studies and a push toward broader community diffusion of EBIs. Distinctions between *efficacy* and *effectiveness* are important for the field of cultural adaptation. Whereas efficacy measures how well an intervention works when tested within the controlled conditions under which it was designed, effectiveness measures how well the intervention works in an applied real-world setting (Flay et al., 2005). The shift in emphasis toward evaluating the evidence of effectiveness studies to determine what preventive programs work, for whom, and under what circumstances is greatly needed to move the field forward in launching successful population-based prevention strategies (Brown, Berndt, Brinales, Zong, & Bhagwat, 2000). Further, a rigorous evaluation of broad dissemination and implementation has important implications for the future of research on cultural adaptation. Moving EBIs into widespread, population-level diffusion requires attention to cultural diversity yet also puts a high priority on adaptation to meet the needs of the local community as well as the service settings in which interventions are delivered (Brownson, Colditz, & Proctor, 2012). Such integration of cultural adaptation and implementation science ultimately will be needed to meet the challenge of reducing racial and ethnic disparities in health (Cabassa & Baumann, 2013).

A final, emergent historical force likely to shape future research on cultural adaptation is the spread of evidence-based practice throughout the world. Countries across the globe are increasingly implementing interventions developed in other countries to address their own public health problems. This trend has already resulted in the wide adoption of EBIs and a growing number of controlled trials of imported and culturally adapted EBIs. The government of Norway implemented a countrywide dissemination of the Parent Management Training–Oregon Model (Forgatch, 1994; Patterson, Reid, & Dishion, 1992) and multisystemic therapy (MST; Henggeler, Schoenwald, Borduin, Rowland, & Cunningham, 1998) to target youth behavior problems (Ogden, Amlund-Hagen, Askeland, & Christensen, 2009). The Swedish National Board of Health and Welfare (2011; cited in Ferrer-Wreder, Sundell, & Mansoor, 2012) reported that, in the last 10 years, imported EBIs comprised over 40% of the total number of social and behavioral interventions tested in Sweden. A recent review of randomized trials narrowly focused

on parent training (PT) interventions to reduce family violence in low- to middle-income countries (Knerr, Gardner, & Cluver, 2013) identified trials spanning 9 countries (Brazil, Chile, China, Ethiopia, Iran, Jamaica, Pakistan, South Africa, and Turkey); given the strict criteria for inclusion in this study of randomized trials, the actual practice of international implementation is likely to be even more extensive.

Although the majority of imported EBIs are based on interventions initially developed in the United States, programs such as the Olweus Bullying Program developed in Norway (Schroeder et al., 2012) and the Triple P parenting intervention from Australia (Sanders, 2012) are examples of programs successfully transported to the United States and several other countries. Thus, the proliferation of interventions crossing international boundaries is well under way, and many of the same debates are taking place about the need to adapt interventions for international diffusion, with some experts cautioning against the trend for global dissemination without careful attention to cultural adaptation (Bernal & Scharró-del-Río, 2001) and others challenging the assumption that such adaptations are necessary (Knerr et al., 2013). Noting some examples of imported EBIs that have failed to show expected benefits, Ferrer-Wreder, Sundell, and Mansoori (2012) recently advocated systematic, theory-driven international adaptation studies to give researchers an improved ability to empirically test degrees of intervention adaptation in relation to intervention effectiveness and thereby improve the field's ability to learn from both its successes and failures. Their approach draws on models that primarily have been used for domestic adaptations, yet also highlights additional challenges for the future of cultural adaptation as a global endeavor.

Propelled by these historical forces and attempts to address the underlying tensions and controversies they have engendered, cultural adaptation of EBIs has evolved over the past three decades into a well-defined area of study that now spans multiple disciplinary boundaries. The number of interventions that have been adapted and evaluated has grown and, in the process, systematic, data-driven, consumer-sensitive processes have been established for determining if EBIs should be adapted, how they should be adapted, and the results of those adaptations on engagement and intervention efficacy and effectiveness (Barrera et al., 2013). We summarize these developments in the next section, then follow with a summary of exemplars of culturally adapted EBIs that target a range of child and adolescent psychological, behavioral, and health problems.

THE RATIONALE FOR CULTURAL ADAPTATION: GUIDELINES TOWARD A MORE SYSTEMATIC AND TARGETED APPROACH

In 2006, Lau proposed a compelling framework for identifying circumstances that would justify a cultural adaptation of an EBI. This thoughtful discussion articulated a balanced perspective on cultural adaptation, assuming its importance yet also allowing there are likely to be interventions and conditions for which it is not warranted. For example, prominent examples exist of non-adapted preventive interventions, including some of the most widely disseminated programs in the field, that have demonstrated efficacy in trials with large numbers of ethnic and racial minorities (e.g., MST; Henggeler et al., 1998) or when specifically applied to distinct ethnocultural groups in the United States (e.g., Nurse Family Partnerships; Olds et al., 1997). Lau's Selective and Directed Treatment Adaptation Framework (SDTAF) advocated a theory- and data-driven approach for determining if an EBI should be culturally adapted and, if so, which intervention elements might be altered.

The "selective" dimension of this framework recommended cultural adaptations should be pursued only when evidence suggests that delivering a standard treatment would result in a high probability of failure. Some scholars suggest adaptations should not be made to interventions until the original version has been implemented and problems have arisen (Kumpfer, Pinyuchon, de Melo, & Whiteside, 2008). Sanders (2012) argued for a deployment-focused model of adaptation that privileges the responses from participants who may be in a better position to gauge the acceptability of an intervention, as reliance on the perceptions of professionals serving ethnic minority communities may result in filtering by "cultural gatekeepers" who hold views on cultural acceptability that may differ from intended recipients (Lau, 2012; Morawska et al., 2011). The "directed" part of the Lau's framework addresses the question of what to adapt, indicating that adaptations to content or procedures should be pursued only when evidence suggests such modifications are needed. This framework was additionally important for highlighting that adaptations directed at attracting and engaging distinct ethnocultural groups are equally important as those directed at improving intervention outcomes.

Lau's framework prompted dialogue among cultural adaptation scholars that led to significant changes in the conceptualization of cultural adaptation. In a response to Lau (2006), Barrera and Castro (2006) elaborated on

the conditions in which an adaptation would be justified, which they organized into four categories: (1) ineffective engagement, such as poor recruitment, retention, program attendance, or participation in intervention activities; (2) unique risk or resilience factors underlying the intervention target (e.g., discrimination experiences), or that function differently across cultural groups; (3) unique symptoms of a common disorder; and (4) poor intervention efficacy for a particular subcultural group. Other models of cultural adaptation to be discussed later have since incorporated these suggestions by placing a high priority on initial data gathering as a way to determine whether and what to adapt. However, despite these aspirational frameworks, it has been rare for such systematic decisions to be articulated in the cultural adaptation literature. EBI adaptations often have been conducted without first demonstrating diminished engagement or benefit, and often without theory or evidence about unique risk and resilience factors guiding adaptation efforts.

THE CONTENT OF CULTURAL ADAPTATION: SURFACE STRUCTURE, DEEP STRUCTURE, AND CORE COMPONENTS

Focusing on the content of cultural adaptations, Bernal, Bonilla, and Bellido (1995) were among the first to identify the intervention elements that may require adaptation to increase ecological validity or congruence between the client's ethnic and linguistic context and the cultural properties embedded in an intervention. The Ecological Validity Model identified eight dimensions in which intervention adaptations are likely to be made. Some of these dimensions are more likely to involve adaptations to specific aspects of the intervention itself (e.g., language, metaphors, content, concepts, goals, and context), while others are more relevant to the intervention's implementation support structures (e.g., persons and methods). Resnicow, Soler, Braithwaite, Ahluwalia, and Butler et al. (2000) also focused on the content of cultural adaptation, describing aspects of the program, protocol, or delivery system that might require adaptation; they introduced the influential distinction between *surface structure* and *deep structure* adaptations. In their words, "Surface structure establishes feasibility, whereas deep structure determines program impact" (p. 274). Surface structure adaptations typically involve change in an original EBI's materials or activities that address observable and superficial aspects of a target population's culture, such as language, music, foods, clothing, and other observable aspects.

An intervention's deep structure involves the processes related to the problem of interest and is more likely to involve the theory-based mediators of the intervention that are thought to affect therapeutic change mechanisms or outcomes.

There is wide agreement that surface structure changes are not only permissible but often necessary to convey respect and achieve "cultural attunement," a term coined by Falicov (2009) to refer to additions to EBIs that are intended to boost engagement and retention of subcultural groups in treatment. Providing services in clients' native language, including familiar cultural traditions and teaching methods (e.g., storytelling, games), and utilizing individuals of color as educational facilitators are examples of features that might attract subcultural groups into treatment and increase their sustained participation. Surface structure changes are also important to maximize comprehension (e.g., understandable content that is matched to the linguistic, educational, and/or developmental needs of the consumer group), motivation (e.g., content that is interesting and important to this group), and cultural relevance (content and materials that are applicable to participants' everyday lives), as described by Castro et al. (2004).

Advisability of deep structure change is a matter of debate among adaptation experts. The reluctance to make deep adaptations is related to the belief that there are features of a program that should not be changed because they reflect the core components or the internal change processes of the intervention responsible for program effects (Falicov, 2009). However, the boundaries of deep structure are not always clear and are generally a matter for program developers to define. Evidence from mediation studies that aim to identify core components (MacKinnon, Fairchild, & Fritz, 2007) can provide guidance, but tests of mediation are not always available nor do they include all components that program developers might include in their definition of the program's core components. For example, in the cultural adaptation of the New Beginnings Program for divorcing parents that was conducted to prepare for an effectiveness trial in family courts (Wolchik et al., 2009), mediation analyses identified specific parenting strategies to retain because they accounted for both immediate and sustained program effects (Tein, Sandler, MacKinnon, & Wolchik, 2004; Zhou, Sandler, Millsap, Wolchik, & Dawson-McClure, 2008), but, in addition, program experts identified intervention processes (e.g., home practice assignments and review) as additional aspects of what they considered to be "core components" to be preserved in the adaptation process. Kumpfer et al.

(2008) drew a strong line at making changes to program components, the timing, and the overall structure of the Strengthening Families Program (SFP) when adapting it for international dissemination. They defined these as deep structure adaptations based on experiences and evidence from effectiveness studies of SFP with American ethnic subgroups that showed diminished effects when such adaptations were made (Kumpfer, Alvarado, Smith & Bellamy, 2002).

A second major class of adaptation involves the introduction of novel or augmented intervention content to target risk factors for youth psychopathology that fall outside the foci of the original EBI. In this way, interventionists targeting culturally diverse populations supplement intervention content to address ecological risk and protective factors pertinent to specific cultural communities (e.g., Coard, Wallace, Stevenson, & Brotman, 2004; Lau, Fung, Ho, Kiu, & Gudiño, 2011; C. R. Martinez & Eddy, 2005). Intervention development studies that draw from existing EBIs but infuse additional elements in which culture is a primary consideration sometimes favor the term *culturally informed* (Falicov, 2009) or *culturally tailored* (Bernal & Domenech Rodríguez, 2012) over *cultural adaptation*. Bernal and Domenech Rodríguez (2012) highlighted that incorporation of cultural dimensions such as interdependence, spirituality, relational views of self, and particular worldviews into an intervention can serve goals of supporting the culture and language of origin and the development and maintenance of cultural values, both of which are empowerment strategies. These added elements may not necessarily change core components but can enrich program effects in ways that are specific to the struggles, needs, and strengths of distinct ethnocultural groups. Even the Kumpfer et al. model (2008), which drew a hard line against deep structure change, did not tacitly prohibit augmented content so long as core components are not displaced. When cultural augmentation occurs, clear documentation of added components and inclusion of measures to assess change on these cultural components can facilitate understanding of whether, how, and to what extent they contribute to intervention outcomes, yet these types of analyses are rare in the literature (Castro et al., 2010).

The admonition against changing core components in the process of cultural adaptation is based on the assumption that the theory of change involved is universally relevant and powerful. To the extent that these assumptions are not accurate, more extensive modifications may be needed. Castro et al. (2004) described a continuum among interventions that incorporate cultural elements

that varies between the extremes of making minor surface structure alterations to an original EBI to the complete rejection of the EBI in favor of a novel, culturally grounded approach; the latter falls outside the boundaries of what some scholars would define as “culturally adapted.” Unlike cultural adaptations that begin with an established EBI and then work to integrate cultural elements that increase relevance and fit for subcultural groups, culturally grounded approaches begin with an assessment of need obtained from members of a particular subcultural group who then participate actively in developing the intervention from the ground up (McKay et al., 2004). The latter approach is based in models of community-based participatory research (CBPR; Minkler & Wallerstein, 2003) and is contrasted with the traditional Prevention Research Cycle, which includes a series of stages that begins, instead, with (a) basic research typically conducted in a university setting to identify a disorder’s risk and protective factors, then continues to (b) the design of an intervention that changes modifiable risk and protective factors, (c) empirical trials of the intervention to determine efficacy and to inform intervention modifications, (d) effectiveness trials to assess generalizability to real-world applications, and (e) large-scale intervention dissemination (Mrazek & Haggerty, 1994). Interventions based on the CBPR framework have gained in popularity recently, partly in response to problems that university-based EBIs developed in the Prevention Research Cycle tradition have encountered in going to scale (Barrera et al., 2011). The belief is that programs developed through a CBRP model may be more likely to be adopted and sustained because they are responsive to the cultural and local conditions, organizational structure, and history that influence implementation and because the community has an established investment in the program from the outset (Wandersman, 2003).

PROCESS MODELS OF CULTURAL ADAPTATION: SYSTEMATIC APPLICATION OF BEST PRACTICES

Perhaps the most significant advance in the cultural adaptation literature over the past two decades is the proliferation of conceptual frameworks that describe the processes that can be used in designing, implementing, and evaluating a culturally adapted EBI. To date, more than 13 different models of cultural adaptation spanning both prevention and treatment have been published, almost all within the past 10 years (for reviews, see Castro et al., 2010; Domenech Rodríguez & Bernal, 2012; Ferrer-Wreder et al., 2012). Frameworks relevant to

prevention are listed in Table 17.1, which also identifies similarities and differences in their content, comprehensiveness, and scope. Although developed independently, these frameworks exhibit considerable consensus in their recommendations. For example, most models delineate specific stages or phases in the adaptation process, ranging from three to nine steps across models, but generally covering the four phases recommended by Barrera and Castro's (2006) Heuristic Model (HM): gather information, make preliminary adaptations to the intervention, test the preliminary adaptations, and refine adaptations.

Like most models, the first step outlined by Barrera and Castro (2006) involves a formative stage of information gathering to be used in planning along the lines of Lau's selective and adaptive recommendations. Across models, this phase relies on multiple sources of both quantitative and qualitative information, such as reviews of the literature to understand common and unique risk factors and focus groups to assess perceived positives and negatives of the original EBI. Domenech Rodríguez and Bernal (2012) highlighted the importance of blending a positivist paradigm in which randomized controlled trials (RCTs) are the gold standard for testing the utility of interventions with a pragmatist view that also uses qualitative sources of knowledge to guide decisions. Collaboration with relevant community stakeholders, such as program developers, agency staff, and community members, is another common feature at this stage and throughout the cultural adaptation process. The second phase in Barrera and Castro's model involves program design in which changes are made along the many dimensions described by Bernal et al. (1995) and Resnicow et al. (2000), including to the intervention protocol, interventionists, training, recruitment, and monitoring systems. This stage often involves multiple iterations of cultural adaptation and feedback from community stakeholders, including intervention participants themselves, in an attempt to maximize cultural fit. In the third step, case studies or pilot studies are used to refine recruitment, intervention, and assessment processes, again using both quantitative and qualitative data to evaluate impact of the intervention on engagement, consumer satisfaction, and intervention processes. This stage also may involve multiple iterations of pilot testing and refinement to prepare for the fourth stage, in which the final intervention package is prepared for a more robust test of the culturally adapted intervention.

Although most of the models listed in Table 17.1 cover similar phases and recommendations, there are important differences between them. Two of the models, the Adaptation for International Transport of Strengthening

Families Program and the International Implementation of Multisystemic Therapy, were developed in relation to the adaptation of specific programs, whereas others offer generic or general guidelines for the field. The models also vary in terms of their intended application in different countries versus a primary focus on domestic ethnocultural groups. Finally, whereas some models are focused on the endpoint of preparing for evaluation of program efficacy, others are oriented toward effectiveness and dissemination. For example, the model for international implementation of MST was developed to support cross-national adaptation and dissemination of multisystemic therapy (Schoenwald, 2008). MST, a systems-oriented family therapy designed to benefit troubled adolescents who have exhibited conduct and/or substance abuse problems, has been implemented in 34 states in the United States and 15 countries to date (Henggeler & Schaeffer, 2010). In contrast to cultural adaptation models focused primarily on program content and design, the international implementation of MST model attends to implementation issues, such as the selection of appropriate clinical staff, training materials and procedures, and the clinical delivery system. Kumpfer and colleagues (2008) delineated a model based on their experience disseminating the SFP around the globe. Their 9-step model (see Table 17.1) begins with a needs assessment and selection of an appropriate EBI and ends with a model that is ready for cross-national dissemination.

The Planned Intervention Adaptation (PIA) protocol is unique because it is a generic model for cross-national adaptation of interventions and is geared toward the design of effectiveness and dissemination trials of imported EBIs (Sundell & Ferrer-Wreder, & Fraser, 2014). In PIA, cultural groups are conceptualized as nested, recognizing that distinct subgroups (e.g., Swedes with a Latino family background) are nested within a larger cultural context (e.g., a Swedish community), thus attempting to address the challenge of attaining cultural sensitivity to the unique conditions produced by these intersecting aspects of culture. PIA begins with formative research (phase I), followed by an effectiveness trial (phase II), which provides evidence that may indicate a need for another round of adaptation or a dissemination study. At the conclusion of PIA's phase II, the feasibility of a dissemination trial is evaluated by program developers and new stakeholders. PIA also specifies a time period for the initial adaptation stages and a sampling strategy for formative studies. Phase I of PIA is recommended as a roughly 1.5- to 2-year period when intervention developers and stakeholders (agency staff, potential consumers) agree to collaborate on five preliminary steps that are summarized in a table and detailed

TABLE 17.1 Models of Cultural Adaptation by Date of First Publication

		Content of Adaptation	Steps/Phases of Adaptation	Stage of Research	Cultural Groups Targeted ¹	Generic vs. Specific Application
Ecological Validity Model (EVM)	Bernal et al. (1995)	Language, persons, metaphors, content, concepts, goals, methods, context	None specified	General use in research and practice contexts	Broadly applicable (domestic, international)	Generic
Cultural Sensitivity Model	Resnicow, Baranowski, Ahluwalia, & Braithwaite (1999); Resnicow et al. (2000)	Distinguished 2 adaptation types: Surface structure Deep structure	None specified Recommended use of both qualitative and quantitative data sources	Broadly focused on preparing for implementation	Broadly applicable (domestic)	Substance use and public health programs
Cultural Adaptation Process Model	Domenech Rodríguez & Wieling (2004)	Emphasized EVM dimensions (Bernal et al., 1995)	Set the stage Initial adaptation Adaptation iterations	General use in research and practice contexts	Broadly applicable (domestic)	Generic
Selective and Directed Treatment Adaptation Framework	Lau (2006)	Identified 2 goals: Enhance engagement Enhance program effects	None specified Decisions about whether to adapt and what to adapt should be guided by evidence	Not specified	Broadly applicable (domestic)	Generic
Heuristic Framework for the Cultural Adaptation of Interventions	Barrera & Castro (2006); Castro et al. (2010)	Adaptations address: Ineffective engagement Unique risk or resilience factors Unique symptoms Lack of efficacy	Gather information Preliminary adaptations Pilot test adaptations Refine adaptations	Adaptation of EBI for efficacy trial with distinct group	Broadly applicable (domestic)	Generic
Culturally Specific Prevention Model	Whitbeck (2006)	Culture-specific risk and protective factors must be considered. Target group knowledge equally important as scientific evidence.	Review existing evidence Target group reviews existing evidence Cultural translation of risk and protective factors Identify unique risk and protective factors Conduct efficacy trial	Model starts with local community defining the problem and extends to efficacy trial	American Indian communities	Generic
Adaptation for International Transport of Strengthening Families Program	Kumpfer et al. (2008)	Fidelity requires cultural adaptation but not modification of program components, timings, or overall structure	Needs assessment Select EBI to adapt Implement original EBI with minor modifications Select and train staff Implement with fidelity Adapt continuously through pilot tests Revise to enhance engagement Conduct evaluation Disseminate results	Dissemination	International	Strengthening Families Program
International Implementation of Multisystemic Therapy	Schoenwald et al. (2008)	Adaptations relevant to: implementation processes, clinical staff, training materials and procedures, clinical service delivery	Pre-implementation (assess needs and stakeholders; understand legal and social standards; determine staffing and consultants) Implementation (translate and “tweak” intervention; adapt training protocols) Service delivery (consider service duration; time and caseload standards; on-call systems; cultural advisors)	Dissemination	International	Multisystemic therapy

TABLE 17.1 (Continued)

		Content of Adaptation	Steps/Phases of Adaptation	Stage of Research	Cultural Groups Targeted ¹	Generic vs. Specific Application
ADAPT-ITT	Wingood & DiClemente (2008); Latham et al. (2010)	Adapt program to maintain fidelity to core components	Assess needs of target group and agency capacity to implement. Decide on EBI. Target group input. Adapt program. Topical experts review first draft. Integrate feedback in second draft. Train implementation staff. Pilot test, revise, evaluate.	Dissemination	Broadly applicable (domestic, international)	HIV interventions
Planned Intervention Adaptation Protocol	Ferrer-Wreder et al. (2012)	Use formative research to evaluate relevance of risk and protective processes and the need for surface and deep adaptations	Formative research. Effectiveness trial. Dissemination.	Effectiveness and dissemination	International	Generic

¹Indicates cultural groups targeted and whether the model has been applied domestically, internationally, or both.

in text (e.g., language translations, tests of translated materials, focus group checks on cultural appropriateness of intervention materials and activities). These preliminary steps help shape the adapted intervention. In phase II of PIA, the authors recommend a three-arm effectiveness study in which a minimally adapted intervention (just surface structure changes, such as language translation), a fully adapted intervention (both surface and deep structure changes), and a control condition are tested. They also recommended the inclusion of appropriate measures and data analytic procedures to identify possible mediators and moderators of intervention effects. Unfortunately, based on Sundell et al.'s (2014) article, it is not certain that the PIA framework actually has been used in international adaptations of interventions.

Common to all models is the expectation that cultural adaptations will be evaluated, either in randomized efficacy or in effectiveness studies. Several questions are critical to address, including effects on engagement and program outcomes and test of mediators and moderators. In efficacy trials, effects on program mediators are useful to identify core components of the intervention to guide future dissemination and effectiveness studies in other contexts or with other populations (e.g., Gonzales et al., 2012; Lau et al., 2011; Murry et al., 2013). Studies of culturally adapted interventions also offer important opportunities to test whether cultural elements included in

the adaptation contribute to intervention efficacy (Castro et al., 2010; B. D. M. Wilson & Miller, 2003). Evaluation of cultural adaptation in the context of effectiveness studies can test whether cultural adaptations impact program uptake and fidelity, the best methods to train interventionists and monitor delivery, and program effects on distinct ethnocultural groups receiving services in natural delivery settings, and whether interventionists are more or less likely to adhere to the intervention protocol when delivering culturally adapted interventions (Spath et al., 2013). Test of moderators can identify subgroups that may show differential response in terms of engagement or outcomes based on dimensions such as acculturation or adversity (Gonzales et al., 2011; Lau et al., 2011), thus advancing understanding of the need for population segmentation while also identifying subgroups most likely to benefit and those that may need further adaptations. Perhaps the most important question rarely addressed when evaluating culturally adapted interventions is whether the adapted EBI is more effective or offers other added benefits (e.g., increased likelihood of adoption in the community, higher enrollment rates) when compared with the original EBI (Castro et al., 2010; Resnicow et al., 2000).

Although it remains a largely unaddressed empirical question, it is possible that the extent of optimal changes to an EBI also will depend on the nature of the EBI to be adapted. For example, adaptations may be more important

for interventions that target family processes than for those delivered in schools, or when they target specific mediators or outcomes. In the next sections, we present a qualitative summary of culturally adapted interventions to prevent a range of child and adolescent emotional, behavioral, and health problems that have been evaluated in RCTs. Our presentation features exemplars and is not intended as an exhaustive review. Because the nature and impact of cultural adaptation may vary depending on the nature of the intervention, we focused our review on four distinct domains. Two broad domains—adaptations of PT interventions and adaptations of sexual risk and drug preventive interventions—were chosen because they represent the most active areas for past research on cultural adaptation. We also include a summary of adapted prevention interventions targeting child and adolescent internalizing outcomes that are less common due to a relative lack of focus in prevention on problems such as anxiety and depression. Our final domain includes health-focused preventive interventions, a newly emerging and important area of emphasis for reducing minority health disparities. Our purposes in presenting this collection of exemplars is to examine: (a) the extent to which child and adolescent interventions have been adapted for distinct ethnocultural groups, including for which groups and for what types of preventive interventions; (b) the processes used and the nature of adaptations that have resulted; (c) and the evidence base for addressing key question about efficacy or effectiveness, including tests of underlying cultural dimensions that account for program effects, subgroup differences in intervention responsiveness, and the most salient question: Do cultural adaptations make a noticeable impact on outcomes? In each section, we first describe the prevention domain, the types of EBIs in each domain that have been culturally adapted and the evidence base to support them, and the empirical or epidemiological basis to suggest that cultural adaptations of these EBIs may be relevant to the public health needs of targeted groups.

PARENT TRAINING INTERVENTIONS

Maladaptive parent-child interactions have been found to precede the onset of or to maintain multiple forms of child psychopathology across both externalizing and internalizing spectrums. Even in conditions not primarily driven by environmental socialization factors, research has documented the ameliorative effects of instantiating specific types of parent-child interactions in improving outcomes of various forms of psychopathology (e.g.,

joint attention and symbolic play for autism spectrum disorders). As such, family-based interventions often form the foundation of prevention and intervention programs across a range of psychopathology targets. In particular, EBIs for conduct disorder and related externalizing behavior problems in youth almost uniformly include PT as the sole or first line intervention, or as a major component of a multisystemic intervention. The purpose of PT is to prevent and decrease disruptive behavior and increase prosocial behaviors in children. The efficacy of PT is well established with robust evidence of clinically significant improvements in child conduct problems (Eyberg, Nelson, & Boggs, 2008; Sheldrick, Kendall, & Heimberg, 2001). Treatment aims to improve on parents' skills and increase the consistent use of those skills. PT is rooted in social learning theory (Bandura, 1977) with specific strategies designed to interrupt cyclical coercive interactions during which aversive behaviors emitted by parents and children are negatively reinforced (Patterson, Chamberlain, & Reid, 1982). Common elements across PT interventions serve to reinforce appropriate child behavior (e.g., positive attention, praise, tangible rewards) while removing reinforcement or punishing negative child behavior (e.g., differential attention, time-out, response cost) (Garland, Hawley, Brookman-Frazee, & Hurlburt, 2008). Parents are also instructed to structure expectations for appropriate behavior through limit setting, routines and rules, and effective commands. Consistent with the Hanf (1969) two-stage model, PT interventions typically begin with instruction, modeling, and rehearsal of positive reinforcement skills to increase prosocial behavior before introducing elements intended to address child misconduct. The rationale is that a more positive parent-child relationship sets the stage for more child compliance and better uptake of positive discipline strategies. Yet some experimental evidence refutes this assumption (Eisenstadt, Eyeberg, McNeil, Newcomb, & Funderburk, 1993).

The application of interventions prescribing fundamental changes in child-rearing practices at the population level has been questioned. Parenting sensibilities and routines are embedded within a system of meanings, shaped by cultural context and values regarding the socialization of the moral child and the expected roles of adult caregivers and children within family structures (Harkness & Super, 1996; Weisner, 2002). Discourse on the application of PT with ethnic minority families has enumerated potential cultural barriers to engagement among parents whose own socialization experiences fall outside middle-class European American heritage (e.g., Barker, Cook, & Borrego, 2010; Forehand & Kotchick, 1996; Lau, 2006). Given

that parent-child relations and discipline practices are the proximal targets of change, many have cautioned that cultural barriers may threaten the generalizability of PT effects. Wide cultural variation in parenting practices and values across ethnic groups may influence receptivity to PT elements, perhaps accounting for increased attrition (e.g., Kazdin & Whitley, 2003), low enrollment (e.g., August, Lee, Bloomquist, Realmuto, & Hektner, 2003; Cunningham et al., 2000) and reduced participation (e.g., Orrell-Valente, Pinderhughes, Valente, & Laird, 1999) among ethnic minorities and immigrants.

Central questions concerning the application of PT with culturally diverse families emerge from sociopolitical and moral perspectives as well as from epistemological and developmental science perspectives. First, widespread dissemination of PT might be viewed as a form of cultural hegemony or colonialism to the extent that it promotes assimilation to European American child rearing as superior (Dionne, Davis, Sheeber, & Madrigal, 2009). Second, the dissemination of PT interventions that were largely developed with middle-class European American families to culturally diverse communities reflects the assumption that treatment effects will generalize across distinct social and cultural family contexts (Bernal & Scharró-del-Río, 2001). Yet some research suggests diversity in relations among parenting behaviors, family processes, and child developmental outcomes (e.g., Deater-Deckard & Dodge, 1997; Halgunseth, Ispa, & Rudy, 2006; C. Ho, Bluestein, & Jenkins, 2008; Lansford et al., 2005).

For these reasons, PT has come under scrutiny as a class of interventions with effects that may be subject to boundary conditions. Accordingly, PT has been well represented among efforts in the cultural adaptation of behavioral interventions. Two distinct themes have emerged in endeavors to enhance the outcomes of PT with ethnic minority and international populations (Lau, 2006). Beyond surface adaptations and basic translation, the range of adaptations that fundamentally alter intervention content or process can be organized into those that work toward enhancing engagement versus those that focus on tailoring intervention content to augment skills training.

Cultural Adaptations to Enhance Engagement in PT

Limited engagement in PT represents a major threat to effective dissemination and implementation in disadvantaged and ethnic minority families. Clinical observations from single-case design experiments and controlled trials of PT with ethnic minority families previously suggested notable cultural distance between skills typically taught in

PT and values governing parent-child relations in ethnic minority families (Borrego, Ibanez, Spendlove, & Pemberton, 2007; J. Ho, Yeh, McCabe, & Lau, 2012; T. Ho et al., 1999; Lau, Fung, L. Ho, Liu, & Gudiño, 2011; Lau, Fung, & Yung, 2010). For example, because of a cultural priority to avoid losing face and the need for constant correction of improper behavior, Chinese parents objected to ignoring misbehavior based on principles of differential attention (Lieh-Mak, Lee, & Luk, 1984). Likewise, Mexican American parents have rated differential attention as less acceptable than other PT practices (Borrego et al., 2007), with some parents objecting because ignoring may harm the parent-child relationship (Pemberton & Borrego, 2007). While praise is often rated as among the most acceptable PT techniques (J. Ho et al., 2012) some interventionists have noted that praise is problematic for Chinese parents owing to beliefs that it results in a lack of humility and complacency over persistence (Crisante & Ng, 2003; T. Ho et al., 1999).

Given that many of the concerns about PT for culturally diverse populations have focused on issues of acceptability, and most disparities in PT controlled trial research relate to differential enrollment and premature attrition, PT interventions targeting ethnic minority families have been culturally adapted by attending to barriers to engagement (e.g., Dillman Carpentier et al., 2007; Matos, Bauermeister, & Bernal, 2009; McCabe & Yeh, 2009). Engagement-focused adaptations encompass surface structure modifications (Resnicow et al., 2000) intended to sufficiently engage the target community so that the intervention can be delivered with fidelity to the so-called deep structure. Fidelity to deep structure maintains the key elements of the intervention's theory of change, or logic model specifying the theoretical relations between the treatment procedures (e.g., teaching parents to praise children for appropriate behavior), mechanisms of change (e.g., alteration of reinforcement schedules), and ultimate outcomes (e.g., increased child compliance) (McKleroy et al., 2006). Most such adaptations focus on adopting a collaborative stance by soliciting and defusing concerns about specific skill elements in PT, reframing techniques to optimize fit with parents' values, and using motivational strategies to stoke parental investment in agreed-on treatment goals.

There is, of course, within-group heterogeneity in attitudes toward PT interventions, with these attitudes covarying with cultural values in distinct ways for different groups. Among Chinese American parents, low levels of acculturation and endorsement of traditional values concerning strict discipline and shaming are associated

with lower perceived acceptability of PT (J. Ho et al., 2012). Yet endorsement of traditional Latino values that emphasize strong orientation and commitment to family, *familismo*, among Mexican Americans has been associated with positive attitudes toward PT (Pemberton & Borrego, 2007). Notably, at least one culturally adapted intervention has adopted an adaptive assessment-driven strategy to tailor the delivery of PT to promote optimal participation and acceptance among parents with a range of baseline attitudes (McCabe, Yeh, Garland, et al., 2005).

Augmenting Intervention Content to Improve Outcomes for Ethnic Minority Families

A second major class of adaptations to PT involves the introduction of novel intervention content to target risk factors for youth psychopathology or family adjustment difficulties that fall outside the original foci of PT. In this way, interventionists targeting culturally diverse families have supplemented PT content to address ecological risk and protective factors associated with child behavior problems and parenting competence pertinent to specific cultural communities (e.g., Coard et al., 2004; Lau et al., 2011; C. R. Martinez & Eddy, 2005). As with other efforts to augment PT, these adaptations address ancillary family stressors that can heighten child vulnerability and interfere with parent skill acquisition (G. E. Miller & Prinz, 1990). These culturally adapted PT protocols have addressed risk processes such as parental acculturative stress, family-level acculturation dissonance, and experiences of discrimination (Lau, 2006). In this way, these adaptations revise the intervention's deep structure or theory of change within a causal model informed by studies of risk and protective factors that highlight potential mediators of change that can improve target outcomes in the population of interest.

Cultural Adaptations of Evidence-Based Parent Training Interventions

Parent-Child Interaction Therapy

Parent-Child Interaction Therapy (PCIT) is designed to treat 2- to 7-year-olds with conduct problems (Herschell, Calzada, Eyberg, & McNeil, 2002). Parents are taught skills to establish a nurturing and secure relationship while reinforcing prosocial behavior and decreasing negative behavior. It is conducted in the context of parent-child play interactions during which the therapist actively coaches the parent in the use of skills, often unobtrusively from behind an observation window. PCIT is assessment driven and terminates when parents demonstrate

mastery of the skills and their child's behavior is within the normal range on a validated scale of child conduct problems (Eyberg Child Behavior Inventory; Eyberg & Pincus, 1999). Treatment begins with the child-directed interaction (CDI) phase designed to create a mutually rewarding parent-child relationship. Toward this end, parents are taught to praise, imitate, and describe the child's appropriate play behavior, reflect appropriate child speech, ignore inappropriate behavior, and allow their child to lead the activity. Parents are also taught to refrain from criticism, commands, and questions. The goal of the second, parent-directed interaction (PDI), phase is for parents to decrease problematic behavior by directing child activity using clear, positively stated, direct commands and consistent consequences (e.g., praise for compliance, time-out for noncompliance).

PCIT has been used with diverse ethnic minority families in the United States with positive results. For example, the effects of PCIT in reducing recidivism in child maltreatment appeared to generalize across racial/ethnic groups including non-Hispanic Whites, Latinos, and African Americans, with no apparent intervention by ethnicity interactions (Chaffin et al., 2004). However, some data reveal disparities in rates of engagement and outcomes in PCIT. Fernandez, Butler, and Eyberg (2011) reported a 60% rate of attrition from PCIT in a sample of low-income African American families of preschoolers with disruptive behavior disorders. Further, only half of the treatment completers showed clinically significant change in child conduct problems, and there was no significant effect of treatment on parenting stress, as has been shown for non-Hispanic White families in PCIT. Previous trials have revealed associations between low SES and dropout from PCIT (Fernandez & Eyberg, 2009); motivational enhancement appears to prevent such attrition among ethnically diverse and disadvantaged families (Chaffin et al., 2009).

Multiple investigators have culturally adapted PCIT using varying approaches to increase responsiveness to the needs of Latino families. Borrego et al. (2006) first demonstrated the feasibility of delivering PCIT in Spanish. Matos, Torres, Santiago, Jurado, and Rodríguez (2006) went further to adapt PCIT for Puerto Rican families of young children with hyperactivity. This also represented primarily a language-based adaptation that included translation of written materials and modification of examples to reflect the daily experiences and idiomatic expressions of Puerto Rican families. Unstructured additional time was allotted to the beginning of each PCIT session to establish and maintain therapeutic rapport through personal interactions and to discuss any issues that could be affecting

treatment progress. Matos et al. (2009) reported large effect sizes (Cohen's $d = 1.37$ to 2.04) in their waitlist controlled trial, with 65% to 75% of treated families evincing clinically significant decreases in conduct problems.

McCabe and colleagues (2005) described a more comprehensive adaptation of PCIT for Mexican American families that adopted a tailoring approach to address barriers to engagement. The adaptation design involved the integration of studies of treatment barriers among Latino families and original qualitative research. The authors noted that cultural barriers for Mexican American parents included feared negative consequences of treatment (e.g., breached confidentiality), disapproval from family members, and doubts of effectiveness or alternately unrealistic expectations of rapid improvement. Specific adaptations of PCIT were informed by focus groups and interviews with Mexican American parents of youth with disruptive disorders. PCIT was described, and participants were invited to comment on its features and suggest improvements. The considerable variability in attitudes (e.g., some Latino parents viewed time-out as not punitive enough, while others felt it was too severe) suggested that adaptations must flexibly accommodate variation in values and acculturation. Yet several clear themes emerged. Given concerns about blame and stigma, mothers preferred the program to be framed as educational rather than therapeutic. Mothers highlighted the need for clear explanations for how activities, such as play, can reduce behavior problems. They also requested additional intervention materials, such as father testimonials, and augmented content, such as discussion of the negative consequences of paternal absence, to promote involvement of fathers and extended family.

Based on these data and consultation with an expert panel and the PCIT developer, McCabe et al. (2005) developed *Guiando A Ninos Activos (GANA)* to include an intensive engagement protocol beginning with the initial phone contact to problem solve barriers to attendance, increase investment in treatment, and create a plan to enlist fathers and other caregivers. GANA is framed as an educational program, offered by a teacher (*maestro*). CDI teaches communication skills (*Ejercicios de Comunicacion*) while PDI teaches consistent discipline (*Disciplina Consistente*). The use of assessment-based tailoring is unique. Given the heterogeneity in attitudes, McCabe et al. designed an initial assessment to direct the implementation of treatment elements. GANA teachers receive a computer-generated report that contains guidance for addressing potential issues (e.g., unrealistic expectations for treatment) and highlights elements that may require reframing to increase acceptability. For example, among

parents favoring strict discipline, time-out is framed as a punishment (*silla de castigo*) whereas for parents who disavow punitive methods, time-out is framed as a time for calming and reflection (*silla de pensativa*).

In a controlled trial that was noteworthy as one of few that compared the culturally adapted intervention with the original EBI, McCabe and Yeh (2009) randomized 58 Mexican American parents with a child with a disruptive disorder to receive standard PCIT, GANA, or treatment as usual (TAU). McCabe and Yeh found that both PCIT and GANA outperformed TAU at posttreatment, and although the differences between PCIT and GANA were not statistically different, the mean effect sizes associated with the GANA-PCIT planned contrast were encouraging (Cohen's $\delta = .28$ for conduct symptoms, $\delta = .42$ for parenting). At follow-up, McCabe, Yeh, Lau, and Argote (2012) reported that GANA outcomes were superior to TAU on most parent-report measures, while PCIT did not maintain an advantage over TAU. Thus, the efficacy and maintenance of gains in GANA have been supported, and there is some limited evidence to suggest that GANA may yield greater longer-term benefit than standard PCIT. However, although GANA adaptations were explicitly designed to enhance engagement, there was no effect of condition on premature termination.

The Incredible Years

The Incredible Years (IY) group intervention is effective in reducing child behavior problems, improving parents' child management skills, and decreasing parenting stress (Reid, Webster-Stratton, & Baydar, 2004). Videotape modeling is an important feature of IY, to both demonstrate techniques and stimulate discussion. As with other PT programs, skill rehearsal is emphasized through role play and homework. IY incorporates many features that may promote parental engagement by introducing skills in a collaborative rather than a didactic manner. As each skill is introduced, parents discuss the benefits and barriers to using the skill, opening up issues of perceived acceptability and fit with parents' values. The group leader encourages investment by highlighting how the benefits of the skill fit with the parents' own goals. The group leader uses Socratic questioning to allow parents to identify the principles underlying effective parenting skills. Treatment manuals orient group leaders to common concerns about each PT skill (e.g., concerns that praise will "spoil" children, worries that time-out is not severe enough). Group process leverages peer-to-peer teaching and experiential learning, such as praising and rewarding parents in session as they are instructed to do with their children. Thus, IY builds in multiple, effective

strategies for enhancing engagement for culturally diverse families. Data from prevention trials fielded in Head Start preschools have shown that IY achieves parity in parenting and child behavior outcomes and parent satisfaction across non-Hispanic White, Latino, African American, and Asian American families, with the number of ethnic differences in attrition and outcomes not exceeding that expected by chance (Reid, Webster-Stratton, & Beauchaine, 2001).

However, some independent investigations of IY suggest that treatment gains among African American and Latino families may not be lasting. For example, Gross et al. (2003) showed improvements in parenting behaviors from IY among low-income, largely African American (57%) and Latino (29%) parents of toddlers in Chicago. However, there were no lasting reductions in child behavior problems based on parent reports or observer ratings at the 1-year follow-up. Similarly, Brotman et al. (2003) found significant initial improvements in parent responsiveness and parent-reported behavior problems following IY in a sample of 20 African American and 10 Latino preschoolers, but these effects were not maintained over 6 months.

IY has been augmented in order to address risk factors that appear relevant to the development of disrupted family relations and child behavior problems in ethnic minority populations. Lau et al. (2011) adapted IY to address risk factors for maladaptive discipline in high-risk immigrant Chinese families in southern California. Lau (2010) found that children's school problems are strongly associated with harsh physical discipline among Chinese immigrant families and that parent-child acculturation conflicts were related to physical discipline when parents held traditional values about firm parental authority. In a waitlist controlled trial including 54 families, PT was augmented with three skill components to reduce contextual stress to allow parents to follow through with effective PT strategies. First, cognitive restructuring was added to help parents to control upsetting thoughts about children's bids for autonomy and school problems that lead to ineffective parenting. Second, psychoeducation and communication skills training helped parents reframe and resolve acculturation-related conflicts. Parents were taught active listening skills and problem-solving steps to elicit the child's perspective and communicate their own concerns effectively. Third, to prevent punitive responses to school problems, parents were taught to increase positive involvement in children's schooling by showing interest in their child's learning, structuring a homework routine, and limiting (television and computer) screen time.

Results of the trial revealed good retention with 83% of families completing the 14-week intervention. Relative to

parents in the delayed-treatment condition, parents who received the intervention responded with lower levels of negative discipline and child internalizing and externalizing problems. Posttreatment effect sizes were in the medium to large range for child behavior problems, with good durability of results at 6-month follow-up. Mediation analyses revealed that decreases in negative discipline accounted for improvements in child externalizing problems. Treatment effects were moderated by baseline parenting stress, such that more distressed parents benefited less from treatment. In contrast, an analysis of outcomes of IY among 514 families across six randomized trials did not find baseline parenting stress to moderate outcomes (Beauchaine, Webster-Stratton, & Reid, 2005). Moreover, despite the inclusion of augmented PT content to address stress in immigrant parent-child relations, intervention effects on parenting stress were not observed at posttreatment. This finding suggests that the cognitive restructuring and communication training components were not sufficient to produce effects among the most distressed immigrant parents in the sample.

ParentCorps is a culturally adapted PT protocol, strongly informed by IY, to prevent child behavior problems and promote school readiness in low-income African American and Latino families of preschoolers in New York (Brotman et al., 2008). Intervention content and process aim to be universally relevant and engaging to all families yet sufficiently intensive and tailored for families residing in disadvantaged, urban communities with risk factors such as community violence, overcrowded schools, and diverse immigrant populations. Guided by the assumption that culture informs all aspects of caregiver psychology (i.e., expected family roles, appraisals of child behavior, future aspirations, and appropriate parenting), ParentCorps focuses explicitly on cultural values, beliefs, and norms and encourages parents toward treatment goals that are culturally relevant. For example, session plans outline discussions concerning values related to obedience and respect for elders and the influence of parents' own childhood experiences on their beliefs about parenting. As in IY, videotape modeling introduces each skill in the context of the daily lives and interactions of three ethnic minority families. Group leaders individualize the use of the skill for each family's unique context and use group activities and role-plays and troubleshoot homework. Parents also receive a ParentCorps item each session to spur uptake of the assigned skill (e.g., a timer for time-outs).

A randomized trial of ParentCorps including 171 families of 4-year-olds in a university preschool program in a large New York public school district revealed effect sizes

in the .50 range for parenting practices and child behavior problems (Brotman et al., 2011). Recruitment results in this trial were good for a universal prevention design: 71% of parents in the intervention schools attended at least one session. However, in terms of retention only 54% attended at least five of 13 sessions.

Indian Family Wellness Program

In their cultural approach to implementing IY, Dionne et al. (2009) designed an initial motivation phase to engage American Indian parents in PT by acknowledging the historical role of colonization in the disruption of traditional parenting and resultant social illnesses. This phase is a three-session motivational interview designed to help parents recognize: (a) how intergenerational transmission of parenting knowledge has been disrupted by colonization, (b) the impact of historical trauma and ongoing injustices on parenting, (c) strengths across generations despite this history, and (d) their responsibility for summoning strengths and adapting their parenting to protect their children from social illnesses. In the intervention phase, IY was delivered with fidelity, but at the beginning of each session, the interventionists drew connections between the PT skill and Central American Indian traditions, beliefs, and values. For example, when teaching alternatives to corporal punishment, parents were reminded that this tactic was introduced by European colonists and in boarding schools and was not consistent with Indian teachings that children were gifts who were never to be hurt. In a controlled trial, 49 American Indian families received the motivational interview and were then randomized to receive the culturally adapted IY or no treatment. Results at posttreatment revealed significant reductions in ineffective parenting and child externalizing and internalizing problems in the treatment group relative to controls.

The Chicago Parenting Program

The Chicago Parenting Program (CPP) is another intervention building on IY that capitalizes on the strengths of videotaped modeling followed by group discussion to elicit principles of effective behavior management and a collaborative approach to engage parents. The CPP was designed in partnership with a parent advisory council of African American and Latino parents from different Chicago neighborhoods. The council advised the program developers on challenges they faced as parents, situations that should be portrayed in videotape modeling, and how best to depict PT skills to be congruent with their cultural values and lifestyle (Gross, Garvey, Julion, & Fogg,

2007). Videos were shot in homes and in public places and depicted African American, Latino, and White families in situations that are familiar, relevant, and stressful. In a departure from IY, the CPP intervention did not proscribe spanking, given the views in some communities. The 11 sessions covered child-centered time; family routines and traditions; praise and encouragement; rewards for reducing challenging behavior; setting clear limits and following through; consequences for misbehavior; strategies of ignore, distract, and time-out; stress management; and problem-solving skills. A booster session is offered 2 months later to help parents maintain gains.

The efficacy of CPP was demonstrated in a controlled trial of families from Chicago area preschools serving low-income families. Families were randomized to receive CPP or assessment only. The study had a recruitment rate of 34%, with average attendance of 4.3 (39%) of 11 sessions. Results indicated that CPP reduced corporal punishment, commands, and observed child behavior problems up to 1-year postintervention. Dosage effects were clear; mean effect sizes for observed child behavior problems, parent behaviors, and parenting self-efficacy increased by 50% when dose was included in the model. Parents of children with more behavior problems at baseline attended more sessions, suggesting that CPP was differentially reaching families who need it most.

Parent Management Training—Oregon Model

Parent Management Training—Oregon Model (PMTO) was derived from social interaction learning theory and grounded in direct observation of family interactions with a critical focus on reducing coercive family processes (Forgatch & Patterson, 2010). Child behavior problems are thought to begin with small negative behaviors that escalate over time (e.g., noncompliance, to shouting, to tantrums) through repeated escape conditioning sequences. Common parental responses to child aversive behaviors inadvertently increase their frequency. For example, a mother finds her son watching television instead of doing homework. The mother scolds, the boy complains and claims he has no homework, the mother relents and changes the topic, and the boy stops arguing. The interaction has increased the likelihood of the boy protesting; he has learned how to turn off the mother's aversive behavior. In ensuing interactions, he may escalate his protest behavior. This is conceptualized as a coercive cycle. Blame is not directed toward individuals in the family but to the interaction between them. Five core parenting practices are identified as mechanisms for change: (1) skill encouragement (promoting prosocial behavior through

teaching and contingent positive reinforcement), (2) limit setting (decreasing deviant behavior through appropriate and contingent use of mild sanctions), (3) positive involvement (ways to invest time and plan activities with children), (4) problem solving (negotiating disagreement, establishing rules and consequences for rule violations), and (5) monitoring (protecting children from involvement in risk behaviors and affiliation with deviant peers).

Cultural adaptations of the PMTO model have been described in the literature. Domenech Rodríguez, Baumann, and Schwartz (2011) described the process by which they developed, *Criando con Amor: Promoviendo Armonía y Superación (CAPAS)* for Latino parents, using their Cultural Adaptation Process Model mentioned on Table 17.1. CAPAS is a product of systematic adaptations to PMTO guided by Bernal's (1995) ecological validity model and a three-stage process employing community engagement and deployment focused iterative adaptation. CAPAS incorporated adaptations to content and process to promote active parental engagement in PMTO. Adaptations focused on six areas:

1. Careful language translation, with simplified terminology and greater reliance on visual aids;
2. Inclusion of Latino-specific *dichos*, proverbs and sayings in Spanish, to provide familiar metaphors for concepts;
3. Incorporating Latino values (e.g., *respeto*, *buena educación*, *valerse por si mismo*) to frame treatment goals, discussion of parent versus child values reflecting emphasis on Latino versus American orientations;
4. Framing PT as promoting family harmony and success;
5. Inviting extended family members; and
6. Attending to contextual challenges in parenting (e.g., fear of children calling child protective services, deportation).

These adaptations preserve the core PMTO elements and the putative mechanisms of change—increasing skill encouragement, positive involvement, problem solving, limit setting, and monitoring.

Consistent with the CAPAS example, C. R. Martinez and Eddy (2005) attended to many of the same cultural and contextual considerations in their adaptation of PMTO called *Nuestras Familias: Andando Entre Culturas (Our Families: Moving Between Cultures)* for Latino families. *Nuestras Familias* was developed over a 2-year period through a collaborative effort between the Oregon Social Learning Center project team, Centro Latino/Americano, other community partners, and senior

Latino family intervention researchers around the country. During the intervention development phase, five Latino family interventionists from the community were provided training in basic Oregon Social Learning Center PMT. After training (which included applying PMT principles to real and fictional cases), project staff, together with community experts and trained interventionists, began the process of adapting the existing PMT intervention model. In addition to keeping the core PMTO components and framing the intervention with consonant cultural values, Martinez and Eddy augmented intervention content with two modules that focused explicitly on bridging two cultures. This augmentation focused on differential acculturation between Latino parents and youth as a specific risk factor for youth maladjustment and behavior problems. Differential acculturation is related to increases in family stress and decreases in effective parenting practices, and these processes mediated the association between discrepant acculturation and adolescent substance risk (C. R. Martinez, 2006). *Nuestras Familias* targets “acculturation gaps” that can increase stress and disrupt effective parenting. C. R. Martinez and Eddy (2005) reasoned that for PT to be maximally effective in this context, it must address how parents manage family relations in the context of differential acculturation. In addition to the core PMTO content, the program includes four components to provide psychoeducation about differential family acculturation and empower parents to bridge acculturation gaps through the use of effective parenting practices associated with standard PMTO. In a randomized trial of *Nuestras Familias*, parents in the intervention condition reported an increase in overall parenting effectiveness and a decrease in child behavior problems compared to the no-intervention control. Furthermore, some evidence indicated that the intervention resulted in stronger effects for families with U. S.-born youth, for whom family adjustment difficulties associated with differential acculturation may be more prominent (Martinez & Eddy, 2005).

PMTO was implemented nationwide in Norway in 1999 with an effectiveness trial indicating positive intervention effects. Yet the sample enrolled in the trial was predominantly native Norwegian families, with few ethnic minorities represented (Ogden & Hagen, 2008). Developmental research was conducted in 2005 to 2007 to investigate strategies for recruiting immigrant Somali and Pakistani families and to determine what types of cultural adaptations may be necessary. The investigators studied the cost-effectiveness of three strategies for recruiting ethnic minority mothers to participate: (a) referrals from human service professionals from public agencies, (b) open

information meetings in community settings, and (c) social network referrals from research staff link workers. All three recruitment strategies yielded families eligible with regard to risk for child conduct problems. However, the information meetings were the most cost-effective strategy, and the highest proportion of the sample was recruited this way. The intervention groups were not mixed gender owing to Islamic religious norms requiring gender segregation. Groups were ethnically homogeneous and led by a Norwegian therapist assisted by trained bilingual assistants (i.e., link workers) who provided direct translation in Somali/Urdu. Link workers also served as “culture-builders” to strengthen relationships between therapists and mothers. In terms of intervention content, there was an expanded focus on certain topics relevant in the context of the immigrant communities targeted, including large sibling groups, and emphasis on cultivating control over emotions.

Findings from intent-to-treat analysis of a randomized waitlist controlled trial of 96 Somali and Pakistani mothers revealed that PMTO was effective in increasing positive parenting and decreasing harsh discipline and child conduct problems, according to maternal report (Bjørknes & Manger, 2013). The largest effect sizes were found among mothers who attended more than 50% of sessions. However, teacher reports showed no significant intervention effects on conduct problems and social competence. This study provides support for contextual and cultural adaptation in order to secure recruitment to and participant attendance in available interventions.

The Triple P Positive Parenting Program

Triple P spans the prevention-intervention continuum and aims to prevent behavioral, emotional, and developmental problems in children by enhancing the knowledge, skills, and confidence of parents (Sanders, 1999). Within a public health model, Triple P incorporates five levels of intervention on a tiered continuum of increasing intensity and narrowing reach. Level 1 includes media-based psychoeducation campaigns targeting whole communities. Level 2 is Selective Triple P delivered in 1 to 2 brief consultations during well child pediatric care. Level 3 Triple P is a four-session consultation model for use in primary care. Level 4 Standard Triple P is an eight- to ten-session parenting skills program that can be delivered in a group, individual, or self-help format. Level 5 is Enhanced Triple P, which provides adjunctive interventions for families in which parenting concerns are situated in the context of other problems, such as marital conflict and parental depression. Most trials with diverse populations have

examined Level 4 Standard Triple P. Meta-analyses have concluded that Triple P has a positive effect on children's behavior and adjustment, with evidence being strongest in the toddler, preschool, and elementary-school age groups (de Graaf, Speetjens, Smit, de Wolff, & Tavecchio, 2008). Effect sizes range widely from small to large positive effects, which is not surprising that Triple P contains multiple levels of intensity across prevention and treatment trials.

Data from controlled trials, focus groups, and survey methods support the acceptability and effectiveness of parenting strategies used in Triple P with samples of parents from Australia, the United States, New Zealand, Japan, Singapore, Hong Kong, Iran, Scotland, England, Ireland, Sweden, Belgium, the Netherlands, Germany, Turkey, Switzerland, South Africa, and Panama (Sanders, 2012). The International Triple P Research Network coordinates a network of scientists and facilitates communication about research activity around the world involving Triple P. The network has created a data repository for outcome studies and coordinates an international conference for researchers, practitioners, and policymakers in a different country annually. Training in Triple P encourages practitioners to work collaboratively with parents and to be responsive to parent values, preferences, and context while preserving core elements. These needs for tailoring can be met by adapting examples used to illustrate teaching points and by customizing homework assignments.

Perhaps given the foregoing discussion of workforce training in flexible and tailored delivery, there have been few cultural adaptations of Triple P. A notable exception is Turner, Richard, and Sanders's (2007) Group Triple P tailored for Australian Indigenous families. The authors conceded that mainstream parenting programs find it difficult to recruit and engage Indigenous parents. In collaboration with Indigenous workers over 5 years, Triple P was tailored to reduce barriers for Indigenous families and to develop resources. The team assumed that the core principles of positive parenting would transcend cultures but the goals and target behaviors, practical implementation of skills, and ways of sharing information would require adaptation. Modifications were made to the language, images, and examples used to illustrate skills in the production of tailored videos, workbooks, and presentation aids. In addition, more time was devoted in sessions to sharing personal stories, developing trust, discussing the social and political context for parenting, and slowing the pace of presentation.

In a waitlist controlled trial of the adapted Triple P program with 51 Indigenous families, treatment parents

reported decreased child behavior problems and less frequent dysfunctional parenting compared to waitlist families. Effects were primarily maintained at 6-month follow-up. Qualitative data revealed high satisfaction with the intervention. Yet engagement was lower than in previous trials of Group Triple P, with only 60.9% families commencing groups. Moreover, only 28% of families randomized to waitlist subsequently attended treatment, suggesting the importance of engaging families upon first contact, reducing barriers to care, and providing care whenever family circumstances permit.

The Effective Black Parenting Program

The Effective Black Parenting Program was an early example of a culturally adapted PT intervention developed by Myers and colleagues (1992) and adapted from the Confident Parenting Program (Eimers & Aitchison, 1977). Specific skills training is provided in describing and monitoring child behavior, specific labeled praise is used for appropriate behavior, special incentives are used for respectful child behavior, and mild social disapproval, active ignoring, and time-outs are used for misbehavior. Parents are instructed in two general approaches to parenting within which these skills can be utilized. First, parents are taught a Family Rule Guideline Strategy, in which they examine the rules they have for their children and the reasoning behind their rules. This orients parents toward developing rules that have an articulated rationale. Next, the Thinking Parents Approach encourages parents to think before they act and to entertain multiple possible attributions for their child's behavior, including stage of development. The intervention is introduced in a discussion that asks parents to contrast the goals of traditional Black discipline (i.e., discipline is achieved through punishment and spanking to instill obedience) versus modern Black self-discipline (i.e., internalizing standards of effective behavior). Discussion aims to situate coercive parenting practices as once protective in African American families but perhaps no longer as adaptive in empowering African American children to strive for achievement and social change. Additional discussion is devoted to instilling pride in Blackness, avoiding racial self-disparagement, and promoting strategies for coping with racism. The program is delivered by African American professionals and organized within a framework called the Pyramid of Success for Black Children.

In a quasi-experimental design, seven schools with high African American enrollment were assigned to the intervention condition (five schools) or to the control condition (two schools). All African American families with children

in the first or second grade were invited to participate in fifteen 3-hour sessions of the Effective Black Parenting Program. Across two cohorts, 109 parents were enrolled in treatment and compared over time to 64 parents in control schools. Results suggested that the intervention produced improvements in some target outcomes, including parental rejection, family relationship quality, and child behavior outcomes. One-year follow-up data from the first cohort suggested maintained effects in parental rejection and selected child behavior problems, although some return to coercive parenting practices was noted. Ultimately, the authors noted that the intensive short-term Effective Black Parenting Program was promising but that results were modest and circumscribed, even when the program was delivered in a culturally sensitive manner.

The Black Parenting Strengths and Strategies

The Black Parenting Strengths and Strategies (BPSS) program (Coard, Wallace, Stevenson & Brotman, 2004; Coard, Foy-Watson, Zimmer & Wallace, 2007) is a cultural adaptation of the Parenting the Strong-Willed Child program (Forehand & Long, 2002). BPSS teaches behavioral management skills (i.e., attending, rewarding and ignoring, giving effective directions, and how to use time-out appropriately) while also mobilizing racial socialization practices found to be protective among African American families. Coard's work is guided by the premise that for African American youth, developmental competence requires the ability to make sense of and cope with prejudice and discrimination. Coard and colleagues (2004, 2007) reviewed research linking parent racial socialization practices to child outcomes including a positive racial identity, academic functioning, self-esteem, positive parent-child interactions, and decreased depression and anger. Coard et al. (2004) conducted qualitative interviews to analyze socialization messages related to racial achievement, preparation for bias, racial equality, and racial pride emphasized by mothers of young African American children living in low-income neighborhoods. The narratives provided the basis for program components for BPSS, a PT intervention augmented to mobilize culturally specific protective factors. Content adaptations included lessons in six areas of how to:

1. Deal with common in-group events (e.g., overt and subtle incidences of racism, prejudice, and discrimination);
2. Problem-solve negative peer interactions (e.g., social exclusion because of skin color);
3. Promote achievement despite barriers (e.g., advocating for a child when low expectations are evident);

4. Reflect on one's own experiences as an African American woman or man;
5. Discuss race-related content in a developmentally appropriate manner; and
6. Access culturally affirming resources for children.

The five delivery-related adaptations included:

1. Using African American language expression and the African American–perspective use of “we”;
2. Emphasis on African American values about collective responsibility, cooperation, and interdependence;
3. Use of African proverbs, sayings and affirmations, poems, quotes, symbols, and pledges;
4. Use of prayer, role-playing, storytelling, extended family participation, and humor; and
5. Use of a setting and motifs representative of the population.

Throughout the adaptation process, consultation with the original developers helped to ensure that key elements of the standard PT program were maintained and not diluted when content was augmented.

In a waitlist controlled trial of BPSS, 32 parents from urban, low-income neighborhoods were randomized to immediate or delayed 12-week treatment. Attendance and retention were strong with 88% of parents being retained in the study, with participants attending 85% of sessions on average and all participants attending at least six sessions. Results indicated significant effects of treatment on six of eight parent-reported outcomes assessed posttreatment. Relative to controls, intervention parents used significantly more racial socialization strategies, positive parenting practices, and less harsh discipline. Parent-reported externalizing problems increased in the control group but decreased in the immediate treatment group. Effect sizes were large for positive parenting, harsh discipline, racial socialization, and conduct problems.

Culturally Grounded Parent Training Interventions

The African Migrant Parenting Program

The African Migrant Parenting Program is a PT intervention for immigrant and refugee families from Africa who resettle in high-income countries, such as Australia (Renhazo & Vignjevic, 2011). The program aims to enhance effective parenting and relationship skills in order to help parents to raise their children confidently and understand their children's needs across

developmental stages in the new cultural, social, and educational environment. Resettlement challenges encountered by refugee and migrant families include: adaptation to new work demands; needs for housing and schooling; changing family structure, roles, and responsibilities; establishing social networks; and learning English (Renhazo, 2002). Previously accepted practices, such as reliance on sibling care and corporal punishment, may no longer be viable. Differential family acculturation and acculturative family distancing are pertinent for these families who are transitioning from collectivistic cultures into high-income countries with social environments promoting individualism. In this context, strain on family relations and parenting are pressing (Renhazo, Swinburn, & McCabe, 2008).

The African Migrant Parenting Program consists of eight 2-hour sessions cofacilitated by African parenting educators and experienced family counselors; three 45-minute home visits by African parenting educators were used to monitor barriers to learning and skill enactment. The sessions were modeled on the Parent Skills module of the Parenting in a New Culture Guide developed for Australian Samoan parents. Session activities incorporate computer slide presentations, discussions, mini case studies, and videotape modeling. Sessions were devoted to a variety of issues including improved family relations, family stress reduction, and even family finances and legal issues.

In an open trial, 44 families receiving counseling for parenting issues at a social service agency were invited to participate. All agreed to participate, and 88.6% of families completed treatment despite a very low level of literacy and multiple barriers to treatment (e.g., housing, transportation, legal problems). Pre-post analyses suggested change on key dimensions: appropriate expectations of children, increased empathy toward children's needs, and attitudes toward corporal punishment. Treatment dose was associated with degree of change on developmental expectations and views on corporal punishment. However, key outcomes of parenting behavior and child adjustment were not examined.

The Exploring Together Program

The Exploring Together Program is a preventive intervention adapted to reduce disparities in the developmental profiles of Australian Aboriginal children relative to the general Australian population. Disparities include the higher prevalence of emotional and behavioral disorders and lower performance of Aboriginal children on national tests for literacy and numeracy. The program combines

elements of PT and children's social skills training for parents of children ages 4 to 6 years old (Littlefield et al., 2005). Children attend with one parent for 10 weekly 2-hour sessions. The first hour consists of an interactive group with all parents and children facilitated by four group leaders and focuses on parent-child interaction. Components of group activity involve turn taking, talk and play based on themes that complement content of the other groups. Then the dyads split into separate groups for parents and children for an hour. Facilitated by two group leaders, parent groups discuss observations of children's behavior, triggers for problem behaviors in the context of family interactions, and factors affecting parenting. Children play together in a separate group, with a focus on social skills and the learning of rules and consequences.

Collaboration with Aboriginal team members resulted in adaptations that drew on songs and storytelling in Aboriginal languages. Drawing activities were used to promote discussion of families, networks, and relationships and to tap into cultural symbols and idioms. These practices aimed to encourage parents to talk about the place of parent and child in the family and values of shared responsibility for children among caregivers in extended family networks. The focus was on identifying strengths and sources of difficulty in each family.

The delivery of interventions for this population faces formidable challenges in geographical dispersal and the linguistic and cultural diversity of Aboriginal communities. An open trial evaluation study was mounted for both Aboriginal and non-Aboriginal children in urban Darwin and for Aboriginal children in three communities of the Tiwi Islands, near Darwin (G. Robinson, Tyler, Jones, Silburn, & Zubrick, 2012). A wide variety of implementation strategies was used to promote reach of the intervention. Children were referred from schools; parents were invited by teachers and then visited at home when they expressed interest. Community leaders explained the program at public meetings, spoke to parents privately, and sought referrals from schools. Group leaders were mature community members with backgrounds in health, child care, or education.

Following the intensive and multipronged approach to recruitment, over 250 children were referred over 3 years, yielding an evaluation sample of 225. However, only 110 (48.9%) commenced the intervention, and only 86 (38.2%) attended at least half of the sessions. The lowest commencement and completion rates were observed among urban Aboriginal families (14% commenced and 11.5% attended at least half of the sessions). Despite the high attrition, outcomes suggested significant reduction in problem

behavior and parental distress at posttreatment. However, improvement was significantly greater for non-Aboriginal families, with the least change observed among urban Aboriginal families. Despite highly flexible strategies of engagement and cultural adaptation of the intervention, results suggested the adapted program still had limited reach and impact on urban Indigenous families.

Discussion

In summary, the research base on cultural adaptation in the area of parenting interventions is relatively mature. The four most frequently cited PT interventions—Triple P, IY, PCIT, and Parent Management Training—Oregon model—have all been subject to cultural adaptations. In addition, other extensive adaptations and culturally grounded PT interventions have been developed and evaluated. We now have ample data suggesting that culturally adapted PT can be efficacious, yielding positive effects in the improvement of parenting behaviors and the prevention and treatment of externalizing as well as internalizing child behavior problems. Thus, the effects of PT interventions appear robust to cultural adaptations.

Enhancing Engagement: Remaining Questions

Most PT adaptations tested to date target changes to promote engagement of underserved or culturally diverse groups. However, as yet there is no conclusive evidence that these engagement-focused adaptations improve the attendance and retention of ethnocultural groups for whom disparities in engagement in standard PT have been observed. Benchmarking studies may be called for since it typically would be neither pragmatic nor ethical to randomize to conditions in which PT is not optimized to engage targeted ethnocultural groups (Cardemil, 2010). Only a single small controlled trial has addressed this question, with negative results on engagement outcomes. McCabe and her colleagues (2008, 2012) found culturally adapted PCIT (GANA) did not outperform standard PCIT in terms of promoting attendance or retention even though adaptations focused on engagement through enhanced treatment orientation, framing PCIT skills based on a pretreatment values assessment, or situating PT as educational rather than therapeutic.

Moreover, although positive engagement results were found in many studies, our review noted examples of trials of PT adapted for engagement that yielded disappointing recruitment, commencement, and completion rates. For example, the CPP adaptation of IY yielded a

recruitment of 37% of African American and Latino parents in low-income preschools, and the average family attended only 39% of sessions (Gross et al., 2009). Once recruited into the trial, there was high attrition from the Indigenous adaptation of Triple P for Australian Aboriginal families with nearly 40% of families failing to attend a single session in the immediate treatment group (Turner et al., 2007). Even lower commencement and attendance rates were found in the Exploring Together PT program also targeting Aboriginal families in urban and rural Australia (G. Robinson et al., 2012). These and other findings demonstrate that even complex and concerted efforts to connect with families and communities are at times insufficient for reducing barriers to PT.

Indeed, engagement may remain an impediment to the public health impact of PT. It can also be argued that modifications to intervention content and process do little to address problems with low recruitment and intervention commencement rates. Parents' perceptions of acceptability of PT when they do arrive at treatment may help to explain low retention, which may be a smaller part of the implementation problem. Triple P developer, Sanders (2012), argues that the single major threat to the reach of PT stems from the fact that they are commonly restricted to small numbers of vulnerable parents with problems. As such, the interventions are seen as for "failed" parents, such as those involved in child protection or mental health systems, and participation is stigmatized. Thus, he argues that a public health approach offering PT to all can result in greater engagement at the population level and that cultural adaptations to intervention content need not be emphasized.

Reliance on self-referred parents and mass media outreach may result in self-selected samples who are motivated and able to attend and whose parenting may more closely resemble PT skill targets (G. Robinson et al., 2012). These recruitment strategies may screen out disadvantaged groups with high need. When Triple P moved beyond universal outreach to target a disadvantaged underserved group with an Indigenous adaptation of the program, substantially lower commencement rates were observed than in general community trials (Turner et al., 2007). Again, the challenge may be getting parents into the room rather than engaging them once they arrive.

Augmented Intervention Content: Paucity of Data

Relatively few PT interventions have been augmented to address the etiological context of parenting dysfunction or child behavior problems in diverse families. Frameworks

guiding cultural adaptation often mention the need to address unique etiological factors but provide little guidance on the development of this content. The few examples of augmented PT interventions have addressed: (a) the protective factor of racial socialization in African American families (Coard et al., 2004; Myers et al., 1992); (b) the school achievement context of ineffective discipline in Asian American immigrant families (Lau et al., 2011); and (c) the risks associated with differential acculturation between immigrant parents and their children (Lau et al., 2011; C. R. Martinez & Eddy, 2005; Renhazo & Vignjevic, 2011). Although the absolute efficacy of these augmented PT interventions has been established, dismantling and process studies are needed to understand whether the additional content remediates the risk processes targeted and whether those changes facilitate ultimate outcomes on child psychopathology.

Additional Directions for PT Adaptation

Although some models of cultural adaptation stress that changes to the pacing, sequence, and length of standard PT should not be made (Kumpfer et al., 2008), these prescriptions appear to guard against unwanted omission of core content. However, some observations from controlled trials suggest that pacing and dosing of PT might be an explicit focus of cultural adaptation to help bridge the cultural distance between cultural traditions of diverse families and prescribed changes to parent-child interactions in PT. In general, controlled trials have revealed dose-response effects associated with standard and culturally adapted PT (e.g., Baydar, Reid, & Webster-Stratton, 2003; Gross et al., 2007; Renhazo et al., 2008). In trials of PCIT, which uses mastery-based criteria to determine length of treatment, trials have shown that the number of sessions required for ethnic minority families to reach performance criteria on CDI and PDI skills exceeded that found in trials of predominantly European American families (Fernandez et al., 2011). For example, in Matos et al.'s (2009) trial, the investigators adapted their mastery criteria using different criterion goals that they felt could be more easily attained by the Puerto Rican families in their sample, and hoped to limit CDI to eight sessions and PDI to nine sessions (doses that are still above published averages), but many families required additional sessions above this limit. McCabe and colleagues (2008) likewise found that Mexican American parents required four to five—more sessions than is typical to complete PCIT. Differences in treatment duration may reflect the need for more behavioral rehearsal of practices that are

unfamiliar and culturally foreign. These problems may be pronounced for immigrant and ethnic minority parents who, given their diverse upbringing, are unacquainted with skills like child-directed play, labeled praise, differential attention, and time-out. Qualitative data from group leaders in Lau et al.'s (2011) trial of IY adapted for Chinese Americans suggested that slowing the pace of skill lessons and increasing the dosage of behavioral rehearsal may be a promising adaptation to achieve meaningful and enduring changes in parenting in immigrant families. This is, of course, not a culturally specific proscription for enhancing PT and can be said of parents from across cultural groups. The issue of dosing may be vital in implementation of PT with diverse parents domestically and internationally, and yet this consideration has generally not entered discussions of cultural adaptation.

YOUTH RISK PREVENTION PROGRAMS

National data from the 2011 Youth Risk Behavior Survey revealed that youths in grades 9 to 12 participated in multiple risky behaviors. During the 30 days before the survey, 38.7% had drunk alcohol, 23.1% had used marijuana, 47.4% of the students had ever had sexual intercourse, 33.7% had had sexual intercourse during the 3 months before the survey, and 15.3% had had sexual intercourse with four or more people during their life. Further, 18.1% of high school students had smoked cigarettes and 7.7% had used smokeless tobacco (Eaton et al., 2012). Studies also consistently find that youth's risky behaviors co-occur; specifically, drinking, smoking, and risky sexual behaviors might have occurred in the same situation because youth lost control of themselves. The co-occurrence of substance use and risky sexual practices has been associated with increased prevalence of HIV/AIDS among youth. In 2009, youth ages 13 to 29 accounted for 39% of the new cases of human immunodeficiency virus (HIV) infection, but they represented only 21% of the American population. African American youths disproportionately represented 65% of the new youth HIV infection cases (Centers for Disease Control, 2011).

Further, the report published by the Centers for Disease Control and Prevention in 2011 showed that, among all new cases of HIV infection in the United States, more than 45% are African American adolescents and adults and more than 20% are Hispanic/Latino adolescents and adults. African American females are particularly at higher risk, representing 63% of all new female cases of HIV infection. More than 50% of the new cases were

caused by male-to-male sexual contact (38.2% are African American, 23.5% are Hispanic/Latino), with other new cases due to heterosexual contact (65.3% are African American, 18.1% are Hispanic/Latino) and injection drug use (47% are African American, 25% are Hispanic/Latino).

One unambiguous implication of these statistics is that African Americans and Hispanics are confronted with high risk of HIV infection. Consequently, they are in more urgent need of access to HIV/AIDS knowledge, skills to negotiate and refuse, and support from family and peer leaders. A second clear implication is that unprotected sexual intercourse and drug use are the primary causes of HIV/AIDS infection among minority youth. Whenever these behaviors co-occur, the risk grows tremendously. Multiple sexual partners worsen the situation as it increases the risk of exposure for youth who do not have a direct relationship with an HIV/AIDS-infected partner (Santelli, Lindberg, Abma, McNeely, & Resnick, 2000). It has been suggested that the consequences of poverty and racism are the most salient predictors of risk-taking behaviors. Education, urbanization, social norms, values, and life goals influence academic aspirations and future orientation. Lack of opportunities for success may cause these youth to seek nontraditional methods of achieving adult status, such as early sexual initiation and the initiation and escalation of substance and drug use (Bakken & Winter, 2002; Stanton et al., 1994). African American and Hispanic youth have traditionally reported high rates of these risk behaviors (Eaton et al., 2008). For example, among adolescents surveyed in 2007, only 67.3% of African Americans and 61.5% of Hispanics reported using condoms during the last time they had sexual intercourse. Additionally, greater percentages of African American (27.6%) and Hispanic (17.3%) youth have had four or more sex partners than non-Hispanic Whites (11.5%). Studies also show that Hispanic and African American youth report higher rates of lifetime drug use than do non-Hispanic White youth (Johnston, Bachman, O'Malley, & Schulenberg, 2008). For example, 25.0% of Hispanic, 19.4% of Black, and 17.9% of non-Hispanic White eighth graders report using drugs in their lifetime (Johnston, O'Malley, Bachman, & Schulenberg, 2009).

Racial-ethnic substance and drug use comparison data that include American Indians and Alaska Natives, however, also show that these youth initiate tobacco, alcohol, and other drugs earlier, at higher rates and with more severe health, social, and economic consequences than adolescents of other ethnic-racial groups in the United States. The prevalence rate of smokeless tobacco use is

especially high among American Indian and Alaska Native children, adolescents, and young women (Schinke et al., 1986). Further, rates of cigarette smoking and heavy use of cigarettes—consumption of at least a pack of cigarettes a day for the last 30 days—are dramatically higher among American Indian and Alaska Natives, exceeding rates for every other ethnic-racial group (Schinke, Moncher, Holden, Botvin, & Orlandi, 1989). Drunkenness and use of substances, including alcohol and nearly all other drugs, is higher among American Indian youth compared to youth in other racial-ethnic groups (Nelson & Nazbah, 2011; Young, 1988). Collectively, these findings underscore the importance of identifying effective ways to prevent engagement in high risk behaviors that compromise the health and well-being of all youth and especially that of racial-ethnic minorities.

A plethora of youth risk preventive intervention programs have been developed and have taken many forms, including school-based prevention programs, mass media campaigns, youth clubs and activities designed as alternatives to substance use, family interventions, and comprehensive, community-wide prevention strategies (Sloboda & Bukoski, 2003). Earlier school-based substance abuse prevention approaches included either resistance-skills training, alone or in combination with competence-enhancement or life-skills training (Botvin, Baker, Dusenbury, Tortu, & Botvin, 1990), that were based on social influence models of risk taking and Bandura's (1977) social learning theory. In this approach, drug use and abuse were conceptualized as socially learned and functional behavior resulting from an interplay between social (e.g., modeling, imitation, reinforcement, pro-drug norms) and individual (e.g. personality) factors. In contrast to drug-specific resistance strategies, competence-enhancement approaches aim to teach a repertoire of skills that adolescents can use to deal with many of the challenges confronting them in their everyday lives, not just drug and alcohol use, such as decision-making and problem-solving skills, cognitive skills for resisting interpersonal and media influences, skills for enhancing self-esteem (goal-setting and self-directed behavior-change techniques), adaptive coping strategies for dealing with stress and anxiety, general social skills (conversational skills and skills for forming new friendships), and assertiveness. These skills are typically taught using a combination of proven cognitive-behavioral methods, such as instruction and demonstration, group feedback and reinforcement, behavioral rehearsal, and extended practice through behavioral homework assignments.

Although resistance and life skills training remain central features, prevention programming expanded over the years to better address the co-occurrence of multiple risk-taking behaviors (e.g., alcohol and drug use, sexual risk taking, HIV) and a widening range of risk and protective processes identified as possible targets (Hawkins, Catalano, & Miller, 1992). Programming enhancements also were informed by developmental and social learning research and theories suggesting that adolescents face pressures to engage (or motivations to avoid) in risk taking from multiple sources and transactions outside of school (e.g., peers, parents, other adults in the community, mass media, community policies) and by evidence that single-channel programs (e.g., school-based drug prevention) may not be powerful enough to produce lasting prevention effects against these broader influences (Dryfoos, 1993). Alternative models, particularly family interventions to be used as the primary approach (Dishion, Kavanagh, Schneiger, Nelson, & Kaufman, 2002) or in combination with youth-focused programs (Spath, Redmond, & Shin, 2001) were developed to strengthen family relationships, bonding, communication about difficult topics (e.g., sex, drugs) and parent management practices and to engage parents in supporting skills taught in youth programs (Anderson et al., 1999). Comprehensive prevention strategies that combine school-based interventions with those affecting the family, social institutions, and the larger community (e.g., Pentz et al., 1989) were recommended in the National Institute of Drug Abuse (1997) publication titled *Preventing Drug Abuse Among Children and Adolescents: A Research-Based Guide* that is recognized for marking the beginning of the evidence-based drug abuse prevention movement. This report identified the tailoring of interventions to address risks specific to ethnic minority populations as one of 14 principles guiding the field.

In this section we assemble the scattered collection of culturally adapted youth risk prevention programs that have been evaluated to date, all but one published since the 1997 NIDA report, and the effectiveness of such programs for preventing and reducing risky sexual behavior and alcohol and substance use among racial/ethnic minority adolescents. As in the prior section on PT, our review is organized to distinguish culturally adapted programs, including those that might be more aptly described as “culturally tailored” (Bernal & Domenech-Rodríguez, 2012), from culturally grounded programs in which community members were involved throughout the initial program inception, design and evaluation, although we acknowledge again that these are not precise distinctions.

Cultural Adaptations of Youth Risk Preventive Interventions

Aban Aya Youth Project

Aban Aya Youth Project is a 4-year-long culturally tailored intervention that is designed to prevent multiple risky behaviors among urban African American youth (Flay, Graumlich, Segawa, Burns, & Holliday, 2004). The program screened the public schools in metropolitan Chicago and built the initial sampling pool based on these standards: enrollment greater than 500; more than 80% of the students are African American and less than 10% are Latino or Hispanic; not on probation or slated for reorganization; not a special designed school; and moderate mobility. The 155 schools selected were then stratified into four quartiles of risk (based on school report card data, such as enrollment, attendance and truancy, mobility, family income, and achievement scores), and 12 of them were sampled (nine inner city schools and three near-suburban schools).

The participants were 153 fifth graders in the 1994–1995 school year. Using schools as units, the students were randomly assigned to one of the three conditions: The social development curriculum (SDC), the school/community intervention (SCI), and the control condition. Students in the SDC arm received 16 to 21 lessons each year, which taught them cognitive-behavioral skills to avoid drug use, risky sexual behaviors, and other risky behaviors. SDC is the Afrocentric arm intervention and was based on *Nguzo Saba* (a Swahili term that means the Seven Principles), which emphasizes the importance of promoting “unity, self-determination, and responsibility” (Flay et al., 2004, p. 377) in youth risk prevention approaches (Flay et al., 2004, p. 377). Instructional methods were based on oral traditions of African American heritage including storytelling and the use of proverbs, history, and literature. Finally, the project assigned homework that aimed at encouraging parental involvement and communicative skill building. The authors did not indicate whether community members and parents were involved in the curriculum development and design of SDC.

The SCI arm is a comprehensive intervention designed to enhance the SDC classroom intervention by providing community empowerment sections that include programs for parents, school staff, and youth. This component of the intervention also has an overarching community element to promote positive connections among parents, schools, local businesses, and agencies. Parent support sought to increase skills to boost children’s learning in school and to enhance parent-child communication. Students in the

control condition received similar amounts of classes as those assigned to the SDC with topics focused on promoting healthy behaviors, such as nutrition, physical activities, and general health care. The intervention classes were taught by university-based health educators. Data were collected at multiple points for each condition. The pretest was done in the fall of 1994, and the follow-up posttest was done in the fall of 1995. Further follow-up tests were done annually until 1998.

Results revealed intervention effects differed for boys and girls. For boys, both the intervention arms (SDC and SCI) led to a salient reduction on drug use and recent sexual intercourse and improved condom use compared to boys in the control condition. It was worth noting that boys exposed to the SCI intervention arm fared better than those assigned to the SDC arm. Neither intervention condition was shown to be effective for girls (Flay et al., 2004). Lack of program effectiveness for girls was attributed to low baseline risk compared to males, as boys demonstrated higher baseline levels than girls on each of the targeted outcomes (Flay et al., 2004).

Adult Identity Mentoring

Adult Identity Mentoring, or Project AIM, is a school-based intervention program designed to reduce risky sexual behaviors among African American youth. The primary goals of Project AIM include delaying initiation of first sexual intercourse, promoting abstinence, and increasing condom use (Clark et al., 2005). The conceptual framework was based on the theoretical tenets of the Possible Selves program (Oyserman & Markus, 1990), which posits that people’s current lives are motivated by their mental images of their future selves. Such mental images can be enhanced by a balance between positive images (desired selves) and negative images (selves to be avoided). AIM was delivered in small groups in which role modeling was used as a teaching strategy to highlight negative connections between desired future life and risky behaviors. Project AIM targeted seventh-grade, primarily low-income African American families in a suburban town near a southeastern metropolitan area. African American students who were previously enrolled in a required 9-week health education class were enrolled in the program. A total of 248 students completed pre-intervention assessment, and 6 were dropped due to missing data; 211 completed posttest, 19 weeks after baseline, and 156 completed 1-year follow-up assessment.

In terms of implementation, AIM consisted of 10 sessions delivered over the course of 6 consecutive weeks during the school year and offered in 20 health education

classes. Of the 20 classes, 11 were randomly assigned to receive AIM intervention, and 9 were assigned to receive standard health education curriculum. AIM sessions were conducted by trained African American college students, while the health education classes were taught by the regular classroom instructor. Besides inclusion of African American educational facilitators, Project AIM also had a session that focused on the importance of “legacy, positive role models, and peers” for youth identity development. In this session, African American youths saw examples of African Americans who left legacies, and each participant was required to compose thank you letters to the visiting positive role models and peers. The inclusion of positive role models and peers was based on the premise that African American youth “lack strong role models and mentors to guide them through the explorations that naturally occur as a part of adolescent self-identity development” (Clark et al., 2005, p. 337). Youth also participated in discussions on topics such as how peers, family, and other environmental factors influence their lives and how their choices affect future goals. The authors gave limited information about ways in which the program was specifically tailored for African American youths. There were no details, for example, on how the role models were recruited to visit the classes. Findings revealed that adolescents in the intervention group reported higher rates of abstinence and less sexual intentions than those in the control group when assessed 19 weeks after the baseline survey. For the youth who did not have any sexual behaviors before the start of the study, those in the intervention groups were less likely to report the initiation of sexual activity than were controls. At 1-year follow-up, the abstinence effect was still significant among the male participants in the intervention group.

The Strengthening Families Program

The Strengthening Families Program (SFP) is a family skills training approach developed in the United States that uses 2-hour-long sessions with parallel Parent Skills Training and Children’s Skills Training groups in the first hour, followed by multifamily family skills training groups in the second hour. This multicomponent approach combines youth enhancement with a focus on family strengthening and improve parenting to support and guide youth. Designed to increase resilience and reduce risk factors for a broad range of behavioral, emotional, academic, and social problems in children 3 to 16 years old, the program is widely recognized as an effective substance abuse prevention program. SFP has been evaluated many times by independent researchers in randomized control

trials or health services research with very positive results in reducing substance abuse and delinquency risk factors by improving family relationships (e.g., Aktan, Kumpfer, & Turner, 1996; Gottfredson, et al., 2006; Kumpfer, Alvarado, Tait, & Turner, 2002; Spoth & Molgaard, 1999).

During the 1990s, cultural adaptations of SFP were made and evaluated among five ethnic minority groups (African, Hispanic, Asian, Pacific Islander, and Native American). Findings generally supported the efficacy of these adapted versions; however, important insights were gained. Although the culturally specific adaptations did not yield better effects compared to the core version, improvements were shown on recruitment and retention. Kumpfer et al. (2002) also reported that an adaptation of SFP for Hawaiian families (Kameoka, 1996) resulted in weakened prevention effects when behavioral skills training was curtailed to allow for extended focus on Hawaiian cultural values. This was an early exemplar of the principle of maintaining fidelity of deep structure or core intervention components while including adaptations to promote engagement. In 2003 the Cochrane Collaboration Reviews published a meta-analysis identifying SFP having demonstrably larger effects than other alcohol prevention programs (Foxcroft, Ireland, Lister-Sharp, Lowe, & Breen, 2003). This prompted international dissemination efforts in a number of nations. Several international adaptations that have used the Kumpfer model of cultural adaptation (see Table 17.1) have been subject to controlled trials, and all have demonstrated positive results among families with substance using parents (Kumpfer et al., 2008).

The Multiethnic Drug Abuse Prevention Program

The Multiethnic Drug Abuse Prevention Program was developed by Botvin, Griffin, Diaz, and Ifill-Williams (2001) using a cognitive-behavioral approach to enhance risk resistance skills among youth. Specific components of the curriculum include building self-esteem, resisting advertising pressure, managing anxiety, communicating effectively, developing personal relationships, and self-advocacy. The project was first tested on White youths in the state of New York in the 1990s and adapted for multicultural implementation with African American and Hispanic youth in 2001. Although the curriculum content and core elements of both programs were similar, several adaptations were made in preparation for delivering the program to youth of color. Focus groups were used to obtain feedback on ways to modify the intervention manual and adopt culturally appropriate and relevant teaching strategies. Adaptations included multiple strategies, such as creating graphics that contained the image

of minority youth, translating and adjusting the reading levels of program materials and questionnaires, and adding culturally relevant stories and role-play scenarios to the curriculum.

To test the culturally adapted program, participants were recruited from 29 public schools in New York City. Of the 5,222 seventh graders recruited, 3,621 completed the pre-intervention test, the posttest, and the 12-month follow-up survey. More than 60% of the participants were African American, and 22% were Hispanic. The schools were randomly assigned to an intervention condition or a control condition. Students in the intervention arm were taught by trained classroom teachers in 15 sessions in the seventh grade and 10 booster sessions in the eighth grade, while youths in the control group did not receive any project interventions (Botvin et al., 2001). The findings in the 3-month and 12-month postintervention tests revealed that the youth assigned to the drug prevention program reported greater reduction in smoking, drinking, and use of inhalant and polydrugs compared to youth in the control group. Programmatic effects observed in the culturally tailored program were similar to what Botvin et al. found among White, middle-class youth (Botvin et al., 1990; Botvin, Baker, Dusenbury, & Botvin, 1995). This program is one of few to advocate for programs that are multicultural rather than monocultural. Further, Botvin et al. (1995) found positive programmatic effects for both inner city minority and White, middle-class youth by making surface changes to the generic curriculum. These surface structure changes included inserting language and words to ensure that the reading level was appropriate for the targeted audiences and using photos of each racial/ethnic group to increase a sense of program relevance for each targeted audience.

Life Skills Training for Native American Youth

Schinke, Tepavac, and Cole, (2000) adapted the core curriculum from the Life Skills Training Program (Botvin et al., 1990, 1995, 2001) to prevent drug abuse among Native American adolescents. The program aimed to teach cognitive and behavioral skills to resist smoking of cigarettes or smokeless tobacco, drinking alcohol, and marijuana use. The participants ($n = 1,396$) were recruited from 27 tribal and public schools in 10 reservations in five states: North Dakota, South Dakota, Idaho, Montana, and Oklahoma. Participating schools were randomly assigned to one of three arms: two intervention arms (a skills-only intervention group and skills plus community involvement intervention group) and a control arm. Youth assigned to the two interventions attended fifteen

50-minute weekly sessions. The intervention sessions were designed to contain the important common cultural elements of Native American tribes, including values, legends, and stories that reflected issues in the Indian society. Specific consideration was given to including cultural content that celebrated Native American traditions, which were further tailored to be reflective of various reservations, tribes, and Indian groups. Similar to other culturally tailored programs, Schinke et al. (2000) included role-playing, storytelling, programmatic activities, and small-group discussions to foster cognitive-behavioral skill development. In addition, homework was assigned to encourage youth to use lessons learned in sessions in their everyday lives. Native American leaders facilitated the group sessions and also served as coaches and to facilitate resistance skills usage and efficacy. Family and community members who had frequent contact with youth also were involved in the intervention to elevate their awareness of substance use prevention and ways to support and encourage youth to avoid drugs and substances. Youth in the intervention arms received boosters semiannually, delivered in two 50-minute sessions. Youth in the control did not receive any intervention. The authors did not provide detailed descriptions of how the curriculum itself was culturally tailored or pilot-tested prior to implementing the program.

All youth completed a pretest, immediate and 6 months posttests, and follow-up assessments every 12 months over the course of 3 years. Outcome analyses showed lower levels of alcohol, smokeless tobacco, and marijuana use among youth exposed to the skills-only intervention compared to those assigned to the skills plus community involvement arm and those in the control group. The absence of intervention effects for the skills plus community involvement condition was thought to have been associated with potential dilution of the program effectiveness. The skills-only program allowed youth to focus more on internalizing core elements of the program, thereby enhancing resistance skills to prepare youth to avoid risky situations. Given the prior lack of culturally sensitive interventions for Native American youth, these findings were important for demonstrating that benefits of culturally tailored risk prevention programs for tribal communities. There remains a great need for conceptual models to guide the development and testing of new prevention approaches and to refine existing interventions for Native American youth.

Keepin' It R.E.A.L.!

Keepin' It R.E.A.L. is a program designed to reduce sexual risk among African American adolescents, with specific

emphasis on relationship development between mothers and sons (DiIorio et al., 2006). Mother-son dyads are randomly assigned to one of two conditions: a program based on social cognitive theory (SCT) or the life skills program (LSK) based on problem behavior theory. Participants in both conditions meet for seven sessions over 14 weeks for 2 hours each session. In the SCT condition, mothers and adolescents attended joint sessions together for four of the seven sessions and separately in concurrent sessions for the remaining three. Sessions included information about reducing HIV transmission, general communication skills, as well as specific communication about sex, values, decision making, and peer influence. In the LSK condition, participants attended separate concurrent sessions, except for a portion of the first and last sessions, where they attended joint sessions. In addition, mothers participated in a stress reduction activity and discussion about ways to apply session topics to personal life. Youth also received a stress reduction activity at the beginning of session accompanied by discussion about a specific risk behavior. The LSK condition also included community service activities, such as an overnight trip to a historically Black college or university. Both the SCT and LSK conditions were highly interactive, and home practice exercises were assigned at each meeting.

Keepin' It R.E.A.L.! was evaluated in a RCT with the two described intervention conditions (SCT: $n = 194$; LSK: $n = 187$) and a control condition ($n = 201$) that received 1 hour of training on HIV prevention. Assessments were given at baseline, followed by 4-, 12-, and 24-month postintervention assessments. Results revealed greater change in HIV knowledge and more recent conversations about sex with their sons among mothers assigned to the SCT condition compared to those in the LSK and control conditions. Mothers in the SCT and LSK conditions, however, scored higher on intent and comfort about discussing sex with their sons than mothers in the control group. Regardless of conditions, sons demonstrated significant increases in sexual behavior over time and increased sense of self-efficacy for abstinence. Further, SCT youth reported greater increase in HIV knowledge, twice the rate reported by those in the LSK condition. Further, compared to youth in the control condition, youth in both SCT and LSK conditions who were sexually active were more likely to use condoms and reported having intentions to end their sexual activity until they were older.

Keepin' It R.E.A.L.! was subsequently adapted as a father and son program, based on the SCT approach, which was similar in format and content to the mothers' program (DiIorio, McCarty, Resnicow, Lehr, & Denzmore, 2007).

The father-son program was evaluated with a randomized cluster design comparing effects of the SCT condition ($n = 141$) with a control condition focusing on nutrition and exercise ($n = 132$). Fathers in the SCT condition reported more communication and greater intent to communicate. Sons in the SCT condition reported less sexual behavior and more condom use among those who were sexually active. Increased condom use was a more robust finding than abstinence, despite being addressed in only one of the program sessions. The authors suggest that for African American male adolescents, condom use may be a more acceptable HIV prevention strategy than abstinence. The implications of cultural tailoring for study findings were not addressed.

Culturally Grounded Interventions

Strong African American Families

Strong African American Families (SAAF) is a family-based preventive intervention program, specifically designed for rural African American families to delay sexual debut and prevent the onset and escalation of alcohol and substance use during early adolescence. SAAF's theoretical and empirical underpinnings were based on data that Murry, Brody, and colleagues (Brody et al., 1994; Brody, Flor, & Neubaum, 1998; Brody, Murry, Kim, & Brown, 2002; Brody Murry, Gerrard, et al., 2004; Gibbons, Gerrard, Cleveland, Wills, & Brody, 2004; Wills, Gibbons, Gerrard, & Brody, 2000; Wills, Gibbons, Gerrard, Murry, & Brody, 2003) gathered for more than a decade from the target population. Community members were involved in numerous aspects of SAAF's design, development, and implementation. Community members, who are residents of the counties in which SAAF families live, act as contacts between Murry's research group and the prevention-targeted communities. These individuals are selected on the basis of their positive reputations and extensive social contacts in their communities. Their feedback is sought on the cultural relevance and sensitivity of measures, protocols, and procedures for rural African American parents and youth. They also provide feedback about families' and communities' cultural identities, meanings, customs, religious practices, languages, expectations, worldviews, and cultural values, all of which impact the successful design and implementation of the SAAF program. Community members reviewed and provided feedback on each session, and the program was pilot tested with rural African American families and refined prior to implementation. Further, to enhance rapport, minimize cultural bias, and enhance cultural understanding, African

American community members serve as home visitors to collect data, and African American community members were trained to implement the program.

The SAAF program consists of 2-hour sessions that take place for 7 consecutive weeks. After a meal catered by local African American providers, caregivers and children attend concurrent breakout sessions and then join together for the last hour of large group discussion. Parent sessions focus on both general (e.g., monitoring and communication with children about sex) and racially specific (e.g., racial socialization) parenting practices via videotaped vignettes and group discussions designed to foster social support. Adolescent sessions focus on topics such as dealing with difficult situations (e.g., racism), images of risk behaviors, and peer pressure resistance strategies via games and activities. Family sessions reinforce the messages from parent and adolescent sessions and build mutual understanding and communication. Each session includes videotapes of African American family interactions that were specifically produced for the SAAF program to illustrate key points. Further, each participating family receives program worksheets, homework assignments, and publications designed to reinforce program content. Every session includes a review of homework, didactic presentation of information, group discussion, and skills training activities designed to reinforce learning.

The SAAF program was evaluated by randomizing counties in the rural South to either the SAAF condition or a minimum-control condition in which parents were mailed information about early adolescent development, stress management, and techniques to promote exercise. Pretest assessment occurred prior to SAAF exposure, followed by 3-month posttest, with follow-up assessments at 9, 27, 54, and 65 months postintervention. The trial included 677 families with an 11-year-old child and has yielded favorable behavioral outcomes and supported SAAF's efficacy in delaying sexual debut and substance and drug use avoidance 54 months postintervention (Murry, Berkel, Brody, Miller, & Chen, 2009), with sustaining effects 65 months postintervention (Murry, Berkel, Pantin, & Guillermo, 2011) among SAAF youth. The researchers tested the mediating mechanisms of the program and found that participating in SAAF evoked increases in regulated-communicative parenting and intrapersonal protective processes in youth (racial identity, self-esteem, self-regulation, and positive self-image). SAAF-induced effects on parenting behavior deterred not only precursors to risk behavior, such as risk-related attitudes, future orientation, self-regulatory capacity, and resistance efficacy, but also immediate HIV-related risk behavior, including

early onset of substance use and sexual intercourse and alcohol use trajectories (Berkel et al., 2009; Brody, Murry, Gerrard, et al., 2004; Brody, Murry, McNair, et al., 2005; Murry et al., 2011; Murry et al., 2012;).

Mexican/Mexican American Keepin' It R.E.A.L.

Mexican/Mexican American REAL is a school-based program to reduce drug/substance use, alcohol use and other risky behaviors that included a multicultural version and a specific adaptation of Keepin' It R.E.A.L. for Hispanic youth. The developers describe the program as a culturally grounded intervention (Hecht et al., 2003) because the core elements of the curriculum were selected and designed to specifically reflect narratives and stories of indigenous youth (Hecht, Graham, & Elek, 2006). The curriculum focuses on five core components: communication competence, narrative knowledge, motivating norms, social learning, and resistance skills. In addition, cognitive skills, appropriate behaviors, and communication for effective use of four "REAL" resistance strategies (*refuse, explain, avoid, and leave* are emphasized (Kulis et al., 2005).

Multiple strategies were used to adapt the curriculum to fit the ecological culture of the targeted audience. Each session was translated into Spanish for the Mexican American version, while English was the language included in programs targeting White and African American youth. Local youth were involved in the production of curriculum videos, as part of the development team, and also served as actors for the videos. Examples from local mass media also were included as supplemental materials. For the boosters, activities were added that were developmentally appropriate for eighth graders, such as group poster projects, mural paintings, neighborhood nights out, and an essay contest (Kulis et al., 2005). The adapted curricula were subsequently tested on the three ethnic groups.

Three versions of Keepin' It R.E.A.L. were taught by trained classroom teachers of seventh graders in an efficacy trial to address the question as to which version was most effective for Mexican American youth: a Mexican American-focused version, a White and African American version, and a multicultural version. The program was delivered in ten 40- to 45-minute sessions to a sample of 3,402 seventh graders from 35 middle school in Phoenix, Arizona (Hecht et al., 2006; Kulis et al., 2005). Participants in the multicultural version received half of the lessons from the Mexican American-focused version and the other half from the White and African American version. After the intervention sessions were conducted, the youth in the intervention group received monthly boosters for the following year. Youth in the control group were not

exposed to programmatic materials or information. Data were collected at four time points, with a pre-intervention survey and three follow-up surveys. Results revealed significant reduction in substance and drug use, intentions for future drug/substance use, and positive images of peers who use drug/substance among participants across each of the program versions. Further, Hecht et al. (2006) noted that Mexican youth exposed to the program, regardless of version, delayed onset of drug/substance use and reported lower rates of increase in drug/substance use, over time, compared to youth in the control groups. On the contrary, the White and African American version had no significant effect. The authors highlighted the importance of cultural inclusion rather than exclusion in program content, targeting risk reduction among Mexican American youth. The positive behavioral outcomes manifested in a multicultural risk prevention curriculum suggests that programs narrowly tailored to each specific subgroup, in this instance Mexican Americans, may not be necessary.

Familias Unidas

Familias Unidas is a preventive intervention for Hispanic youth based on ecodevelopmental theory (Pantin, Schwartz, Sullivan, Prado, & Szapocznik, 2004) that promotes four major family processes that are protective against substance use and unsafe sexual behavior: positive parenting, parent-adolescent communication, parental involvement, and family support (Pantin et al., 2003; Prado et al., 2007). Consistent with Hispanic cultural expectations, Familias Unidas places parents in positions of leadership and expertise in helping to prevent problematic outcomes in their adolescents.

Familias Unidas has been evaluated in two separate RCTs, and a third and a fourth are ongoing. In the first RCT, Pantin, Coatsworth et al. (2003) found that Familias Unidas was efficacious (in a sample of 167 Hispanic youth), relative to a no-intervention control group, in (a) increasing parental involvement, parent-adolescent communication, and parental support for the adolescent; and in (b) reducing adolescent behavior problems, a risk factor for drug use and HIV risk behaviors. Results also indicated that increases in parental investment were related to decreases in behavior problems. In the second RCT, Prado et al. (2007) evaluated the efficacy of Familias Unidas relative to a family-centered HIV preventive intervention and a family-centered adolescent cardiovascular health intervention over a 3-year period. The results of the study found that, over time, Familias Unidas was efficacious in reducing illicit drug use relative to the cardiovascular preventive intervention. Familias Unidas was also found

to be efficacious in reducing cigarette use over time when compared to both the HIV preventive intervention and the cardiovascular preventive intervention. Finally, Familias Unidas was efficacious in reducing unprotected sexual behavior and preventing the incidence of sexually transmitted diseases. The effects of Familias Unidas on smoking and illicit drug were partially mediated by improvements in family functioning. Specifically, improvements in positive parenting (reward contingencies offered by parents) and in parent-adolescent communication explained some of the effects of intervention condition on cigarette and illicit drug use.

CHAMP Family Program

CHAMP was designed in close collaboration with community members, who played major roles in the program's design, implementation, and evaluation to reduce HIV risk among urban African American adolescents (McKay et al., 2004). CHAMP utilized a community participatory approach and framework to design, develop, and implement the program. The framework focuses on several core principles: (a) linking collaborators from outside and within a setting to design a program; (b) creating a stakeholder advisory group to oversee program activities; (c) designing programs that integrate scholarly and indigenous knowledge and perspectives relevant to a setting to shape prevention messages and activities; and (d) using credible messengers, such as those already connected to a setting, to implement interventions in their own neighborhoods (Baptiste et al., 2006).

All CHAMP participants ($n = 227$ families) were chosen randomly from a roster of 550 youth attending four inner city public elementary schools. Families were recruited via research staff making presentations to fourth- and sixth-grade classes, personal contact with parents at schools, telephone calls, or home visits to parents. As with other family-based interventions, CHAMP was designed as a multiple-family group format. This delivery format was thought to be especially useful in increasing social support and mutual aid, creating strategies to deal with common problems, and reducing the stigma associated with talking about sensitive issues (McBride et al., 2007). The format is somewhat unique in that pretest data collection occurs in session 2, after the participants have had the opportunity to meet facilitators and other families and find out more about the goals of the program. Posttest data collection occurs in session 11, before the celebration in the final session. In the non-data collection sessions, families meet for 20 minutes to discuss the topic of that night's meeting and then break out for 45-minute concurrent parent and

child sessions. In youth sessions, participants learn and role-play peer-pressure scenarios and communication with parents. The parent sessions focus on monitoring, social support, problem solving, and communication with young people about sensitive topics. At the conclusion of each meeting, parents and children reconvene to make an action plan for their home practice activities.

CHAMP was initially evaluated with a pretest-posttest design that included a no-treatment comparison group from the CHAMP longitudinal study (McKay et al., 2004). Results from parents demonstrated a positive increase in family decision making relative to the comparison group, as did comfort in communicating about sensitive topics with their children. Among adolescents, those in the intervention group were less often exposed to mixed-gender settings without parental supervision (McBride et al., 2007). Youth in the intervention condition reported higher family conflict, perhaps because the intervention encouraged them to communicate more with their parents about sensitive topics. In addition, CHAMP has demonstrated effects on externalizing behaviors. Moreover, the CHAMP Family Program has been adapted for and piloted in several other contexts, including HIV-positive youth (McKay et al., 2007), in multiple international settings, such as South Africa (Bell et al., 2008) and Trinidad and Tobago (Baptiste, Voisin, Smithgall, Martinez, & Henderson, 2007). The program developers attribute positive results to the use of core community participatory principles to inform and guide efforts to refine a disease prevention and health promotion model and to refine research methods that embed culture as well as context in the design, development, and implementation of the CHAMP program.

The initial CHAMP outcome study did not include a control arm, but subsequent adaptations of the program have added experimental control comparisons, showing continued strong programmatic effects on increased HIV transmission knowledge, less stigmatizing attitudes toward people with HIV, increased sexual conversation between parents and youth, and increased community protective influences, via more social control and the promotion of social action to foster health-enabling communities for youth (McBride, Baptiste, Traube et al., 2007).

Puentes a la Secundaria/Bridges to High School

Puentes a la Secundaria/Bridges to High School (Puentes/Bridges) is a family-based program originally intended to prevent mental health disorders and prevent school dropout among Mexican American adolescents attending schools in low-income, urban communities (Gonzales,

Dumka, Millsap, et al., 2012); however, it also produced reductions in adolescents' overall substance use, sexual risk taking, and alcohol use disorder (Germán, Gonzales, West & Dumka, in review; Gonzales et al., 2012; Gonzales, Jensen, Wong et al., in review; Gonzales Wong, Toomey, Millsap, Dumka, & Mauricio, 2014; Jensen, Wong, Gonzales, et al., 2014). Content and structural elements of Puentes/Bridges were based on: (a) qualitative interviews and focus groups with Mexican-origin families including Spanish-speaking and English-speaking parents who were born in the United States or Mexico; (b) key informant interviews with school personnel and service providers; (c) more than a decade of programmatic, longitudinal research on risk and protective processes within the targeted communities (Dumka, Gonzales, Woods, & Formoso, 1998; Gonzales, Dumka, Mauricio, & Germán, 2007); and (d) extensive pilot testing and refinement with schools and families in these communities (Dumka et al., 1998; Gonzales, Dumka, Deardorff, Jacobs-Carter, & McCray, 2004). The program's nine weekly sessions and two home visits targeted parenting practices, child coping, family cohesion, and school engagement. Puentes/Bridges was guided by an ecodevelopmental framework (Szapocznik & Coatsworth, 1999) that recognizes unique cultural factors and experiences that impact youth development at multiple levels and the challenges they encounter in their families, neighborhoods, and schools and with peers. In particular, school engagement can be a challenge for Mexican American youth in the United States. For example, since 1980, Mexican Americans have had the lowest rates of high school completion compared to Whites, Blacks, Asian, and Pacific-Islander groups and also compared to youth from other Latino groups (U.S. Census Bureau, 2010). Mexican American youth in low-income communities find it difficult to envision positive future possibilities (possible selves) in a context of low-wage jobs, unemployment, experiences of discrimination, and reduced expectations directed at them (Oyserman & Markus, 1990). These constraints combine with other features of low-income communities, such as the availability of drugs, deviant peers, and parents' diminished capacity for effective parenting, to reduce investment in education and increase risk for mental health problems (Barrera et al., 2002). Immigrant youth also encounter cultural conflicts in their schools and families, and their parents often have a poor understanding of U. S. schools and are therefore ill prepared to monitor and intervene when they have academic difficulties (Suarez-Orozco & Suarez-Orozco, 1995). Puentes/Bridges aimed to promote family and youth competencies shown in the general literature to protect against youth risk behaviors while also

empowering families to address these unique challenges and support their adolescents' school engagement across the middle school and high school years. In addition to program content specifically designed to increase knowledge and engagement of families in promoting their adolescents' educational success, the program integrated the values of *familismo* (e.g., strong orientation and commitment to family) through both content (cohesiveness, coparenting, adolescent family support seeking) and structure (recruiting multiple caregivers, joint sessions with parents and children). Furthermore, traditional values were used as a motivational tool to encourage parents and adolescents to internalize and incorporate elements of the program into regular practice (Gonzales et al., 2007).

Puentes/Bridges was evaluated via an RCT with 516 families from four middle schools in Phoenix, Arizona (Gonzales et al., 2012). An efficacy trial showed desired effects on adolescent externalizing, internalizing, substance use, school disciplinary actions, and grades and also provided evidence that these effects were mediated by targeted changes in parenting, adolescent coping efficacy, family cohesion, and school engagement. However, tests of moderation also showed that effects on program mediators varied on the basis of family acculturation level. Greater changes in parental monitoring and family cohesion were found for more acculturated English-dominant families, whereas the less acculturated Spanish-dominant families showed greater intervention effects to reduce harsh parenting and to increase adolescent coping efficacy and school engagement. Patterns of engagement also varied, with less acculturated families enrolling and attending at higher rates. However, despite acculturation-related differences at immediate posttest, long-term follow-up showed similar improvements across groups on the quality of parent-child relationships (Jensen et al., 2014) and school engagement (Gonzales et al., 2014) following transition to high school (ninth grade), and these changes mediated subsequent effects to reduce internalizing and externalizing symptoms substance use, diagnosed alcohol use disorder, and school dropout at 5 years posttest when most adolescents were in the twelfth grade (Gonzales, Jensen, Wong et al, in review).

Be Proud! Be Responsible!

Be Proud! Be Responsible!, developed by Jemmott, Jemmott, and Fong (Jemmott, 1992; Jemmott, Jemmott, Fong, & McCaffree, 1999; Jemmott, Jemmott, Fong, 2010), was designed to increase minority youth knowledge on sexually transmitted diseases (STDs), HIV, and acquired immune deficiency syndrome (AIDS); to affect their attitudes and

beliefs about safer sex (condom use and other prevention strategies); to inspire their understanding on individuals' vulnerability to STDs, HIV, and AIDS; and to build their skills on negotiation and resistance to risky sexual behaviors. This culturally grounded intervention was subsequently adapted and evaluated with different ethnic groups and subpopulations. Be Proud! Be Responsible! The African American Male version (Jemmott, 1992) sampled 157 Black male adolescents (mean age 14.4 years) from three sites in Philadelphia, Pennsylvania: a medical clinic, a local high school, and a local YMCA. Following a pretest evaluation, a total of 27 small groups were formulated of which 14 were assigned to the intervention condition and 13 to the control condition. Both conditions consisted of small group interventions lasting 5 hours in which the curriculum was delivered via videotapes, small-group discussions, exercises, and games. Prior to program implementation, curriculum materials were pilot tested and revised to fit the interests of African American boys in urban regions. Although limited information was provided about specific adaptations for African American males, it appears that the cultural tailoring primarily involved the use of African American images in the videos and a gaming format to deliver program content. For example, they revised the AIDS knowledge content to be delivered as "AIDS Basketball," with the game organized around how to avoid risky sexual behaviors. The program developers also characterized the selection of program implementers as evidence of cultural tailoring; all were African American adults with at least a 4-year college degree and were employed as educators, social workers, or health professionals. Posttests were administered immediately after the intervention, and a follow-up assessment was conducted 3 months later. Findings revealed that intervention targeted youths reported greater change in AIDS knowledge, less favorable attitudes toward risky sexual behaviors, and fewer intentions to participate in such behaviors. Intervention effects remained significant at 3 months follow-up.

Be Proud! Be Responsible! The New Jersey version (Jemmott et al., 1999) targeted 496 African American youth (male and female) residing in Trenton, an economically distressed urban community with high prevalence rates of unintended pregnancies and sexually transmitted infections (STIs). The participants were recruited from local public schools in seventh and eighth- grades (mean age 13.8 years). Similar to the original version, the students were randomly assigned to the intervention groups (HIV reduction intervention) and the control groups (general health promotion intervention). Each small group

contained six to eight adolescents, with youth attending a 5-hour session in one day. The intervention targeted “hedonistic beliefs about the positive consequences of condom use on sexual enjoyment and self-efficacy to implement condom use” (Jemmott et al., 1999, p. 168). Prior to implementation, this curriculum was pilot tested to ensure that the program content and activities were sensitive to urban African American adolescents’ interests. For instance, the videos selected for the program included multiethnic actors talking about AIDS, such as Ruben Blades, a Latino actor, demonstrating how to put on a condom using a banana as a prop (Jemmott et al., 1999, p. 168). Participants in the general health promotion intervention groups were taught about nutrition, exercises, and diseases. Local African Americans were hired to deliver the program. Posttest data collection occurred immediately after the intervention, and follow-up tests were conducted 3 and 6 months postintervention. Results from the immediate postintervention test revealed more positive beliefs about condom uses, stronger condom use intentions, and higher self-efficacy among the participants in the intervention groups compared to those assigned to the control condition. However, no group differences on targeted behaviors were observed at 3-month follow-up. Finally, the 6-month follow-up tests found decreased HIV-related unprotected sexual behaviors and fewer sexual partners compared with youth in the general health promotion intervention groups.

The Children’s Health and Responsible Mothering—Project CHARM

The Children’s Health and Responsible Mothering—Project CHARM is another adaptation of Be Proud! Be Responsible! specifically designed to reduce repeated pregnancy among Hispanic and African American adolescent mothers. CHARM is based on the core elements of Be Proud! Be Responsible! Project, which aims to reduce the number of episodes of unprotected sexual intercourse and number of sexual partners as well as to facilitate increased willingness and intentions to use condoms in order to prevent repeated pregnancies (Koniak-Griffin et al., 2003). A total of 572 adolescents, age 16.7 on average, who were mothers or pregnant were recruited, with a total of 497 retained for all the data collection cycles. The evaluation sample was primarily Hispanic (78%), with a smaller representation of African American females (18%). Youth were grades 7 through 12 from four school districts in Los Angeles County, California. Schools were randomly assigned to treatment (HIV program) or control conditions (general health program). Each program consisted of four 2-hour sessions.

Compared to the original curriculum, Project CHARM focused more on the impact of STDs and HIV on the lives of pregnant girls and their babies, including STI prevention during pregnancy and the importance of avoiding risky sexual practices postpartum. Similar to the original program, supplemental materials included culturally and developmentally relevant videos, skill-building activities, and role-playing. In addition, an HIV-infected adolescent mother was invited to speak in the second intervention session to motivate participants to make healthy sexual decisions and to encourage them to be sexually responsible and accountable as well as politically aware of the effects of HIV/AIDS on inner-city communities and their children (Koniak-Griffin et al., 2003). Assessment occurred immediately after the intervention sessions ended, followed by 3-month, 6-month, and 1-year postintervention assessments. Findings revealed that at 6-month follow-up, compared to females in the control group, those assigned to the HIV program demonstrated greater change in AIDS knowledge, scored higher in behavioral intentions to use condoms, and reported greater reduction in number of sexual partners.

Following several demonstrations of positive program effects with varying subpopulations in the context of efficacy trials, Jemmott, Jemmott, Fong & Morales (2010) launched an implementation effectiveness trial of Be Proud! Be Responsible! wherein members of community-based organizations (CBOs) were randomly assigned to different training conditions to test the role of different program implementation training strategies on intervention effectiveness. The Be Proud! Be Responsible! effectiveness trial was implemented through CBOs with 1,707 youth age 13 to 18 residing in Philadelphia, Pennsylvania, and Trenton, New Jersey. There were three training conditions: (1) CBOs received the two project manuals (HIV prevention and health and wellness curricula) and no face-to-face training; (2) CBOs received the project manual and attended a 2-day face-to-face training session; and (c) CBOs received the project manual, 2 days of training on the manual, followed by a practice intervention-implementation session and 2 days of additional training in which program implementers were videotaped and received feedback on their intervention implementation practice sessions. Similar to their other program designs, Jemmott et al. randomly assigned youth to one of two conditions: Youth in intervention group received the HIV/AIDS prevention program, and those in the control group received general health promotion intervention sessions. Subsequently, trained CBOs at each site were randomly assigned to each youth group.

Preintervention data were collected and follow-up tests were administered at 3, 6, and 12 months postintervention exposure. However, results revealed no statistically significant program impacts on frequency of sexual intercourse in the past 3 months or condom use at last intercourse, and there were no effects of training condition on intervention effectiveness. The researchers concluded that there was no evidence that the amount of time training community members improved facilitators' efficacy in program delivery. Further, they offered several explanations for lack of significant program effects. First, they surmised that there may have been a selection bias among CBOs, as many who declined to participate held strong values about teaching abstinence that conflicted with the program's philosophy. Further, they speculated that the community-based delivery site may have affected study findings as prior school-based evaluations consistently showed expected benefits on risk behaviors for youth that received the intervention compared to those in the control group.

Respeto/Proteger

The Project Respecting and Protecting Our Relationships (Respeto/Proteger) is an adapted version of Be Proud! Be Responsible! designed to prevent HIV/AIDS among Latino adolescent couples who are new parents. The program focuses specifically on enhancing parental protectiveness as a mechanism to foster resiliency and motivate youth to reduce risky sexual practices (Lesser, Koniak-Griffin, Huang, Takayanagi, & Cumberland, 2009). Respeto/Proteger was informed by results from Project CHARM, which revealed that young Latino mothers felt less empowered to negotiate condom use, compared to their male partners. Lesser, Koniak-Griffin, Huang, Takayanagi, and Cumberland (2009) concluded from these findings that there is a need to include young fathers in pregnancy and sexual risk reduction program to increase sexual protection among young mothers.

Adolescent mothers and their partners were recruited from 28 Women, Infants, and Children (WIC, a federal assistance program of the Food and Nutrition Service of the United States Department of Agriculture (USDA) for healthcare and nutrition of low-income women, infants, and children) sites, eight alternative schools, two community-based organizations, and one community-based clinic in Los Angeles (Lesser et al., 2009, p. 92). Through randomization, 84 couples were assigned to the treatment condition and 86 couples were assigned to the control condition. The mean age was 20 years for males and 18 years for females; 78% of the male parents and 86% of the female parents were Latino. Young parents in

the intervention group received a 12-hour curriculum (six 2-hour sessions) while their counterparts in the control group received a 1.5-hour didactic HIV prevention education program. A total of 41 program series were held in local settings, such as CBOs and health clinics. Evaluative surveys were conducted at pre-intervention, immediately after the intervention, and during follow-up periods.

Respeto/Proteger program design and development were informed by qualitative data received from Latino mothers attending health promotion projects for young mothers, and intervention approaches and strategies adopted from Project CHARM, Project EL Joven Noble, and Project Con los Padres. The Project EL Joven Noble and Project Con los Padres were programs designed to influence Latino youth understanding of romantic relationships, the "macho" man stereotype, and strategies to deal with daily challenges. Prior to implementing Respeto/Proteger, the program was pilot tested with groups similar to the targeted audience. Programmatic changes were made based on pilot test results. Findings revealed that the couples, both adolescent mothers and their partners in the intervention arm of the study, reported less unprotected sexual behaviors 6 months after the intervention sessions. Further analysis indicated that, compared to the adolescent mothers in the control group, those in the intervention group whose male partner scored higher on protectiveness at the baseline reported even less unprotected sexual behaviors. Interestingly, female protectiveness did not show significant influences on males' unprotected sexual behaviors.

¡Cuidate!

¡Cuidate! (English translation means: take care of yourself) is a program to increase parent-child communication about sex in Mexican-heritage families (Villarruel, Cherry, Cabriales, Ronis, & Zhou, 2008). The theory of planned behavior was combined with a focus on Mexican cultural values, including familism and religiosity, in the design of the program. Familism was utilized as a motivational impetus to frame the importance of talking with children about sex. Religiosity was considered for its influence on decision making about sexual behavior, for example, whether to address condom use or abstinence. The 6-hour program was conducted with separate sessions for parents and adolescents on two consecutive Saturdays. The intervention group (n = 404) was provided with information about the prevention of HIV and pregnancy as well as support for general and sexual communication with children. The control group (n = 317) received a 6-hour general health intervention, which also emphasized the principal

role of parents in determining adolescents' health but did not discuss parent-child communication. Both groups were assigned homework to complete with their adolescents. Participants in the intervention group reported more communication, communication about sex, and comfort in communication than parents in the control group. Moreover, this effect was mediated by the perceived utility of communication about sex; attitudes of the adolescent, family, and church related to communication about sex; and parents' comfort in discussing sex with adolescents. Effects were not moderated by familism, religiosity, or the gender of parents or children.

Discussion

Several observations emerged from our review of youth risk prevention programs. The majority of programs were delivered and tested in school based settings, with a significant proportion using community- and family-based approaches, and the majority of programs included multiple intervention components. Our review also revealed that the majority of programs targeting African Americans and Hispanics were designed to prevent risky sexual behavior and its consequences (e.g., STIs and HIV/AIDS). The program targeting Native American youth focused solely on substance and drug use prevention. Findings are generally positive in showing that these programs fostered significant reductions in risky sexual practices, increased sexual and HIV/AIDS knowledge, and reduced youth intentions to participate in risk behaviors, including reduced intent to use drugs/substances. Evidence of condom use and positive attitudes about condoms as a means for protecting against unintended pregnancies, STIs, and HIV was also noted. We also can conclude these programs foster sustained effects, extending from 12 months to 6 years after the intervention, on outcomes such as sexual debut, substance and drug use, number of sexual partners, and diagnosed substance use disorder. Thus, our review offers support for the usefulness of culturally tailored programs to prevent the consequences of early sexual activity onset and substance and drug use among youth of color. Finally, while each of the programs included in our review focused on the prevention of sexual initiation, there was a lack of consideration given to the inclusion of culturally tailored alcohol and substance use programs for youth of color.

Content of Adaptations: Beyond Surface Structure

In contrast to adaptations of PT interventions that have augmented well-established EBIs previously validated

with White samples, the majority of programs reviewed here were originally designed and developed for targeted youth of color. Only two programs, Life Skills Training (Botvin et al., 1990, 1995, 2001) and the SFP (Spoth, Redmond, & Shin, 2001), were first evaluated with White youth prior to being adapted for other ethnocultural groups (African American and Hispanic). In addition, our review determined that 10 of the 16 programs emerged from culturally grounded strategies involving extensive, in-depth community involvement and input on program design, measures, and various aspects of the program's deep structure. The integration of cultural values, beliefs, norms, and information about legacy and ancestry were apparent in most of the adapted and culturally grounded program. In addition, role-playing, role models, the use of storytelling, and the inclusion of culturally relevant materials, videos with images familiar to targeted audiences, and games were incorporated in the cultural adaptations. A noteworthy observation, however, is that none of the studies in this review, with the exception of SAAF program developers (Berkel et al., 2009; Murry, McNair, Myers et al., 2014), included measures to assess cultural dimensions in summative assessments of program effects.

Subgroup Differences

Our review also revealed evidence of subgroup differences in program responsiveness. For example, the Aban Aya Youth program demonstrated greater improvements in sexual risk and drug reduction among males than females. Projects that involved both teen mothers and fathers indicated that the intervention is more effective on the young fathers for condom use. In the one study that examined within-group variability based on culture, Puentes/Bridges showed differences in levels of engagement and program effects between low-acculturated (primarily immigrant Spanish-speaking) and high-acculturated (English-speaking) families. However, it was noteworthy that similar long-term effects on targeted mental health, substance use outcomes, and school dropout were found across levels of acculturation, suggesting that diverse families may benefit in different ways, depending on their unique needs when participating in comprehensive, multicomponent interventions. This finding underscores the need for greater attention to within-group variability in tests of efficacy and effectiveness in order to determine whether and how EBIs can address subgroup differences for racial/ethnic groups that are neither monolithic nor static on cultural dimensions.

Remaining Gaps and Questions

To further advance this area of research, there is a need to address several gaps and unresolved questions. Several groups are not visible in preventive intervention research—namely Alaska Native, Asian Americans, and Asian Pacific Island adolescents. No programs targeting these youth emerged in our comprehensive literature search, and very few have been evaluated with American Indian youth. American Indians and Alaska Natives are disproportionately affected by all major youth risk behaviors, including alcohol, tobacco, drug use, risky sex, and suicidal behaviors. Further, available studies have shown that while Asian Pacific Island youth are less likely to engage in risky sexual practices than any other group of adolescents (Grunbaum, Lowry, Kann, & Pateman, 2000), when this population has been included, findings reveal greater endorsement of having used alcohol or drugs at last intercourse compared to other racial/ethnic groups. Sexual risk protection was comparable to other ethnocultural groups in reports of using a condom at last intercourse. That there is a major void in targeting these subgroups in preventive interventions is of great concern. That the majority of risk prevention programs have been developed and tested on urban youth suggests the need to determine how to translate these programs in rural contexts. This information can further our understanding of the utility of developing programs to specifically address the needs of rural youth. The paucity of studies in our review that targeted rural youth is of concern, given that rural youth are at increased risk for using a variety of substances, including tobacco (Zollinger, Saywell, Overgaard, Przybylski, & Dutta-Bergman, 2006), alcohol (Hutchison & Blakely, 2003), and methamphetamines (Johnston et al., 2009). Several researchers have noted that rural Whites, especially those living in the Appalachian region, appear to be at greater risk for early onset of substance use than their more frequently studied urban counterparts (Johnston et al., 2009; Pruitt, 2009; Spoth et al., 2001; Zollinger, Saywell, Overgaard, Przybylski, & Dutta-Bergman, 2006). Not only are these youth infrequently included in epidemiological studies, but, to our knowledge, few programs specifically target White youth residing in southern rural communities.

Finally, only one study included in the current review directly compared whether a culturally adapted program is more effective than a generic or multicultural program (Keepin' It R.E.A.L.), and the findings did not support the cultural adaptation with Hispanic youth. Notwithstanding these limitations, our review offers support that it is possible to provide effective programming for youth from

a variety of cultural backgrounds with proven, culturally generic, evidence based programs. However, it remains unknown when, how, for whom, and under what circumstances culturally tailored youth risk prevention program will foster targeted programmatic effects.

PREVENTION OF ANXIETY AND MOOD DISORDERS

Anxiety and mood disorders are prevalent among children and adolescents, causing significant impairment at home, with peers, and at school (e.g., Kessler, Avenevoli, & Merikangas, 2001; Pina, Zerr, Gonzales, & Ortiz, 2009). Data from the National Comorbidity Survey Adolescent Supplement showed that among youth 13 to 17 years old, anxiety disorders were the most prevalent class of disorders (24.9%) with mood being the third most prevalent class of disorders (10%) (Kessler et al., 2012). In addition, Kessler, Avenevoli, and Ries Merikangas's (2001) seminal review of epidemiological and large-scale prospective longitudinal studies (e.g., Costello, Angold, & Keeler, 1999; Kovacs, Gastonis, Paulauskas, & Richards, 1989) revealed that anxiety disorders were not only the conditions most commonly comorbid with depression but that about three-fourths of depressed youth had a history of at least one anxiety disorder. There also are data suggesting some variations in the prevalence and co-occurrence of anxiety and mood problems (e.g., age, sex), albeit epidemiological or large scale studies focusing on ethnicity-race are scant.

In the adult literature, for example, Karno et al. (1989) found that the lifetime prevalence for any anxiety disorder was higher among U. S.-born Mexican Americans compared to Mexican Americans born in Mexico (22.6 versus 13.7), based on the Los Angeles Site of the National Institute of Mental Health Epidemiologic Catchment Area study. In another study using data from the National Epidemiologic Survey on Alcohol and Related Conditions, Grant et al. (2004) found that the lifetime prevalence of any anxiety disorder was higher for U. S.-born Mexican Americans (16.3%) than for foreign-born Mexican Americans (9.1%). In addition, for any mood disorder, the lifetime prevalence was higher for U. S.-born Mexican Americans (19.3%) than for foreign-born Mexican Americans (10.2%). Turning to the National Health and Nutrition Examination Survey-III, Riolo, Nguyen, Greden, and King (2005) found that Mexican Americans and Caucasians had a significantly earlier onset of major depressive disorder compared to African Americans. In addition, education (less than 8 years of school) was associated with

major depressive disorder for Mexican Americans only. Also, dysthymic disorder was greater among Mexican Americans and African Americans, even after controlling for the effects of poverty and education. (No age by ethnicity data were presented, albeit the youngest responders were 15 to 19 years old.) No comparable research focusing on Native Americans and Asian Americans has examined variations in the prevalence and co-occurrence of anxiety and mood problems.

When it comes to ethnic minority children and adolescents, there also is a scarcity of anxiety and depression disparities research from large-scale and methodologically robust studies; the little work that has been conducted has focused on African American and Hispanic/Latino youth. Bird (1996), for example, reported prevalence rates of psychiatric disorders in children and adolescents (age 4–16 years) in Puerto Rico. Findings showed that depression was the second most common diagnosis (5.9%) with only separation anxiety disorder and specific phobias emerging from the anxiety disorder modules at 4.7% and 2.6%, respectively. More recently, Alegria et al (2012) reported data from the National Comorbidity Survey Adolescent Supplement. Findings showed that for non-Latino Black youth, the lifetime prevalence of separation anxiety disorder was significantly higher (10.8%) compared to non-Latino Whites (6.8%) or Latinos (7.9%). No additional differences relevant to ethnicity and anxiety-depression were found.

The high prevalence of anxiety and depressive disorders in children and adolescents (Costello, Egger, & Angold, 2005), coupled with the fact that these conditions also are typically chronic and lead to significant impairment (including early initiation of the use of alcohol, tobacco, and other illicit drugs; Conger & Rueter, 1996; Glantz & Leshner, 2000), has prompted much interest in developing EBIs (Beidel & Turner, 2007; Silverman, Pina, & Viswesvaran, 2008). In addition, data showing some ethnic variations in the prevalence of these disorders prompted some research focused on culturally adapted interventions focused on reducing and preventing disparities in these psychiatric conditions among youth (Cardemil, Reivich, & Seligman, 2002; Cooley-Strickland, Griffin, Darney, Otte, & Ko, 2011).

In the clinical child and adolescent area, several randomized, controlled prevention trials have been published at the universal, selective, and indicated levels focusing on anxiety and/or depression. In the anxiety area, most anxiety prevention studies have been conducted at the universal level, and only a handful of studies have tested selective or indicated protocols. Indicated programs have

been directed at children due to a family history of anxiety (Ginsburg, 2009), a known predisposition (e.g., behavioral inhibition; Rapee, Kennedy, Ingram, Edwards, & Sweeney, 2005), or some anxiety symptoms (Pina, Zerr, Villalta, & Gonzales, 2012). Selective or indicated depression prevention programs have been directed to youth at risk for depression due to a family history or diagnosis of depression (Beardslee, Gladstone, Wright, & Cooper, 2003) or exposure to stressors that place them at high risk, such as urban poverty (Cardemil et al., 2002) or the loss of a parent (Sandler et al., 2003). Anxiety prevention strategies show content overlap with prevention protocols, typically focusing on negative cognitions, social skills training, and problem-solving skills (Barrett & Turner, 2001; Dadds, Spence, Holland, Barrett, & Laurens, 1997; Essau, Conrath, Sasagawa, & Ollendick, 2012; Lock & Barrett, 2003; Lowry-Webster, Barrett, & Dadds, 2001; Pina et al., 2012; Simon, Bogels, & Voncken, 2011; Warner, Weissman, Fendrich, Wickramaratne, & Moreau, 1992).

In a nutshell, research shows that pathological levels of childhood and adolescent anxiety and depression may be preventable. Using meta-analyses, Fisak, Richard, and Mann (2011) found a significant overall mean weighted effect size of about .18 as well as a slightly greater effect for targeted (i.e., selective or indicated) anxiety prevention over universal protocols. In another meta-analysis, Teubert and Piquart (2011) found significant effect sizes on anxiety at posttest (about .22 for symptoms and .23 for diagnoses) and follow-up (about .19 for symptoms and .32 for diagnoses) as well as significant variations of program effects, giving targeted programs an advantage over universal programs. When it comes to the prevention of depression, targeted (selective/indicated) programs also appear to be more effective than universal programs. In Horowitz and Garber's (2006) meta-analyses, the pretest-to-posttest effect size for targeted programs ranged from about .23 to .30 compared to only .12 for universal programs. At follow-up, the effect size for targeted programs ranged from about .31 to .34 compared to .02 for universal programs. Similarly, Stice, Shaw, Bohon, Marti, and Rohde's (2009) meta-analyses found an average effect size ranging from .14 to .23 for targeted programs compared to .06 to .04 for universal programs. It is important to note, however, that there was some overlap in the trials included in the anxiety and depression meta-analyses, making these effect size evaluations nonindependent. For example, the FRIENDS for Life program was included in both anxiety and depression meta-analyses (e.g., Barrett, Farrell, Ollendick, & Dadds, 2006). Nonetheless, based on these findings, preventing anxiety and depressive disorders appears to be both an

effective and an efficient way to reduce the incidence of these psychiatric conditions, especially when the protocols are targeted.

Cultural Adaptation and the Prevention of Anxiety and Depression

Despite the promising overall effects of childhood anxiety and depression prevention programs, evidence seems to suggest that prevention efforts need to be strengthened, particularly for ethnic minority or culturally diverse child populations (e.g., Barrett, Sonderegger, & Xenos, 2003; Cardemil et al., 2002; Cooley-Strickland et al., 2011; I. D. Miller et al., 2011). Studies testing cultural adaptations of anxiety and depression prevention programs are far less common than for substance use, externalizing problems, or other youth risk behaviors. In this section we highlight the few prominent exemplars that have been evaluated to date.

The Penn Resiliency Program

The Penn Resiliency Program is a school-based depression prevention program that teaches cognitive and social problem-solving skills to groups of middle school students who might be at risk for developing depressive symptoms by virtue of their low-income status. The program is an adaptation of an EBI established with predominately middle-income Caucasian children (Gillman, Reivich, Jaycox, & Seligman, 1995) that Cardemil et al. (2002) applied to low-income African American and Latino children. In their study, the original structure of the program was maintained, but cultural modifications were made. In particular, the characters and examples used in the program were made culturally and contextually relevant by illustrating the skills in the context of challenges associated with growing up in a single-parent home and managing peer conflict in urban settings. For Latino youth, fewer depression symptoms, negative automatic thoughts, and hopeless thoughts were found from pretest to posttest and at 6-month follow-up. In addition, at the follow-up only, Latino youth also evidenced higher self-esteem over time compared to no-treatment controls. For African American youth, however, findings were not encouraging. There were no differences between the intervention and the control conditions on any of the measures or at any measurement point. At the 2-year follow-up (Cardemil, Reivich, Beevers, Seligman, & James, 2007) Latino youth continued to show program-related gains but African American youth did not. The finding with African American children was interpreted as possibly being related to six potential factors, two of which seem especially plausible: cross-ethnic

differential reporting of depressive symptoms and/or differential expression of depression symptoms. However, with no data about the measurement equivalence of the assessment tools used in the study, it is difficult ascertain the viability of these possibilities (Pina, Gonzales, Holly, Zerr, & Wynne, 2012).

FRIENDS for Life (African American Youth)

In another study focusing on African American youth, Cooley, Boyd, and Grados (2004) and Cooley-Strickland et al. (2011) described their adaptation of the FRIENDS for Life childhood anxiety prevention program developed in Australia to prevent anxiety and depression (Barrett et al., 2006). The original EBI is a school-based intervention shown to reduce risk of anxiety disorders by teaching cognitive and emotional skills to cope with feelings of fear, worry, and depression. In the cultural adaptation work of Cooley and colleagues, modifications focused on replacing references specific to Australian culture and including examples relevant to the contexts of inner city African American youth. In terms of language, for example, the word *vegemite* was replaced with *peanut butter*, *sugar glider possum* was replaced with *bat*, and *tucker* was explained as *food*. Also, relaxation exercises were modified, and references to animals indigenous to Australia (e.g., koalas, cattle dogs, crocodiles) were replaced with those found in urban America (e.g., pigeons, mice, dogs, kittens). As far as incorporating the contexts of inner-city African American youth into the intervention, children drew pictures of real-life violence that had occurred in their communities, and examples of incidents relevant to living in violent and chaotic environments (e.g., fights, gangs, drugs, low resources) were integrated into the sessions. In their trial, Cooley-Strickland et al. found no statistically significant program differences compared to waitlist on any of the primary outcome variables including anxiety symptom levels (not even for those children with higher levels of pretest community violence exposure or as a function of the child's sex).

FRIENDS for Life (Aboriginal Children in Canada)

Focusing on Aboriginal children in the West Coast Nations of Canada, I. D. Miller et al. (2011) implemented and tested a culturally adapted FRIENDS for Life protocol. This adaptation focused primarily on cultural enrichments, such as adding a new character (Rusty the Raven) and narrative to facilitate the use of storytelling as part of the protocol. The adaptation also entailed: introducing the craft and use of a medicine pouch decorated with Aboriginal symbols; replacing a support team activity with a circle

of support activity that focused on family, community, culture, and natural and spiritual world resources: and adding a “FRIENDS Wheel” modeled after the medicine wheel. In this study, the FRIENDS program was delivered to elementary school students in Canada (including many Aboriginal children). Findings showed that, for both Aboriginal and non-Aboriginal children, no statistically significant reductions in anxiety levels could be attributed to the intervention.

Acercamiento

Pina et al. (2012) reported on an indicated prevention and early intervention for childhood anxiety trial (about 48% of Latinos who were Spanish speaking and requiring services in Spanish) that included a significant proportion of Mexican-origin families. As such, services were made available in Spanish, and interventionists were trained in a culturally prescriptive protocol (Pina et al., 2009) that calls for personalizing or tailoring program implementation strategies to the unique cultural characteristics of the client. This is achieved by including in the protocol references to norms and values (e.g., *familismo*) that can help the client engage in the use of core program components, such as gradual exposures via motivational procedures. In terms of culture, a focus was on developing cultural enrichments and implementing them prescriptively with high fidelity in the application of standard cognitive and behavioral procedures. The cultural adaptation was conceptualized in terms of program materials and program implementation strategies. In terms of program materials, for example, child and parent handouts were available in Spanish, and cultural enrichments focused on incorporating *dichos* (sayings), proverbs, and refrains that could assist content delivery. In terms of strategies, interventionists were trained both to be attuned to various terminology used by children and families and to use such terminology during the sessions, and to identify “windows of opportunities” to use key cultural values (e.g., *familismo*) potentially relevant to promoting program engagement and practice of program skills, as necessary. Two conditions were compared, “child-only” or “child plus parent,” in Pina et al. (2012), based on knowledge gained from parent focus groups and in-depth interviews showing that Mexican-origin caregivers prefer parental involvement in child-focused interventions (similar findings have been reported by others; Dumka et al., 1998), perhaps because Hispanic/Latinos place great emphasis on caregiving and family. Significant program effects were found in terms of child anxiety symptoms based on a diagnostic interview schedule, parent and child rating scales, and clinically significant change, although the

child-plus-parent condition showed slightly greater gains in terms of child self-reported anxiety and depression levels as well as parent-reported anxiety symptoms about the child. No program moderation effects were found for ethnicity or language (Spanish versus English).

Discussion: Limited Evidence

Despite efforts made to develop culturally robust anxiety and depression prevention programs, overall findings have been somewhat unsuccessful. This appears to be true when regarding African American children in particular (Cardemil et al., 2002; Cooley-Strickland et al., 2011). In contrast, the evidence for Latino youth seems to be more encouraging (Cardemil et al., 2002; Pina et al., 2012). It is important to highlight, however, that the literature is limited to two studies focusing on African American youth and two studies focusing on Latino youth, both exploratory in nature and lacking statistical power for the analyses conducted. For African American youth, deep-level adaptations might be necessary as current findings are not very encouraging. For example, given the importance African Americans place on the family and the caregiving roles grandparents often play, it might be important to enhance anxiety and depression prevention efforts by providing caregivers with substantial PT sessions that focus on evidence-based mediators of child outcomes (e.g., parenting, family functioning variables). Turning to other groups, there was little to no representation in the literature of programs focusing on Asian and/or Pacific Islander children (Barrett et al., 2003; Yu & Seligman, 2002). Barrett et al. (2003), for instance, implemented the FRIENDS program (original nonmodified version) to former Yugoslavian, Chinese, and multi-ethnic children residing in Australia. At posttest, former Yugoslavian youth in FRIENDS showed greater gains than their Chinese counterparts on self-reported anxiety and self-esteem compared to waitlist (along secondary outcomes measures, FRIENDS showed similar cross-ethnic gains including at the 6-month follow-up). Also, there are no anxiety or depression prevention studies focusing on Native American youth, although there is some encouraging research focusing on the prevention of posttraumatic stress reactions in this population (Goodkind, LaNoue, & Milford, 2010).

As we move forward to advance prevention methods focusing on anxiety and depression among ethnic minority children, it would be important to identify program adaptation parameters beyond language and contextually relevant examples. For example, when it comes to Latino children, it would be important to test the extent to which

cultural values such as *familismo* are likely to enhance program effects. Emerging tailoring work by Beardslee and colleagues (D'Angelo et al., 2009) in the area of depression suggests there might be a need to focus on immigration history, adaptation to the host culture, and consideration of key cultural values (trust, respect) during the intervention. However, substantial data about the impact of these types of modifications have not been published to date (D'Angelo et al., 2009; Feinberg et al., 2012). For Asian American children, focusing on cultural values also might hold promise. For instance, Huey and Pan (2006) reported on a small randomized demonstration trial with clinically anxious/phobic Asian American adults. Findings revealed that Asian Americans benefited slightly more from a culturally adapted protocol that considered traditional Asian values compared to a standard protocol. This was true in terms of catastrophic thinking and behavioral approach, two core features of pathological anxiety. Building on these data, it appears that "culture" might definitely need to be considered when targeting anxiety and depression in ethnically and/or culturally diverse children and adolescents. We caution, however, that it might be possible to saturate a program and distract away from critical change processes by infusing too much culture into an intervention (Pina et al., 2009).

HEALTH-FOCUSED INTERVENTIONS

Health disparities for subcultural group children and adolescents are well documented (Flores, 2010; Flores & Tomany-Korman, 2008). In the most extensive analysis of childhood health disparities, Flores (2010) reviewed 111 articles published between 1950 and March 2007. Consistent with the definition provided by the U. S. Department of Health and Health Services, disparities covered by that review included subcultural group differences in disease prevalence, health outcomes, and access to health care. African American, Asian/Pacific Islander, Latino, and Native American children showed disparities relative to non-Hispanic White children in one or more categories of mortality, injuries and illnesses, obesity, physical activity, proper nutrition, and medical care. Culturally adapted interventions often are initiated with the primary intention of reducing disparities by promoting cultural elements that are healthful or modifying those that are associated with health risks. Culture influences health through a variety of mechanisms, such as food preferences, traditional methods of food preparation, celebrations involving food and beverage consumption, perceived desirability of

certain body shapes, consumption of alcohol, and norms for physical activity (MacLachlan, 2006). Understanding how culture influences the health of children informs the development of interventions that incorporate cultural beliefs and practices. By focusing on children's health, cultural adaptations of childhood interventions can not only affect proximal child health outcomes but also can change the trajectory of health outcomes into adulthood. There is a developmental component to some health behaviors and illnesses (such as Type 2 diabetes and cardiovascular disease) that justifies early intervention efforts (Berenson, 2002; Guo & Chumlea, 1999; Magarey, Daniels, Boulton, & Cockington, 2003; Must, Jacques, Dallal, Bajema, & Dietz, 1992; Serdula et al., 1992). Thus, it is worthwhile to develop culturally informed health interventions for children and adolescents that reduce health disparities, incorporate an understanding of culture's influence on health, and disrupt risky health behaviors that threaten to extend from childhood into adulthood.

The Cases of Obesity and Physical Activity

Research on the interrelated topics of obesity and physical activity illustrates the need for culturally adapted interventions and their thorough evaluation. Obesity is a prominent preventable risk factor in the major illness categories of heart disease, cancer, and Type 2 diabetes (Eyre, Kahn, & Robertson, 2004). Unfortunately, across various national studies, there are consistent ethnic group disparities on the prevalence of obesity. For adults, the combined prevalence of overweight and obesity was 64.2%, 76.1%, and 75.8% for non-Hispanic Whites, Blacks, and Mexican Americans, respectively (Wang & Beydoun, 2007). For children, African American, Latino, Native American, and some Pacific Islander subgroups (e.g., Filipinos) show elevated rates of obesity when compared with non-Hispanic Whites (Flores, 2010; Wang & Beydoun, 2007). Wang and Beydoun (2007) reported that the combined prevalence rates for overweight and obesity were 28.2%, 35.4%, and 39.9% for non-Hispanic White, African American, and Mexican American children and adolescents. Disparities were particularly apparent for girls. For instance, in their justification for a targeted preventive intervention, T. N. Robinson et al. (2010) noted that since the 1960s, obesity among African American girls has tripled; they differ from non-Hispanic White girls on body mass index (BMI) as young as age 6, and obesity disparities increase as girls mature into adolescence and young adulthood.

There is a suggestion in the literature that disparities on obesity for African American girls cannot be explained

readily by SES differences (Gordon-Larsen, Adair, & Popkin, 2003). Utilizing data from a nationally representative sample of U. S. adolescents, one report focused on a subsample of over 13,000 that included non-Hispanic White, Hispanic, African American, and Asian American males and females who were 12 to 20 years of age. Ethnic-racial disparities were evident in the prevalence of overweight girls: Asian American: 10.37%; non-Hispanic White: 22.24%; Hispanic: 29.59%; and African American: 37.94%. It was particularly interesting that there was a negative relation between overweight status and SES for non-Hispanic White and Hispanic girls but not for African American girls, whose prevalence of being overweight was relatively elevated at higher levels of SES. For African American girls, it appears that factors such as income and education might increase risk rather than lessen risk as they do for other subcultural groups. Subcultural group differences in risk and protective factors are intriguing because they suggest possible differences in modifiable variables that could be included in culturally adapted interventions for childhood obesity (Barrera et al., 2013).

One chronic illness associated with obesity, Type 2 diabetes, disproportionately affects African Americans, Latinos, and Native Americans (Cowie et al., 2006; Fagot-Campagna et al., 2000; Gahagan & Silverstein, 2003) and is a leading cause of death, blindness, heart disease and amputations (Wong et al., 2005). Particularly disturbing are the estimates of the prevalence of diabetes for the next generation, an anticipated development largely attributable to the rise in obesity. Forecasts suggest that for those born in the year 2000, 45.4% of Latino men, 52.5% of Latino women, 40.2% of Black men, and 49% of Black women will be diagnosed with diabetes within their lifetimes (Narayan, Boyle, Thompson, Sorensen, & Williamson, 2003). Those projections are substantially higher than projections for White men (26.7%) and women (31.2%), which are, nevertheless, still disturbing in their magnitude.

Physical Activity

Comparative research on the physical activity of ethnic minority children is less developed than research on obesity (Flores, 2010). When compared with non-Hispanic White youth, African American and Hispanic youth had lower aerobic fitness, less vigorous physical activity among girls, and more television viewing during weekdays (Flores, 2010). Those conclusions were based primarily on findings from the well-known Monitoring the Future annual surveys (Delva, O'Malley, & Johnston, 2006). The report by Delva, O'Malley, and Johnston (2006) used surveys from 1993–19–94 to 2001–2003 for eighth and tenth

graders and surveys from 1986–1988 to 2001–2003 for twelfth graders. Results indicated that for boys, only one of 14 grade-by-survey year comparisons on frequency of vigorous physical activity showed ethnic/racial differences. For girls, the results were drastically different. African American and Hispanic girls reported less vigorous physical activity than non-Hispanic White girls on all annual surveys and for all grade levels. For hours of TV viewing, African American and Hispanic boys and girls reported more TV viewing than non-Hispanic White boys and girls on all annual surveys and for all grade levels. These findings indicate the clear need for interventions that are designed to increase physical activity and decrease sedentary behavior among African American and Hispanic children and adolescents.

Exemplary Research Programs Integrating Culture and Children's Health

Several model health interventions illustrated extensive and systematic cultural adaptation efforts. In this section we highlight three that are exemplary in the procedures they implemented to become informed about relevant cultural considerations, integrate those considerations into intervention designs, and then evaluate adapted interventions' engagement and efficacy. Other important controlled studies on interventions designed to improve the diet and physical activity of ethnocultural group children have been conducted, but published reports lack detailed descriptions of cultural adaptation procedures (Crespo et al., 2012; J. N. Davis et al., 2012; de Heer, Koehly, Pederson, & Morera 2011; Klesges et al., 2010; Nadar et al., 1999; Resnicow, Taylor, Baskin, & McCarty, 2005; Treviño et al., 2004; D. K. Wilson et al., 2011).

Stanford GEMS

The Stanford GEMS (Girls health Enrichment Multi-site Studies) was a culturally adapted after-school dance program and an in-home sedentary behavior (television and other screen viewing) reduction intervention for African American girls (T. N. Robinson et al., 2010). Stanford GEMS emphasized physical activity as a means for body weight reduction. The research team described their programmatic research in key articles that contained literature reviews, theoretical considerations, pilot research, and a randomized controlled trial efficacy study (Kumanyika et al., 2003; T. N. Robinson et al., 2003, 2008, 2010). Many features of exemplary cultural adaptation procedures (Barrera et al., 2013) were illustrated in the development of Stanford GEMS.

An initial report described a literature review that identified special considerations for obesity interventions directed at African American youth (T. N. Robinson et al., 2003). The authors noted that because African American girls, family members, and men might be less concerned about thinness than White counterparts, African American girls and women might be less motivated to achieve distal goals of weight loss and fitness. They surmised that intervention activities might need to be intrinsically and immediately appealing to sustain engagement of African American girls. The authors explained the considerations they made in identifying dance as a gratifying, culturally appropriate after-school physical activity. They conducted other extensive formative research that they summarized in this way: “The treatment and control interventions were developed through extensive formative research, including focus groups with girls; focus groups with parents/guardians; written questionnaires; interviews of community youth workers and community leaders; and testing the feasibility of the dance classes and TV reduction lessons with small groups of girls and individual families” (T. N. Robinson et al., 2003, S1–67 to S1–77.).

Cultural adaptations included surface structure elements, such as culturally matched participants and staff and participant-selected music and dance as the featured physical activity. The intervention developers indicated that deep structure elements associated with African American culture (e.g., collectivism, importance of family, present orientation, religiosity, sense of historical racism and prejudice, and use of social support) were infused into the intervention. The early cultural adaptation phases of Stanford GEMS contained a 12-week pilot study (T. N. Robinson et al., 2003), another feature of high-quality adaptation efforts (Barrera et al., 2013). Goals of the pilot were to evaluate the recruitment strategy and participants’ satisfaction with intervention activities’ and the intervention’s effects on BMI, physical activity, screen (TV, video) viewing, and other outcomes. Sixty-one African American girls age 8 through 10 were recruited through community centers and after-school programs in low-income neighborhoods of Oakland and East Palo Alto California. To be eligible, girls had to have a BMI that was greater or equal to the fiftieth percentile for their age group and/or one overweight parent or guardian with a BMI of greater or equal to 25. Girls who were randomly assigned to the active intervention could attend dance classes that were offered 5 days per week at neighborhood community centers. They could attend dance classes as often as they liked for the 12 weeks of intervention. In addition, intervention participants received 5 in-home lessons that were designed

to reduce TV and video viewing. Participants assigned to the control condition received a health education program that emphasized healthful eating and physical activity. They also received newsletters containing information about health risks that might lead to obesity, heart disease, and diabetes.

The pilot showed that the researchers were successful in enrolling more than their targeted number of participants, and participants met eligibility criteria. They also retained all but one of the participants over the 12 weeks of the pilot. Satisfaction with the intervention and participation in it were very good with one exception. The lack of affordable transportation in one community resulted in less than half the dance class participation that was achieved in the other community. Due to a lack of power, the intention was to estimate effect sizes rather than to show statistically significant intervention effects. Encouraging effect sizes were found for BMI, waist circumference, physical activity, and TV viewing that favored the intervention condition. Overall, the pilot study demonstrated the feasibility, appeal, and potential efficacy of the combined dance and TV-viewing-reduction intervention.

Ultimately, the Stanford GEMS team conducted a 2-year randomized controlled trial with 261 African American girls and their parents or guardians (T. N. Robinson et al., 2010). As in the pilot, participating girls were between the ages of 8 and 10 and were either overweight or had at least one parent/guardian who was overweight. Participants randomly assigned to the active intervention had access to after-school dance classes 5 days each week for 2 years and received up to 24 in-home sessions for reducing screen viewing. Over the 2-year period, the control condition included health education, monthly newsletters, and quarterly lectures at community centers. Despite the exemplary methods for creating a culturally adapted intervention, extending the intervention period to 2 years, and attracting an adequate number of participants, there was not a statistically significant treatment effect on the primary outcome measure, BMI. However, there were statistically significant intervention effects on total cholesterol, low-density lipoprotein cholesterol, and depression. In tests of moderation when BMI was the outcome variable, the intervention was more effective when parents or guardians were unmarried and when baseline television viewing was high. The research team suspected that treatment effects were not stronger because of barriers participants encountered in attending dance classes due to transportation difficulties and episodes of violent crime near the sites where dance classes were held.

Hip-Hop to Health Jr.

A second exemplary program of research, Hip-Hop to Health Jr., was a community-based obesity prevention program that was culturally adapted for African American and Latino Head Start preschool children (Fitzgibbon et al., 2002). It was an adaptation of Hip-Hop to Health, an intervention for African American 6- to 10-year-old children (Fitzgibbon et al., 1998). In an article that described the rationale and design of Hip-Hop to Health Jr., the research team reviewed the literature on diet and exercise of African American and Latino children to identify specific characteristics (e.g., fat and fiber consumption) that might account for disparities in childhood and adult obesity. They also explained that the theoretical framework guiding intervention development was an integration of social learning theory (Bandura, 1986), self-determination theory (Deci, 1992; Deci, Driver, Hotchkiss, Robbins, & Wilson, 1993), and the transtheoretical model (Prochaska & DiClemente, 1992). Consistent with the guidelines for cultural adaptation phases (Barrera et al., 2013), this first phase of information gathering included a literature review.

Also consistent with adaptation guidelines, the Hip-Hop to Health Jr. research team conducted a 3-week pilot study to test the feasibility and appeal of both the parent and child intervention components. The child portion of the pilot intervention contained three 40-minute sessions per week. The first half of each session was devoted to nutrition, and the second half was devoted to physical activity. In the nutrition lessons, “go and grow” foods (healthful) were contrasted with “slow” foods (unhealthful). In the parent component, parents attended one 30-minute nutrition class and two 30-minute aerobics classes each week.

Several changes were made to the intervention following the pilot study. The distinction between “go and grow foods” and “slow foods” was difficult for preschool children to understand. For the revised intervention, researchers created puppets to represent the food groups (Ms. Grain, Ms. Fruit, Mr. Vegetable, Mr. Fat, Ms. Sugar) in nutrition lessons. They also recognized the need for greater emphasis on the importance of physical activity. In the pilot parent component, nutrition lessons were attended poorly. Based on focus group recommendations that followed the pilot study, weekly newsletters and homework assignments replaced the in-person nutrition lessons in the revised parent intervention component.

A revised intervention was evaluated in two RCTs, one conducted with primarily African American families (Fitzgibbon et al., 2005) and one conducted with primarily Latino families (Fitzgibbon et al., 2006). The intervention was provided for 14 weeks. Children received 20 minutes

of nutrition instruction (utilizing puppets) and 20 minutes of physical activity three times each week. Parents received weekly newsletters that were coordinated with the content of their children’s instruction. In addition, parents could participate in 30-minute aerobic classes twice each week at their children’s Head Start facilities.

The authors summarized the five special considerations they made in developing the intervention for African American and Latino preschoolers and their parents:

1. Offer the sessions in locations that are safe and that can be reached easily.
2. Establish personal relationships between participants and intervention staff.
3. Reduce barriers (e.g., negative attitudes, competing responsibilities) that interfere with improvements in nutrition and exercise.
4. Provide behavioral demonstrations of strategies for lifestyle change.
5. Make intervention accessible for participants at all levels of literacy.

The RCTs found significant intervention effects in predominantly African American Head Start centers (Fitzgibbon et al., 2005) but not in predominantly Latino centers (Fitzgibbon et al., 2006). In African American centers, children in the intervention condition showed smaller BMI increases than children in the control condition 1 year and 2 years after the end of the intervention, and less saturated fat consumption at 1-year follow-up. In contrast, there were no significant intervention effects for children and parents who attended primarily Latino Head Start centers. The researchers noted that children in the Latino centers had higher BMI levels at baseline than children in the African American centers. They speculated that relatively high initial levels of overweight and obesity might have made it difficult for children in the Latino centers to modify their eating and weight. Overall, they recognized that additional changes were needed to assist unacculturated Latino parents in making changes to diet and physical activity. On a positive note, retention of participants was quite high and did not differ between intervention and control groups. The retention rates were 97%, 86%, and 85%, at posttest, Year 1, and Year 2 follow-ups, respectively.

Pathways

Pathways was a multifaceted, multisite obesity prevention program for American Indian schoolchildren, which

started in grade 3 and continued through grade 5. It was developed and evaluated in two 3-year phases. The first 3-year phase was used to systematically develop culturally appropriate intervention and assessment methods. An entire supplemental issue of the *American Journal of Clinical Nutrition* (1999) was devoted to detailed descriptions of the formative research (e.g., S. M. Davis et al., 1999; Gittelsohn et al., 1999; Teufel et al., 1999). An additional 3 years was required to conduct a randomized controlled trial to assess intervention effects on body fat, dietary fat intake, physical activity, knowledge of the diet and physical activity curriculum, and self-efficacy for physical activity and healthful food choices (Caballero et al., 2003). Articles in a special supplemental issue of *Preventive Medicine* (2003) reported detailed results of the massive project.

The intervention consisted of four components: a classroom curriculum, school food services, physical activity during and after school, and family support and involvement. Formative research groups were created to develop each of the components. Unusually thorough input from many constituencies was obtained throughout intervention development. Each section of the intervention was reviewed by “tribal representatives, American Indian members of the Pathways study, intervention committee members, and the Pathways steering committee, composed of the principal investigators from all five participating institutions, the National Heart, Lung, and Blood Institute project scientist, and two American Indian representatives” (S. M. Davis et al., 1999, p. S798).

They instituted an information gathering phase when information relevant for understanding the communities, risk factors, and potential intervention strategies were obtained from school staff members, students, and their caregivers in detailed interviews, focus groups, and direct observation. That information informed a culturally adapted intervention that incorporated American Indian concepts and traditions pertinent to nutrition and physical activity. For example, the classroom component integrated “stories, games, music, artwork, foods, family activities, videos, and events contributed by members of the seven tribes” (S. M. Davis et al., 2003, p. S26). An initial version of the intervention was pilot tested in six American Indian communities. Observers made notes of the intervention’s implementation, and researchers interviewed intervention staff members (e.g., teachers, food service workers) to ascertain strengths, weaknesses, and recommendations for changes. A revised intervention was evaluated in a randomized controlled trial of 1704 children in 41 schools in seven American Indian communities in Arizona, New Mexico, and South Dakota (Caballero et al., 2003). The

intervention began in third grade and extended to fifth grade. Retention from baseline to follow-up assessment was 83% in both the intervention and control conditions.

Despite rigorous cultural adaptation methods, the comprehensive intervention was not more successful than the control condition in reducing body fat, the primary outcome measure. Motion sensor measurement of physical activity also did not show significant intervention effects. Significant intervention effects were found for self-reported and observed dietary fat intake and for self-reported physical activity. In discussing their results, the researchers noted that several other comprehensive interventions used diet and physical activity outcomes but did not treat children’s body weight and adiposity as endpoints (e.g., Luepker et al., 1996; Simons-Morton, Parcel, Baranowski, Forthofer, & O’Hara, 1991).

Stanford GEMS, Hip-Hop to Health Jr., and Pathways were exemplary in their illustration of cultural adaptation procedures. As noted previously, several other studies on the nutrition and/or physical activity of ethnocultural group children did not provide substantive descriptions of cultural adaptation considerations and procedures. For instance, Bienestar was a large-scale study that involved 13 intervention elementary schools and 14 control schools in which 713 fourth-grade students participated in the intervention and 706 students participated in the control condition (Treviño et al., 1998, 2004; Treviño, Hernandez, Yin, Garcia, & Hernandez, 2005). Eighty percent of the participants were of Mexican heritage. Much like Pathways’ comprehensive intervention, the Bienestar comprehensive intervention involved four social systems: parents, school classrooms, school cafeterias, and after-school programs. However, unlike Pathways, it is not certain if formative research was funded to conduct systematically a cultural adaptation of the intervention. The other studies of nutrition and/or physical activity interventions that were directed at subcultural group children (Crespo et al., 2012; J. N. Davis et al., 2012; de Heer et al., 2011; Klesges et al., 2010; Nadar et al., 1999; Resnicow et al., 2005; D. K. Wilson et al., 2011) might have included cultural adaptation phases, but detailed descriptions of adaptation methods did not appear in the literature we were able to locate and review. Many of these studies showed significant intervention effects on physical activity and healthful nutrition measures (Crespo et al., 2012; de Heer et al., 2011; Nadar et al., 1999; Treviño et al., 2004).

Discussion

The three exemplary studies we highlighted—Stanford GEMS, Hip Hop to Health Jr., and Pathways—were

pioneers in using intervention development procedures that we now recognize as high-quality features of cultural adaptation stages: information gathering with literature searches to document health disparities and to identify subcultural risk and protective factors that should be considered in intervention development; formative studies that include focus groups as well as quantitative surveys; pilot studies to test the feasibility and effects of a draft intervention; satisfaction surveys to gauge the appeal of recruitment and intervention procedures; and formal RCTs of adapted interventions. The exemplary projects benefited from funded, multiyear developmental stages that permitted thorough cultural adaptation efforts.

Despite those systematic efforts, culturally adapted interventions were not always successful in changing primary endpoints of BMI or adiposity (see Fitzgibbon et al., 2005, for an exception). Those outcomes are notoriously resistant to change, particularly in primary prevention interventions with nonclinical populations (Caballero et al., 2003). In a meta-analysis of school-based obesity interventions in the United States and United Kingdom, Kanekar and Sharma (2009) found that the interventions were not successful in reducing BMI. Similarly, a review of 20 after-school obesity prevention interventions concluded that the results were generally unfavorable when obesity-related variables were the outcomes (Branscum & Sharma, 2012). Nevertheless, culturally adapted interventions were often successful in improving secondary (yet important) outcomes, such as dietary fat consumption, fiber consumption, and physical activity. Childhood obesity and its subcultural group disparities will persist for some time. They have such critical importance for the nation's health that research must continue to seek solutions that will reach children in all segments of communities.

One of the lessons from the Stanford GEMS project (T. N. Robinson et al., 2010) is that well-conceived interventions must include effective methods for overcoming barriers to participation in intervention components. The GEMS' goal for attendance at after-school dance classes in community centers was three of five school days, but by the end of the intervention, participants averaged only one of five school days. Of 24 in-home TV reduction visits, participating families received an average of 12.4 lessons over a 2-year period. In-school interventions avoid the participation slippage that occurs when children are required to travel to intervention venues or when family participation is required. However, true lifestyle change occurs during school hours and after school hours when the family has a key role in influencing food consumption and physical

activity. Cultural adaptation efforts should continue to address both engagement methods and intervention content (Barrera & Castro, 2006).

CONCLUSIONS AND FUTURE RESEARCH DIRECTIONS

The cultural adaptation of interventions for youth is a maturing and still-vibrant topic in developmental psychopathology. There is a persistent need for interventions to reach and engage ethnic and racial minority youth just as there are persistent disparities in youth problem behavior and health risk. This chapter has highlighted the most influential work on cultural adaptations that has been done to date, leading to several conclusions about what we have learned, limitations to our knowledge, and ways we might advance our understanding in the near future. Basic conclusions and future research directions are integrated in the next sections.

Adaptation Effectiveness

In this chapter, as well as in other reviews (Benish, Quintana, & Wampold, 2011; Griner & Smith, 2006; Huey & Polo, 2008; Ortiz & Del Vecchio, 2013), the emphasis was on evaluating intervention efficacy. Are cultural adaptations effective? That question should be qualified by another question that has been raised repeatedly: Compared to what? There is interest in the question not only of the efficacy of cultural adaptations relative to benign control conditions but also relative to unadapted, recognized interventions, such as the original versions of interventions. For some, demonstrating the value added by modifications to original EBIs is of critical importance (Ortiz & Del Vecchio, 2013). Yet calls for research that compares the effectiveness of culturally adapted EBIs to their original versions are qualified by skepticism: It is hard to imagine providing unaltered original EBIs to ethnocultural groups without making basic adaptations, such as language translations or delivery by bilingual staff; and it would be costly to launch an adequately powered study comparing an original EBI, a culturally adapted intervention, and a control condition, particularly if differences between the original and adaptation are expected to be modest (Barrera et al., 2013). Lau, Chang, Okazaki, and Bernal (in press) questioned the value of original EBI-adaptation comparisons, also citing the substantial statistical power required for such comparisons, the need for at least some cultural adaptations to effectively engage

ethnocultural participants into comparative research, and potential ethical concerns about withholding culturally sensitive care from participants who receive unadapted EBIs. Given the onerousness, methodological challenges, and ethical quandaries of conducting adequately powered relative efficacy trials of adapted EBIs (Cardemil, 2010), Domenech Rodriguez and Bernal (2012) noted that researchers are subject to a catch-22 when expected to demonstrate relative efficacy prior to pursuing questions about mediators and moderators of change within the clinical trials tradition. From our qualitative review of culturally adapted interventions for internalizing disorders, externalizing and high-risk youth behaviors, health, and PT, there was good evidence of absolute intervention effects when compared to control conditions. However, substantial heterogeneity was observed in the efficacy of culturally adapted EBIs across the diverse outcomes that have been studied. Although some outcomes (e.g., obesity) have been notoriously intractable to remediation or prevention even for majority group youth, it was still disappointing when well-conceived, systematic, and community-informed adaptation efforts were unsuccessful in achieving goals.

Our analysis of the literature was qualitative and not intended to be exhaustive or capable of supplying quantitative answers to questions of relative efficacy. Fortunately, four meta-analyses of culturally adapted interventions (Benish et al., 2011; Griner & Smith, 2006; Huey & Polo, 2008; Smith, Domenech-Rodriguez, & Bernal, 2011) and a formal review of PT interventions (Ortiz & Del Vecchio, 2013) have been published. There was some overlap in the studies that were analyzed in these reviews, but the reviews differed in their inclusion and exclusion criteria. Using relatively broad inclusion criteria, Griner and Smith (2006) evaluated 76 published and unpublished studies that made at least some reference to adaptation because of culture, ethnicity, or race. Only studies of EBIs with children were reviewed by Huey and Polo (2008). Psychotherapy studies conducted with children or adults (excluding preventive interventions) were the foci of the meta-analysis by Benish et al. (2011). That review also contrasted effect sizes that were obtained when culturally adapted interventions were compared to some other treatments or benign control conditions and when cultural adaptations were compared to just other bona fide psychotherapies. Bona fide psychotherapies were defined by “a relationship with a therapist tailored to the client and is evidenced by the inclusion of two or more of the following conditions: use of an established psychotherapy approach, incorporation of the use of psychological processes, use of a manual or

training as a guide for therapists, and/or identification of the active ingredients of the therapy” (Benish et al., 2011, p. 280). Smith et al. (2011) conducted a meta-analysis of studies concerned with mental health outcomes. That review excluded research on health, substance abuse, and educational outcomes. Ortiz and Del Viccho’s (2013) review dealt exclusively with PT interventions. Because of differences in scope and reference groups used in comparisons, it is understandable that there were differences in conclusions regarding adaptation efficacy.

The five most basic findings from these reviews are summarized next.

1. Overall, culturally adapted interventions were moderately effective (average effect size was $d = .45$) but were less effective for children than they were for adults (Griner & Smith, 2006).
2. For interventions conducted with ethnic minority children, culturally adapted treatments and unadapted treatments showed comparable effect sizes (Huey & Polo, 2008).
3. On mental health outcomes, culturally adapted therapies showed a moderate effect size (average effect size $d = .46$) (Smith et al., 2011). Similar to the age effect found by Griner and Smith (2006), participants older than 35 appeared to benefit more from adapted therapies compared with benefits for children, adolescents, and young adults. The number of cultural adaptations made in interventions was positively correlated (.28) with intervention efficacy.
4. Analyses comparing culturally adapted psychotherapies to other bona fide (unadapted) psychotherapies showed an average effect size of .32. In contrast to other reviews (Griner & Smith, 2006; Smith et al., 2011), the average effect size was comparable for adults and children (Benish et al., 2011).
5. In studies of parent training, the reviewers concluded that “treatment outcomes from adapted PT interventions are not superior to those unadapted interventions typically achieve” (Ortiz & Del Vecchio, 2013, p. 452).

In general, cultural adaptations show moderate effect sizes when compared with control conditions, but there was some evidence that effects might be attenuated for research conducted with children and adolescents. Age might moderate effects if cultural adaptations are most beneficial for the least acculturated participants who are often older adults rather than children (Griner & Smith, 2006). It was notable, however, that no age effects were observed in

what was arguably the most rigorous comparison, that is, when adapted interventions were found to be moderately more effective than unadapted bona fide therapies (Benish et al., 2011).

Two additional observations provide important context for interpreting these findings. First, there was considerable heterogeneity in treatment effects, which calls for caution in making broad generalizations about efficacy. Second, some analyses were based on a small number of studies. We agree with Smith et al.'s (2011) conclusion that because the full assessment of efficacy "depends on the consistent replication of supportive findings, the single greatest need of the research on culturally adapted interventions is for more evidence to accumulate" (p. 171). They noted that a consistent rate of about two studies each year was produced since 1981 that met the inclusion criteria for their meta-analytic review. Even Ortiz and Del Vecchio (2013), who questioned the evidence supporting the value of cultural adaptations, acknowledged that ethnic minority children and families (particularly Hispanics) were underrepresented in controlled, peer-reviewed studies of parent training.

Cultural Adaptation Process

Researchers and program developers turn to the literature to inform decisions about pursuing cultural adaptations and to learn strategies for conducting them. That literature shows large variability in the thoroughness of adaptation descriptions—descriptions ranging from cryptic accounts of seemingly modest tailoring efforts to extensive efforts that required several years to craft interventions that subsequently were evaluated in controlled trials. Multiyear projects to culturally adapt children's health interventions (e.g., T. N. Robinson et al., 2003) were innovative during their time, implementing sequential steps that are now reflected in the stage models listed in Table 17.1 for developing culturally adapted versions of EBIs. We have learned that when EBIs are identified for modifications, there are generally accepted procedures for conducting them. An alternative approach to tailoring EBIs was exemplified with approaches that began with basic developmental psychopathology research to identify the risk factors that appear to contribute to psychopathology or youth problem behaviors (e.g., Gonzales et al., 2007; Murry et al., 2007). In the future, greater structure in the procedures used to culturally adapt interventions might decrease the heterogeneity in effects that were found in our review and recent meta-analyses. Many of the studies included in reviews were conducted prior to published conceptual models of adaptation procedures.

Reach and Engagement

Lau (2006) made the distinction between the ability of interventions to change outcomes, such as disability and distress, and their ability to draw participants into interventions, engage them in intervention activities, and retain them throughout treatment and maintenance phases. As shown in this review, the effects of cultural adaptations on reach and engagement have not been studied and reported as systematically as intervention efficacy, yet there are examples of how tailoring can improve engagement. Kumpfer and Alvarado (1995) found that original versions of Strengthening Families had slightly better outcomes than adapted versions. However, recruitment and retention of participants receiving culturally adapted versions were 41% better, instantiating the potential of cultural adaptations to substantially improve involvement in interventions for ethnic minority families. Unfortunately, there are countervailing examples of well-conceived efforts to boost the intervention engagement of ethnic minority youth and parents that were not successful (e.g., McCabe et al., 2012; T. N. Robinson et al., 2010). The exigencies of living in stressful, low-resource environments will continue to present barriers to ethnic minority participants in intervention research. Developing and evaluating methods for improving reach, engagement, and sustained participation in interventions are among the most important topics for future research.

Understanding Cultural Components Added to Culturally Adapted Interventions

Several reviewers were critical of cultural adaptations that lacked theoretical frameworks for specifying why cultural concepts could be added to interventions to improve reach, engagement, or efficacy (Castro et al., 2010; Ortiz & Del Vecchio, 2013; B. D. M. Wilson & Miller, 2003). Little progress has been made on this front. When cultural adaptations are explicitly designed to influence a cultural construct that could operate as a mediator of change (e.g., enhancing cultural identity), studies could inform culturally relevant theory by evaluating whether the intervention succeeded in changing that construct and whether that change affected the outcome (e.g., Murry et al., 2007). Toward this important aim, the designers of culturally relevant interventions should be explicit about the hypothesized roles of specific cultural variables and should design evaluation research that tests theoretical assertions. This approach was illustrated in the study by Lau et al. (2011) that included special therapy procedures to reduce negative parenting strategies of immigrant Chinese parents.

Problematic parenting practices identified in preliminary research (Lau, 2010) were improved in a culturally adapted version of an EBI, and those improvements mediated the intervention's beneficial effects on children's externalizing problems (Lau et al., 2011). Murry et al. (2007) provided a mediation analysis of the Strong African American Families Program that could serve as another model for future research on the cultural mechanisms that mediate the effects of cultural adaptations.

Accommodating Within-Group Diversity

A major impetus for conducting cultural adaptations is to address special needs and characteristics of ethnocultural groups that were not addressed adequately by the original versions of the interventions. However, ethnocultural groups are not homogeneous. Within-group diversity is derived from differences in acculturation, nativity, language use, immigration experiences, skin color, religion, and a host of other factors that might be related to adversity exposure and responsiveness to interventions. For example, Gonzales and colleagues (2012) reported differential program effects on the basis of family linguistic acculturation, yet program benefits were nonetheless shown over time for both low- and high-accultured families (Jensen et al., in 2014). A complexity in designing culturally adapted interventions is to accommodate within-group diversity. It is not feasible to create fixed interventions that are relevant and effective for narrow segments of an ethnocultural group. Instead, there is promise in creating assessment and intervention procedures that guide individualized tailoring within the context of a standardized intervention. Standardized decision rules for varying the content and dosage of treatment, depending on the characteristics of participants is the essence of adaptive intervention designs (Collins et al., 2004). In the same spirit of adaptive interventions, Pina et al. (2009) described a culturally prescriptive intervention framework that guides the tailoring of childhood anxiety interventions depending on the cultural features of the individual child or family. Rather than applying a fixed set of adaptation activities, their treatment manual describes a uniform set of guidelines to assist interventionists in determining how to attend to language and other cultural considerations (e.g., familism) in tailoring the intervention. McCabe et al.'s (2005) use of an initial assessment of participants' attitudes and preferences to guide the individualized application of treatment elements was innovative and merits additional research and development as a strategy for addressing within-group diversity.

International Application and Dissemination

It could be said that the strongest test of the need for cultural adaptation may come from international dissemination of interventions to non-Western and developing nations. The need for linguistic adaptations and other cultural considerations is likely to be magnified in international applications with non-Western developing countries in Africa, Asia, and the Middle East. Earlier in the chapter we noted the emerging trend for international dissemination of EBI, including several PT interventions that have been adapted for international dissemination. The same has been true of programs to prevent HIV/AIDS (Wingood & DiClemente, 2008). Undoubtedly, the best examples of international adaptations were concerned with the SFP, a family skills intervention for the prevention of substance abuse (Kumpfer et al., 2008). SFP was initially developed and tested in the United States and subsequently evaluated in non-RCT studies in about 17 countries and RCTs in nine countries (United States, Canada, Australia, United Kingdom, Sweden, the Netherlands, Spain, Italy, and Thailand). Kumpfer and her colleagues (2008, 2012) detailed 10 steps shown in Table 17.1 that should be followed in creating international adaptations of SFP and perhaps other evidence-based programs. In Kumpfer et al. (2008), the steps were illustrated with examples from their extensive experience in many countries. That informative paper is filled with general guidance and best practices for those who are planning international adaptations of EBIs.

International adaptation of interventions is a prime topic for future research and practice. However, in their recent review of PT interventions that have been implemented and evaluated internationally, Knerr et al. (2013) noted that the four PT interventions with the strongest evidence base have been replicated in RCTs in many developed Western countries, but these major programs have not been implemented in low-income or developing countries. On the contrary, many PT interventions transported to these contexts lacked strong efficacy data in the countries in which they originated. Thus, it is unclear whether international efforts to deploy PT or other types of interventions will be informed by domestic efforts in cultural adaptation.

It is important to note that although some EBIs are gaining traction in usual care settings, and the recent federal emphasis on prevention services as outlined in the Affordable Care Act (U.S. Department of Health and Human Services, 2010) has potential to accelerate this pattern, there is little evidence that their culturally adapted counterparts have been adopted outside the original efficacy trials and the occasional dissemination trials.

Zayas et al. (2012) argued that one of the shortcomings in the cultural adaptation literature is the inattention to the multiple service contexts that influence the implementation of interventions and the critical features that enhance the uptake of interventions and services. Larger contextual factors, in addition to sociocultural factors, must be considered in the adaptation process, such as the different service sectors (e.g., child welfare, education, juvenile justice, medical) and the organizational contexts (e.g., agencies, hospitals, clinics, health care) in which interventions are implemented, and the types of modifications required at different levels of the service delivery and implementation chain. It is possible that, in each context, cultural adaptations must be conceptualized and operationalized differently (Schoenwald 2008). An integrated approach to cultural adaptation, which includes the careful consideration of both the service contexts as well as the cultural diversity represented among the families and children within these contexts, may be necessary (Wolchik et al., 2009; Zayas et al., 2012). As the field of child, youth, and family prevention services pushes forward to solve the problem that EBIs have not been adopted broadly and are not accessible, scalable, replicable, and sustainable, intervention adaptation will be a major thrust in the next generation of efforts to design and diffuse EBIs (Rotheram-Borus, Swendeman, & Chorpita, 2012). The role that culture and the cultural adaptation processes described in this review will play in this agenda has yet to be determined.

REFERENCES

- Addis, M. E., & Cardemil E. V. (2006). Does manualization improve therapy outcomes? Psychotherapy manuals can improve outcomes. In J. C. Norcross, L. E. Beutler, & R. F. Levant (Eds.), *Evidence-based practice in mental health: Debate and dialogue on the fundamental questions* (pp. 131–140). Washington, DC: American Psychological Association.
- Aktan, G., Kumpfer, K. L., & Turner, C. W. (1996). Effectiveness of a family skills training program for substance use prevention with inner city African-American families. *Substance Use and Misuse, 31*, 157–175. doi: 10.3109/10826089609045805
- Alegria, M., Lin, J. Y., Green, J. G., Sampson, N. A., Gruber, M. J., & Kessler, R. C. (2012). Role of referrals in mental health service disparities for racial and ethnic minority youth. *Journal of the American Academy of Child & Adolescent Psychiatry, 51*, 703–711. doi: 10.1016/j.jaac.2012.05.005
- Anderson, N. L., Koniak-Griffin, D., Keenan, C. K., Uman, G., Duggal, B. R., & Casey, C. (1999). Evaluating the outcomes of parent-child family life education. *Research and Theory for Nursing Practice, 13*, 211–234.
- August, G. J., Lee, S. S., Bloomquist, M. L., Realmuto, G. M., & Hektner, J. M. (2003). Dissemination of an evidence-based prevention innovation for aggressive children living in culturally diverse, urban neighborhoods: The Early Risers effectiveness study. *Prevention Science, 4*(4), 271–286. doi: 10.1023/A:1026072316380
- Bakken, R. J., & Winter, M. (2002). Family characteristics and sexual risk behaviors among black men in the United States. *Perspectives on Sexual and Reproductive Health, 34*, 252. doi: 10.2307/3097824
- Balcazar, H., Castro, F. G., & Krull, J. L. 1995. Cancer risk reduction in Mexican American women: The role of acculturation, education, and health risk factors. *Health Education Quarterly, 22*, 61–84. doi: 10.1177/109019819502200107
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. *Psychological Review, 84*(2), 191. doi: 10.1037/0033-295X.84.2.191
- Bandura, A. (1986). *Social foundations of thought and action*. Englewood Cliffs, NJ: Prentice Hall.
- Baptiste, D. R., Bhana, A., Peterson, I., McKay, M., Voisin, D., Bell, C., & Martinez, D. D. (2006). Community collaborative youth-focused HIV/AIDS prevention in South Africa and Trinidad: Preliminary findings. *Journal of Pediatric Psychology, 31*, 905–916. doi: 10.1093/jpepsy/31.10
- Baptiste, D. R., Voisin, D. R., Smithgall, C., Martinez, D. D. C., & Henderson, G. (2007). Preventing HIV/AIDS among Trinidad and Tobago teens using a family-based program: Preliminary outcomes. *Social Work in Mental Health, 5*(3–4), 333–354. doi: 10.1300/J200v05n03_05
- Barker, C. H., Cook, K. L., & Borrego, J., Jr. (2010). Addressing cultural variables in parent training programs with Latino families. *Cognitive and Behavioral Practice, 17*(2), 157–166. doi: 10.1016/j.cbpra.2010.01.002
- Barrera, M., Jr., & Castro, F. G. (2006). A heuristic framework for the cultural adaptation of interventions. *Clinical Psychology: Science and Practice, 13*, 311–316. doi: 10.1111/j.1468-2850.2006.00043.x
- Barrera, M., Jr., Castro, F. G., & Holleran Steiker, L. K. (2011). A critical analysis of approaches to the development of preventive interventions for subcultural groups. *American Journal of Community Psychology, 48*, 439–454. doi: 10.1007/s10464-010-9422-x
- Barrera, M., Jr., Castro, F. G., Strycker, L. A., & Toobert, D. J. (2013). Cultural adaptation of behavioral health interventions: A progress report. *Journal of Consulting and Clinical Psychology, 81*, 196–205. doi: 10.1037/a0027085
- Barrera, M., Jr., Prelow, H. M., Dumka, L. E., Gonzales, N. A., Knight, G. P., Michaels, M. L., . . . Tein, J.-Y. (2002). Pathways from economic conditions to adolescent distress: Supportive parenting, stressors outside the family and deviant peers. *Journal of Community Psychology, 30*, 135–152. doi: 10.1002/jcop.10000
- Barrett, P. M., Farrell, L. J., Ollendick, T. H., & Dadds, M. (2006). Long-term outcomes of an Australian universal prevention trial of anxiety and depression symptoms in children and youth: An evaluation of the Friends Program. *Journal of Clinical Child and Adolescent Psychology, 35*, 403–411. doi: 10.1207/s15374424jccp3503_5
- Barrett, P. M., Sonderegger, R., & Xenos, S. (2003). Using friends to combat anxiety and adjustment problems among young migrants to Australia: A national trial. *Clinical Child Psychology & Psychiatry, 8*, 241–260. doi: 10.1177/1359104503008002008
- Barrett, P., & Turner, C. (2001). Prevention of anxiety symptoms in primary school children: Preliminary results from a universal school-based trial. *British Journal of Clinical Psychology, 40*, 399–410. doi: 10.1348/014466501163887
- Baydar, N., Reid, M. J., & Webster-Stratton, C. (2003). The role of mental health factors and program engagement in the effectiveness of a preventive parenting program for Head Start mothers. *Child Development, 74*, 1433–1453. doi: 10.1111/1467-8624.00616
- Beardslee, W. R., Gladstone, T. R., Wright, E. J., & Cooper, A. B. (2003). A family-based approach to the prevention of depressive symptoms in children at risk: evidence of parental and child change. *Pediatrics, 112*, 119–131. doi: 10.1542/peds.112.2.e119

- Beauchaine, T. P., Webster-Stratton, C., & Reid, M. J. (2005). Mediators, moderators, and predictors of 1-year outcomes among children treated for early-onset conduct problems: A latent growth curve analysis. *Journal of Consulting and Clinical Psychology, 73*, 371. doi: 10.1037/0022-006X.73.3.371
- Beidel, D. C., & Turner, S. M. (2007). Clinical presentation of social anxiety disorder in children and adolescents. In D. C. Beidel & S. M. Turner (Eds.), *Shy children, phobic adults: Nature and treatment of social anxiety disorders* (2nd ed., pp. 47–80). Washington, DC: American Psychological Association.
- Bell, C. C., Bhana, A., Petersen, I., McKay, M. M., Gibbons, R., Bannon, W., & Amaty, A. (2008). Building protective factors to offset sexually risky behaviors among black youths: a randomized control trial. *Journal of the National Medical Association, 100*, 936–944.
- Benish, S. G., Quintana, W., & Wamold, B. (2011). Culturally adapted psychotherapy and the legitimacy of myth: A direct comparison meta-analysis. *Journal of Counseling Psychology, 58*, 279–289. doi: 10.1037/a0023626
- Berenson, G. S. (2002). Childhood risk factors predict adult risk associated with subclinical cardiovascular disease: The Bogalusa Heart Study. *American Journal of Cardiology, 90*, L3–L7. doi: 10.1016/S0002-9149(02)02953-3
- Berkel, C., Murry, V., Hurt, T., Chen, Y., Brody, G., Simons, R., . . . Gibbons, F. (2009). It takes a village: Protecting rural African American youth in the context of racism. *Journal of Youth and Adolescence, 38*(2), 175–188. doi: 10.1007/s10964-008-9346-z
- Bernal, G., Bonilla, J., & Bellido, C. (1995). Ecological validity and cultural sensitivity for outcome research: issues for the cultural adaptation and development of psychosocial treatments with Hispanics. *Journal of Abnormal Child Psychology, 23*, 67–82. doi: 10.1007/BF01447045
- Bernal, G., & Domenech Rodríguez, M. (Eds.). (2012). *Cultural adaptations: Tools for evidence based practice with diverse populations*. Washington, DC: American Psychological Association.
- Bernal, G., Jimenez-Chafey, M. I., & Domenech Rodríguez, M. M. (2009). Cultural adaptation of treatments: A resource for considering culture in evidence-based practice. *Professional Psychology: Research and Practice, 40*, 361–368. doi: 10.1037/a0016401
- Bernal, G., & Scharró-del-Río, M. R. (2001). Are empirically supported treatments valid for ethnic minorities? Toward an alternative approach for treatment research. *Cultural Diversity and Ethnic Minority Psychology, 7*, 328. doi: 10.1037/1099-9809.7.4.328
- Betancourt, H., & Lopez, S. R. (1993). The study of culture, ethnicity, and race in American psychology. *American Psychologist, 48*, 629–637. doi: 10.1037/0003-066X.48.6.629
- Bird, H. R. (1996). Epidemiology of childhood disorders in a cross-cultural context. *Journal of Child Psychology and Psychiatry, 37*, 35–49. doi: 10.1111/j.1469-7610.1996.tb01379.x
- Bjorknes, R., & Manger, T. (2013). Can parent training alter parent practice and reduce conduct problems in ethnic minority children? A randomized controlled trial. *Prevention Science, 14*(1), 52–63. doi: 10.1007/s11121-012-0299-9
- Borrego, J., Jr., Anhalt, K., Terao, S. Y., Vargas, E. C., & Urquiza, A. J. (2006). Parent-child interaction therapy with a Spanish-speaking family. *Cognitive and Behavioral Practice, 13*, 121–133. doi: 10.1016/j.cbpra.2005.09.001
- Borrego, J., Ibanez, E. S., Spendlove, S. J., & Pemberton, J. R. (2007). Treatment acceptability among Mexican American parents. *Behavior Therapy, 38*, 218–227. doi: 10.1016/j.beth.2006.08.007
- Botvin, G. J., Baker, E., Dusenbury, L., & Botvin, E. M. (1995). Long-term follow-up results of a randomized drug abuse prevention trial in a white middle-class population. *Journal of the American Medical Association, 273*, 1106–1112. doi: 10.1001/jama.273.14.1106
- Botvin, G. J., Baker, E., Dusenbury, L., Tortu, S., & Botvin, E. M. (1990). Preventing adolescent drug abuse through a multimodal cognitive-behavioral approach: Results of a 3-year study. *Journal of Consulting and Clinical Psychology, 58*, 437–446. doi: 10.1037/0022-006X.58.4.437
- Botvin, G. J., Griffin, K. W., Diaz, T., & Ifill-Williams, M. (2001). Drug abuse prevention among minority adolescents: Posttest and one-year follow-up of a school-based preventive intervention. *Prevention Science, 2*, 1–13. doi: 10.1023/A:1010025311161
- Branscum, P., & Sharma, M. (2012). After-school based obesity prevention interventions: A Comprehensive review of the literature. *International Journal of Environmental Research and Public Health, 9*, 1438–1457. doi: 10.3390/ijerph9041438
- Brody, G. H., Flor, D. L., & Neubaum, E. (1998). Coparenting processes and child competence among rural African American families. In M. Lewis & C. Feiring (Eds.), *Families, risk, and competence* (pp. 227–243). Mahwah, NJ: Erlbaum.
- Brody, G. H., Kim, S., Murry, V. M., & Brown, A. C. (2004a). Protective longitudinal paths linking child competence to behavioral problems among African American siblings. *Child Development, 75*, 455–467. doi: 10.1111/j.1467-8624.2004.00686.x
- Brody, G. H., Murry, V. M., Brown, A. C., & Anderson, T. (2004, May). *Translating research into prevention effects*. Paper presented at the Society for Prevention Research, Montreal, Canada.
- Brody, G. H., Murry, V. M., Kim, S., & Brown, A. C. (2002). Longitudinal pathways to competence and psychological adjustment among African American children living in rural single-parent households. *Child Development, 73*, 1505–1516. doi: 10.1111/1467-8624.00486
- Brody, G. H., Murry, V. M., McNair, L., Chen, Y.-F., Gibbons, F. X., Gerrard, M., & Willis, T. A. (2005). Linking changes in parenting to parent-child relationship quality and youth self-control: The Strong African American Families program. *Journal of Research on Adolescence, 15*, 47–69. doi: 10.1111/j.1532-7795.2005.00086.x
- Brody, G. H., Stoneman, Z., Flor, D., McCrary, C., Hastings, L., & Conyers, O. (1994). Financial resources, parent psychological functioning, parent co-caregiving, and early adolescent competence in rural two-parent African-American Families. *Child Development, 65*, 590–605. doi: 10.1111/j.1467-8624.1994.tb00770.x
- Brotman, L. M., Calzada, E., Huang, K. Y., Kingston, S., Dawson-McClure, S., Kamboukos, D., . . . Petkova, E. (2011). Promoting effective parenting practices and preventing child behavior problems in school among ethnically diverse families from underserved, urban communities. *Child Development, 82*, 258–276. doi: 10.1111/j.1467-8624.2010.01554.x
- Brotman, L. M., Gouley, K. K., Huang, K. Y., Rosenfelt, A., O'Neal, C., Klein, R. G., & Shrout, P. (2008). Preventive intervention for preschoolers at high risk for antisocial behavior: Long-term effects on child physical aggression and parenting practices. *Journal of Clinical Child and Adolescent Psychology, 37*, 386–396. doi: 10.1080/15374410801955813
- Brotman, L. M., Gouley, K. K., Klein, R. G., Castellanos, F. X., & Pine, D. S. (2003). Children, stress, and context: Integrating basic, clinical, and experimental prevention research. *Child Development, 74*, 1053–1057. doi: doi.org/10.1111/1467-8624.00589
- Brown, C. H., Berndt, D., Brinales, J. M., Zong, X., & Bhagwat, D. (2000). Evaluating the evidence of effectiveness for preventive interventions: Using a registry system to influence policy through science. *Addictive Behavior, 25*, 955–964. doi: 10.1016/S0306-4603(00)00131-3

- Brownson, F. C., Colditz, F. A., & Proctor, E. K. (2012). *Dissemination and implementation research in health*. New York, NY: Oxford University Press.
- Caballero, B., Clay, T., Davis, S. M., Ethelbah, B., Rock, B. H., Lohman, T., ... Stevens, J. (2003). Pathways: A school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. *American Journal of Clinical Nutrition*, *78*, 1030–1038.
- Cabassa, L. J., & Baumann, A. A. (2013). A two-way street: Bridging implementation science and cultural adaptations of mental health treatments. *Implementation Science*, *8*, 90. doi: 10.1186/1748-5908-8-90
- Cardemil, E. V. (2010). Cultural adaptations to empirically supported treatments: A research agenda. *Scientific Review of Mental Health Practice*, *7*, 8–21.
- Cardemil, E. V., Reivich, K. J., Beevers, C. G., Seligman, M. E. P., & James, J. (2007). The prevention of depressive symptoms in low-income, minority children: Two-year follow-up. *Behaviour Research and Therapy*, *45*, 313–327. doi: 10.1016/j.brat.2006.03.010
- Cardemil, E. V., Reivich, K. J., & Seligman, M. E. P. (2002). The prevention of depressive symptoms in low-income minority middle school students. *Prevention & Treatment*, *5*, Article 8. doi: 10.1037/1522-3736.5.1.58a
- Castro, G. F., Barrera, M., Jr., & Holleran Steiker, L. K. (2010). Issues and challenges in the design of culturally adapted evidence-based interventions. *Annual Review of Clinical Psychology*, *6*, 213–239. doi: 10.1146/annurev-clinpsy-033109-132032
- Castro, F. G., Barrera, M., Jr., & Martinez, C. R. (2004). The cultural adaptation of preventive interventions: Resolving tensions between fidelity and fit. *Prevention Science*, *5*, 41–45. doi: 10.1023/B:PREV.0000013980.12412.cd
- Centers for Disease Control and Prevention. (2011). Diagnoses of HIV Infection in the United States. Retrieved from <http://www.cdc.gov/MMWR/PDF/SS/SS6104/PDF>.
- Chaffin, M., Valle, L. A., Funderburk, B., Gurwitsch, R., Silovsky, J., Bard, D., & Kees, M. (2009). A motivational intervention can improve retention in PCIT for low-motivation child welfare clients. *Child Maltreatment*, *14*, 356–368. doi: 10.1177/1077559509332263
- Chambless, D. L., & Hollon, S. D. 1998. Defining empirically supported therapies. *Journal of Consulting and Clinical Psychology*, *66*, 7–18. doi: 10.1037/0022-006X.66.1.7
- Cicchetti, D. (1990). A historical perspective on the discipline of developmental psychopathology. In J. Rolf, A. Masten, D. Cicchetti, K. Nuechterlein, & S. Weintraub (Eds.), *Risk and protective factors in the development of psychopathology* (pp. 2–28). New York, NY: Cambridge University Press.
- Cicchetti, D., & Hinshaw, S. P. (2002). Editorial: Prevention and intervention science: Contributions to developmental theory [Special Issue]. *Development and Psychopathology*, *14*, 667–981. doi: 10.1017/S0954579402004017
- Cicchetti, D., & Toth, S. L. (1992). The role of developmental theory in prevention and intervention. *Development and Psychopathology*, *4*, 489–493. doi: 10.1017/S0954579400004831
- Clark, L. F., Miller, K. S., Nagy, S. S., Avery, J., Roth, D. L., Liddon, N., & Mukherjee, S. (2005). Adult identity mentoring: Reducing sexual risk for African-American seventh grade students. *Journal of Adolescent Health*, *37*, 337. doi: 10.1016/j.jadohealth.2004.09.024
- Coard, S. I., Foy-Watson, S., Zimmer, C., & Wallace, A. (2007). Considering culturally relevant parenting practices in intervention development and adaptation: A randomized controlled trial of the Black Parenting Strengths and Strategies (BPSS) program. *Counseling Psychologist*, *35*, 797–820. doi: 10.1177/0011000007304592
- Coard, S. I., Wallace, S. A., Stevenson, H. C., Jr., & Brotman, L. M. (2004). Towards culturally relevant preventive interventions: The consideration of racial socialization in parent training with African American families. *Journal of Child and Family Studies*, *13*, 277–293. doi: 10.1023/B:JCFS.0000022035.07171.f8
- Coatsworth, J. D., Pantin, H., & Szapocznik, J. (2002). Families Unidas: A family-centered ecodevelopmental intervention to reduce risk for problem behavior among Hispanic adolescents. *Clinical Child and Family Psychology Review*, *5*, 113–132. doi: 10.1023/A:1015420503275
- Collins, L. M., Murphy, S. A., & Bierman, K. L. (2004). A conceptual framework for adaptive preventive interventions. *Prevention Science*, *5*, 185–196. doi: 10.1023/B:PREV.0000037641.26017.00
- Comas Díaz, L. (2006). Latino healing: The integration of ethnic psychology into psychotherapy. *Psychotherapy: Theory, Research, Practice, Training*, *43*, 436–453. doi: 10.1037/0033-3204.43.4.436
- Conger, R. D., & Rueter, M. A. (1996). Siblings, parents, and peers: A longitudinal study of social influences in adolescent risk for alcohol use and abuse. In G. H. Brody (Ed.), *Sibling relationships: Their causes and consequences* (pp. 1–30). Westport, CT: Ablex.
- Cooley, M. R., Boyd, R. C., & Grados, J. J. (2004). Feasibility of an anxiety preventive intervention for community violence exposed African-American children. *Journal of Primary Prevention*, *25*, 105–123. doi: 10.1023/B:JOPP.0000039941.85452.ea
- Cooley-Strickland, M. R., Griffin, R. S., Darney, D., Otte, K., & Ko, J. (2011). Urban African American youth exposed to community violence: A school-based anxiety preventive intervention efficacy study. *Journal of Prevention & Intervention in the Community*, *39*, 149–166. doi: 10.1080/10852352.2011.556573
- Costello, E. J., Angold, A., & Keeler, G. P. (1999). Adolescent outcomes of childhood disorders: The consequences of severity and impairment. *Journal of the American Academy of Child & Adolescent Psychiatry*, *38*, 121–128.
- Costello, E. J., Egger, H., & Angold, A. (2005). 10-year research update review: The epidemiology of child and adolescent psychiatric disorders: I. methods and public health burden. *Journal of the American Academy of Child & Adolescent Psychiatry*, *44*, 972–986.
- Cowie, C. C., Rust, K. F., Byrd-Holt, D. D., Eberhardt, M. S., Flegal, K. M., Engelgau, M. M., ... Gregg, E. W. (2006). Prevalence of diabetes and impaired fasting glucose in adults in the US population: National Health and Nutrition Examination Survey 1999–2002. *Diabetes Care*, *29*, 1263–1268. doi: 10.2337/dc06-0062
- Crespo, N. C., Elder, J. P., Ayala, G. X., Slymen, D. J., Campbell, N. R., Sallis, J. F., ... Arredondo, E. M. (2012). Results of a multi-level intervention to prevent and control childhood obesity among Latino children: The Aventuras Para Niños Study. *Annals of Behavioral Medicine*, *43*, 84–100. doi: 10.1007/s12160-011-9332-7
- Crisante, L., & Ng, S. (2003). Implementation and process issues in using group Triple P with Chinese parents: Preliminary findings. *Advances in Mental Health*, *2*, 226–235. doi: 10.5172/jamh.2.3.226
- Cunningham, C. E., Boyle, M., Offord, D., Racine, Y., Hundert, J., Secord, M., & McDonald, J. (2000). Tri-ministry study: Correlates of school-based parenting course utilization. *Journal of Consulting and Clinical Psychology*, *68*, 928–933. doi: 10.1037/0022-006X.68.5.928
- Dadds, M. R., Spence, S. H., Holland, D. E., Barrett, P. M., & Laurens, K. R. (1997). Prevention and early intervention for anxiety disorders: A controlled trial. *Journal of Consulting and Clinical Psychology*, *65*, 627–635. doi: 10.1037/0022-006X.65.4.627
- D'Angelo, E. J., Llerena-Quinn, R., Shapiro, R., Colon, F., Rodriguez, P., Gallagher, K., & Beardslee, W. R. (2009). Adaptation of the preventive intervention program for depression for use with predominantly low-income Latino families. *Family Process*, *48*, 269–291. doi: 10.1111/j.1545-5300.2009.01281.x

- Davis, S. M., Clay, T., Smyth, M., Gittelsohn, J., Arviso, V., Flint-Wagner, H., . . . Stone, E. J. (2003). Pathways curriculum and family interventions to promote healthful eating and physical activity in American Indian schoolchildren. *Preventive Medicine, 37*, S24–S34. doi: 10.1016/j.ypmed.2003.08.011
- Davis, S. M., Going, S. B., Helitzer, D. L., Teufel, N. I., Snyder, P., Gittelsohn, J., . . . Altaba, J. (1999). Pathways: A culturally appropriate obesity-prevention program for American Indian schoolchildren. *American Journal of Clinical Nutrition, 69*(Suppl.), 796S–802S.
- Davis, J. N., Ventura, E. E., Tung, A., Munevar, M. A., Hasson, R. E., Byrd-Williams, C., . . . Goran, M. I. (2012). Effects of a randomized maintenance intervention on adiposity and metabolic risk factors in overweight minority adolescents. *Pediatric Obesity, 7*, 16–27. doi: 10.1111/j.2047-6310.2011.00002.x
- Deater-Deckard, K., & Dodge, K. A. (1997). Externalizing behavior problems and discipline revisited: Nonlinear effects and variation by culture, context, and gender. *Psychological Inquiry, 8*, 161–175. doi: 10.1207/s15327965pli0803_1
- Deci, E. L. (1992). The relation of interest to the motivation of behavior: a self-determination theory perspective. In K. A. Renninger, S. Hidi, & A. Krapp (Eds.), *The role of interest in learning and development* (pp. 43–70). Hillsdale, NJ: Erlbaum.
- Deci, E. L., Driver, R. E., Hotchkiss, L., Robbins, R. J., & Wilson, I. M. (1993). The relation of mothers' controlling vocalizations to children's intrinsic motivation. *Journal of Experimental Child Psychology, 55*, 151–162. doi: 10.1006/jecp.1993.1008
- de Graaf, I., Speetjens, P., Smit, F., de Wolff, M., & Tavecchio, L. (2008). Effectiveness of the Triple P Positive Parenting Program on behavioral problems in children: A meta-analysis. *Behavior Modification, 32*, 714–735. doi: 10.1177/0145445508317134
- de Heer, H. D., Koehly, L., Pederson, R., & Morera, O. (2011). Effectiveness and spillover of an after-school health promotion program for Hispanic elementary school children. *American Journal of Public Health, 101*, 1907–1913. doi: 10.2105/AJPH.2011.300177
- Delva, J., O'Malley, P. M., & Johnston, L. D. (2006). Racial/ethnic and socioeconomic status differences in overweight and health-related behaviors among American students: National trends 1986–2003. *Journal of Adolescent Health, 39*, 536–545. doi: 10.1016/j.jadohealth.2006.02.013
- DiClemente, R. J., Wingood, G. M., Rose, E. S., Sales, J. M., Lang, D. L., Caliendo, A. M., . . . Crosby, R. A. (2009). Efficacy of sexually transmitted disease/human immunodeficiency virus sexual risk-reduction intervention for African American adolescent females seeking sexual health services. *Archives of Pediatric & Adolescent Medicine, 163*, 1112–1121. doi: 10.1001/archpediatrics.2009.205
- DiIorio, C., McCarty, F., Resnicow, K., Lehr, S., & Denzmore, P. (2007). REAL men: A group-randomized trial of an HIV prevention intervention for adolescent boys. *American Journal of Public Health, 97*, 1084–1089. doi: 10.2105/AJPH.2005.073411
- DiIorio, C., Resnicow, K., McCarty, F., De, A. K., Dudley, W. N., Wang, D. T., & Denzmore, P. (2006). Keepin' it R.E.A.L.: Results of a mother-adolescent HIV prevention program. *Nursing Research, 55*, 43–51. doi: 10.1097/00006199-200601000-00006
- Dillman Carpentier, F., Mauricio, A. M., Gonzales, N. A., Millsap, R., Meza, C., Dumka, L. E., . . . & Genalo, M. T. (2007). Engaging Mexican origin families in a school-based preventive intervention. *Journal of Primary Prevention, 28*, 521–546. doi: 10.1007/s10935-007-0110-z
- Dionne, R., Davis, B., Sheeber, L., & Madrigal, L. (2009). Initial evaluation of a cultural approach to implementation of evidence-based parenting interventions in American Indian communities. *Journal of Community Psychology, 37*, 911–921. doi: 10.1002/jcop.20336
- Dishion, R. J., Kavanagh, K., Schneiger, A., Nelson, A., & Kaufman, N. K. (2002). Preventing early adolescent substance use: A family-centered strategy for public middle schools. *Prevention Science, 3*, 191–201. doi: 10.1023/A:1019994500301
- Domenech Rodríguez, M. M., & Bernal, G. (2012). Frameworks, models and guides. In G. Bernal & M. M. Domenech-Rodríguez (Eds.), *Cultural adaptations: Tools for evidence-based practice with diverse populations*. pp. 23–44 Washington, DC: American Psychological Association.
- Domenech Rodríguez, M. M., Baumann, A. A., & Schwartz, A. L. (2011). Cultural adaptation of an evidence based intervention: From theory to practice in a Latino/a community context. *American Journal of Community Psychology, 47*, 170–186. doi: 10.1007/s10464-010-9371-4
- Dumka, L. E., Gonzales, N. A., Woods, J., & Formoso, D. (1998). Using qualitative methods to develop contextually relevant measures and preventive interventions: An illustration. *American Journal of Community Psychology, 26*(4), 605–637. doi: 10.1023/A:1022145022830
- Dryfoos, J. G. (1993). Preventing substance use: Rethinking strategies. *American Journal of Public Health, 83*, 6. doi: 10.2105/AJPH.83.1.6
- Eaton, D. K., Kann, L., Kinchen, S., Shanklin, S., Flint, K. H., Hawkins, J., . . . Wechsler, H. (2012). Youth risk behavior surveillance—United States, 2011. *Surveillance Summaries, Morbidity and Mortality Weekly Report, 61*(4).
- Eaton, D. K., Kann, L., Kinchen, S., Shanklin, S., Ross, J., Hawkins, J., . . . Wechsler, H. (2008). Youth risk behavior surveillance—United States, 2007. *Surveillance Summaries, Morbidity and Mortality Weekly Report, 57*(SS04).
- Eimers, R., & Aitchison, R. (1977). *Effective parents—responsible children: A guide to confident parenting*. New York, NY: McGraw-Hill.
- Eisenstadt, T., Eyberg, S., McNeil, C., Newcomb, K., & Funderburk, B. (1993). Parent-child interaction therapy with behavior problem children: Relative effectiveness of two stages and overall treatment outcome. *Journal of Clinical Child Psychology, 22*, 42–51. doi: 10.1207/s15374424jccp2201_4
- Elliot, D. S., & Mihalic, S. (2004). Issues in disseminating and replicating effective prevention programs. *Prevention Science, 5*, 47–53. doi: 10.1023/B:PREV.0000013981.28071.52
- Essau, C. A., Conradt, J., Sasagawa, S., & Ollendick, T. H. (2012). Prevention of anxiety symptoms in children: Results from a universal school-based trial. *Behavior Therapy, 43*(2), 450–446. doi: 10.1016/j.beth.2011.08.003
- Eyberg, S. M., Nelson, M. M., & Boggs, S. R. (2008). Evidence-based psychosocial treatments for children and adolescents with disruptive behavior. *Journal of Clinical Child and Adolescent Psychology, 37*, 215–237. doi: 10.1080/15374410701820117
- Eyberg, S. M., & Pincus, D. (1999). *Eyberg Child Behavior Inventory and Sutter-Eyberg Student Behavior Inventory-Revised: Professional manual*. Psychological Assessment Resources.
- Eyre, H., Kahn, R., & Robertson, R. M. (2004). Preventing cancer, cardiovascular disease and diabetes: A common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. *Diabetes Care, 27*, 1812–1824. doi: 10.2337/diacare.27.7.1812
- Fagot-Campagna, A., Pettitt, D. J., Engelgau, M. M., Burrows, N. R., Geiss, L. S., Valdez, R., . . . Narayan, K. M. (2000). Type 2 diabetes among North American children and adolescents: An epidemiologic review and a public health perspective. *Journal of Pediatrics, 136*, 664–672. doi: 10.1067/mpd.2000.105141
- Falicov, C. J. (2009). Commentary: On the wisdom and challenges of culturally attuned treatments for Latinos. *Family Process, 48*, 292–309. doi: 10.1111/j.1545-5300.2009.01282.x

- Feinberg, E., Stein, R., Diaz-Linhart, Y., Egbert, L., Beardless, W., Hegel, M. T., & Silverstein, M. (2012). Adaptation of problem-solving treatment for depression among low-income, culturally diverse mothers. *Family Community Health, 35*, 57–67. doi: 10.1097/FCH.0b013e3182385d48
- Fernandez, M. A., Butler, A. M., & Eyberg, S. M. (2011). Treatment outcome for low socioeconomic status african american families in parent-child interaction therapy: A pilot study. *Child & Family Behavior Therapy, 33*, 32–48. doi: 10.1080/07317107.2011.545011
- Fernandez, M. A., & Eyberg, S. M. (2009). Predicting treatment and follow-up attrition in parent-child interaction therapy. *Journal of Abnormal Child Psychology, 37*, 431–441. doi: 10.1007/s10802-008-9281-1
- Ferrer-Wreder, L., Sundell, K., & Mansoor, S. (2012). Tinkering with perfection: Theory development in the intervention cultural adaptation field. *Child Youth Care Forum, 41*, 149–171. doi: 10.1007/s10566-011-9162-6
- Fisak, B., Jr., Richard, D., & Mann, A. (2011). The prevention of child and adolescent anxiety: A meta-analytic review. *Prevention Science, 12*, 255–268. doi: 10.1007/s11121-011-0210-0
- Fitzgibbon, M. L., Prewitt, T. E., Blackman, L. R., Simon, P., Luke, A., Keys, L. C., . . . Singh, V. (1998). Quantitative assessment of recruitment efforts for prevention trials in two diverse black populations. *Preventive Medicine, 27*, 838–45. doi: 10.1006/pmed.1998.0367
- Fitzgibbon, M. L., Stolley, M. R., Schiffer, S., Van Horn, L., Kaufer-Christoffel, K., & Dyer, A. (2002). A community-based obesity prevention program for minority children: Rationale and study design for Hip-Hop to Health Jr. *Preventive Medicine, 34*, 289–297. doi: 10.1006/pmed.2001.0977
- Fitzgibbon, M. L., Stolley, M. R., Schiffer, S., Van Horn, L., Kaufer-Christoffel, K., & Dyer, A. (2005). Two-year follow-up results for Hip-Hop to Health Jr.: A randomized controlled trial for overweight prevention in preschool minority children. *Journal of Pediatrics, 146*, 618–625. doi: 10.1016/j.jpeds.2004.12.019
- Fitzgibbon, M. L., Stolley, M. R., Schiffer, S., Van Horn, L., Kaufer-Christoffel, K., & Dyer, A. (2006). Hip-Hop to Health Jr. for Latino preschool children. *Obesity, 14*, 1616–1625. doi: 10.1038/oby.2006.186
- Flay, B. R., Biglan, A., Boruch, R. F., Castro, F., Gottfredson, D., Kellam, S., . . . Ji, P. (2005). Standards of evidence: Criteria for efficacy, effectiveness and dissemination. *Prevention Science, 12*, 103–117. doi: 10.1007/s11121-005-5553-y
- Flay, B. R., Graumlich, S., Segawa, E., Burns, J. L., & Holliday, M. Y. (2004). Effects of 2 prevention programs on high-risk behaviors among African American youth: A randomized trial. *Archives of Pediatrics & Adolescent Medicine, 158*, 377–384. doi: 10.1001/archpedi.158.4.377
- Flores, G. (2010). Racial and ethnic disparities in the health and health care of children. *Pediatrics, 125*, e979–e1020. doi: 10.1542/peds.2010-0188
- Flores, G., & Tomany-Korman, S. C. (2008). Racial and ethnic disparities in medical and dental health, access to care, and use of services in US children. *Pediatrics, 121*, e286–e298. doi: 10.1542/peds.2007-1243
- Forehand, R., & Kotchick, B. A. (1996). Cultural diversity: A wake-up call for parent training. *Behavior Therapy, 27*, 187–206. doi: 10.1016/S0005-7894(96)80014-1
- Forehand, R., & Long, N. (2002). *Parenting the strong-willed child: The clinically-proven five-week program for parents of two- to six-year-olds* (2nd ed.). New York, NY: McGraw-Hill.
- Forgatch, M. S. (1994). *Parenting through change: A programmed intervention curriculum for groups of single mothers*. Eugene, OR: Oregon Social Learning Center.
- Forgatch, M. S., & Martinez, C. R. (1999). Parent management training: A program linking basic research and practical application. *Tidsskrift for Norsk Psykologforening, 36*, 923–937.
- Forgatch, M. S., & Patterson, G. R. (2010). Parent Management Training—Oregon Model: An intervention for antisocial behavior in children and adolescents. In J. R. Weisz & A. E. Kazdin (Eds.), *Evidence-based psychotherapies for children and adolescents* (Vol. 2, pp. 159–178). New York, NY: Guilford Press.
- Foxcroft, D. R., Ireland, D., Lister-Sharp, D. J., Lowe, G., & Breen, R. (2003). Longer-term primary prevention for alcohol misuse in young people: A systematic review. *Addiction, 98*, 397–411. doi: 10.1046/j.1360-0443.2003.00355.x
- Gahagan, S., & Silverstein, J. (2003). Prevention and treatment of type 2 diabetes mellitus in children, with special emphasis on American Indian and Alaska Native children. *Pediatrics, 112*, e328–e328. doi: 10.1542/peds.112.4.e328
- García Coll, C., Lambert, G., Jenkins, R., McAdoo, H. P., Crnic, K., Wasik, B. H., & Vásquez, H. (1996). An integrative model for the study of developmental competencies in minority children. *Child Development, 67*, 1891–1914. doi: 10.2307/1131600
- Garland, A. F., Hawley, K. M., Brookman-Frazee, L., & Hurlburt, M. S. (2008). Identifying common elements of evidence-based psychosocial treatments for children's disruptive behavior problems. *Journal of the American Academy of Child and Adolescent Psychiatry, 47*, 505–514. doi: 10.1097/CHI.0b013e31816765c2
- Germán, L., Gonzales, N. A., West, S. G., & Dumka, L. E. (in review). An experimental test of parental influences on risky sexual behaviors among Mexican-origin adolescents. Manuscript submitted for publication.
- Gibbons, F. X., Gerrard, M., Cleveland, M. J., Wills, T. A., & Brody, G. (2004). Perceived discrimination and substance use in African American parents and their children: A panel study. *Journal of Personality and Social Psychology, 86*, 517–529. doi: 10.1037/0022-3514.86.4.517
- Gillman, J. E., Reivich, K. J., Jaycox, L. H., & Seligman, M. E. P. (1995). Prevention of depressive symptoms in schoolchildren: Two-year follow-up. *Psychological Science, 6*, 343–351. doi: 10.1111/j.1467-9280.1995.tb00524.x
- Ginsburg, G. S. (2009). The child anxiety prevention study: Intervention model and primary outcomes. *Journal of Consulting and Clinical Psychology, 77*, 580–587. doi: 10.1037/a0014486
- Gittelsohn, J., Evans, M., Story, M., Davis, S. M., Metcalfe, L., Helitzer, & Clay, T. E. (1999). Multisite formative assessment for the Pathways study to prevent obesity in American Indian schoolchildren. *American Journal of Clinical Nutrition, 69*(suppl), 767S–772S.
- Glantz, M. D., & Leshner, A. I. (2000). Drug abuse and developmental psychopathology. *Development and Psychopathology, 12*, 795–814. doi: 10.1017/S0954579400004120
- Gone, J. P. (2004). Mental health services for Native Americans in the 21st century United States. *Professional Psychology: Research and Practice, 35*, 10–18. doi: 10.1037/0735-7028.35.1.10
- Gonzales, N. A., Dumka, L., Deardorff, J., Jacobs-Carter, S., & McCray, A. (2004). Preventing poor mental health and school dropout of Mexican American adolescents following the transition to junior high school. *Journal of Adolescent Research, 19*, 113–131. doi: 10.1177/0743558403258124
- Gonzales, N. A., Dumka, L. E., Mauricio, A. M., & Germán, M. (2007). Building bridges: Strategies to promote academic and psychological resilience for adolescents of Mexican origin. In J. E. Lansford, K. Deater-Deckard, & M. H. Bornstein (Eds.), *Immigrant families in contemporary society* (pp. 268–286). New York, NY: Guilford Press.
- Gonzales, N. A., Dumka, L. E., Millsap, R. E., Gottschall, A., McClain, D. B., Wong, J. J., . . . Kim, S. Y. (2012). Randomized trial of a broad

- preventive intervention for Mexican American adolescents. *Journal of Consulting and Clinical Psychology*, 80, 1–16. doi: 10.1037/a0026063
- Gonzales, N. A., Jensen, M., Wong, J. J., Dumka, L. E., & Millsap, R. (in review). Long term effects of family based prevention on Mexican American adolescents substance use and abuse in emerging adulthood. Manuscript submitted for publication.
- Gonzales, N. A., Wong, J. J., Toomey, R. B., Millsap, R., Dumka, L. E., Mauricio, A. M. (2014). School engagement mediates long term prevention effects for Mexican American adolescents. *Prevention Science*, 15, 929–939.
- Goodkind, J. R., LaNoue, M. D., & Milford, J. (2010). Adaptation and implementation of cognitive behavioral intervention for trauma in schools with American Indian youth. *Journal of Clinical Child and Adolescent Psychology*, 39, 858–872. doi: 10.1080/15374416.2010.517166
- Gordon-Larsen, P., Adair, L. S., & Popkin, B. M. (2003). The relationship of ethnicity, socioeconomic factors, and overweight in U.S. Adolescents. *Obesity Research*, 11, 121–129. doi: 10.1038/oby.2003.20
- Gottfredson, D., Kumpfer, K., Polizzi-Fox, D., Wilson, D., Puryear, V., Beatty, P., & Bilmenay, M. (2006). The Strengthening Washington DC Families project: A randomized effectiveness trial of family-based prevention. *Prevention Science*, 7, 57–74. doi: 10.1007/s11121-005-0017-y
- Grant, B. F., Stinson, F. S., Hasin, D. S., Dawson, D. A., Chou, S., & Anderson, K. (2004). Immigration and lifetime prevalence of *DSM-IV* psychiatric disorders among Mexican Americans and non-Hispanic Whites in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Archives of General Psychiatry*, 61, 1226–1233. doi: 10.1001/archpsyc.61.12.1226
- Griner, D., & Smith, T. B. (2006). Culturally adapted mental health interventions: a meta-analytic review. *Psychotherapy: Theory, Research, Practice, Training*, 43, 531–548. doi.org/10.1037/0033-3204.43.4.531
- Gross, D., Fogg, L., Webster-Stratton, C., Garvey, C., Julion, W., & Grady, J. (2003). Parent training of toddlers in day care in low-income urban communities. *Journal of Consulting and Clinical Psychology*, 71, 261. doi: 10.1037/0022-006X.71.2.261
- Gross, D., Garvey, C., Julion, W. A., & Fogg, L. (2007). Preventive parent training with low-income, ethnic minority families of preschoolers. In J. M. Briesmeister & C. E. Schaefer (Eds.), *Handbook of parent training: Helping parents prevent and solve problem behaviors* (3rd ed., pp. 5–24). Hoboken, NJ: Wiley.
- Gross, D., Garvey, C., Julion, W., Fogg, L., Tucker, S., & Mokros, H. (2009). Efficacy of the Chicago Parent Program with low-income African American and Latino parents of young children. *Prevention Science*, 10, 54–65. doi: 10.1007/s11121-008-0116-7
- Grunbaum, J. A., Lowry, R., Kann, L., & Pateman, B. (2000) Prevalence of health risk behaviors among Asian American/Pacific Islander high school students. *Journal of Adolescent Health*, 27, 322–30. doi: 10.1016/S1054-139X(00)00093-8
- Guo, S. S., & Chumlea, W. C. (1999). Tracking of body mass index in children in relation to overweight in adulthood. *American Journal of Clinical Nutrition*, 70(Suppl.), 145S–148S.
- Halgunseth, L. C., Ispa, J. M., & Rudy, D. (2006). Parental control in Latino families: An integrated review of the literature. *Child Development*, 77, 1282–1297. doi: 10.1111/j.1467-8624.2006.00934.x
- Hanf, C. A. (1969). *A two-stage program for modifying maternal controlling during mother-child (M-C) interaction*. Paper presented at the meeting of the Western Psychological Association, Vancouver, Ontario.
- Harkness, S., & Super, C. M. (1996). *Parents' cultural belief systems: Their origins, expressions, and consequences*. New York, NY: Guilford Press.
- Hartmann, W. E., Kim, E. S., Kim, J. H. J., Nguyen, T. U., Wendt, D. C., Nagata, D. K., & Gone, J. P. (2013). In search of cultural diversity, revisited: Recent publication trends in cross-cultural and ethnic minority psychology. *Review of General Psychology*, 17, 243–254. doi: 10.1037/a0032260
- Hawkins, J. D., Catalano, R. F., & Miller, J. Y. (1992). Risk and protective factors for alcohol and drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychological Bulletin*, 112, 64–105. doi: 10.1037/0033-2909.112.1.64
- Hecht, M. L., Graham, J. W., & Elek, E. (2006). The drug resistance strategies intervention: Program effects on substance use. *Health Communication*, 20, 267–276. doi: 10.1207/s15327027hc2003_6
- Hecht, M. L., Marsiglia, F. B., Elek, E., Wagstaff, D. A., Kulis, S., Dustman, P., & Miller-Day, M. (2003). Culturally grounded substance use prevention: An evaluation of the Keepin' It R.E.A.L. curriculum. *Prevention Science*, 4, 233–248. doi: 10.1023/A:1026016131401
- Henggeler, S. W., & Schaeffer, C. M. (2010). Treating serious emotional and behavioural problems using multisystemic therapy. *Australian and New Zealand Journal of Family Therapy*, 31, 149–164. doi: 10.1375/anft.31.2.149
- Henggeler, S. W., Schoenwald, S. K., Borduin, C. M., Rowland, M. D., & Cunningham, P. B. (1998). *Multisystemic treatment of antisocial behavior in children and adolescents*. New York, NY: Guilford Press.
- Herschell, A. D., Calzada, E. J., Eyberg, S. M., & McNeil, C. B. (2002). Research on Parent Child Interaction Therapy: Past and future. *Cognitive and Behavioral Practice*, 9, 9–16. doi: 10.1016/S1077-7229(02)80034-7
- Ho, C., Bluestein, D. N., & Jenkins, J. M. (2008). Cultural differences in the relationship between parenting and children's behavior. *Developmental Psychology*, 44, 507–522. doi: 10.1037/0012-1649.44.2.507
- Ho, T., Chow, V., Fung, C., Leung, K., Chiu, K., Yu, G., . . . Lieh-Mak, F. (1999). Parent management training in a Chinese population: Application and outcome. *Journal of the American Academy of Child & Adolescent Psychiatry*, 38, 1165–1172. doi: 10.1097/00004583-199909000-00022
- Ho, J., Yeh, M., McCabe, K., & Lau, A. (2012). Perceptions of the acceptability of parent training among Chinese immigrant parents: Contributions of cultural factors and clinical need. *Behavior Therapy*, 43, 436–449. doi: 10.1016/j.beth.2011.10.004
- Horowitz, J. L., & Garber, J. (2006). The prevention of depressive symptoms in children and adolescents: A meta-analytic review. *Journal of Consulting and Clinical Psychology*, 74, 401–415. doi: 10.1037/0022-006X.74.3.401
- Huey, S., Jr., & Pan, D. (2006). Culture-responsive one-session treatment for phobic Asian Americans: A pilot study. *Psychotherapy: Theory, Research, Practice, Training*, 43, 549–554. doi: 10.1037/0033-3204.43.4.549
- Huey, S. J., & Polo, A. J. (2008). Evidence-based psychosocial treatments for ethnic minority youth. *Journal of Clinical Child & Adolescent Psychology*, 37, 262–301. doi: 10.1080/15374410701820174
- Hutchison, L., & Blakely, C. (2003). Substance abuse-trends in rural areas: A literature review. In L. Gamm, L. Hutchison, B. D. Dabney, & A. M. Dorsey (Eds.), *Rural healthy people 2010: A companion document to Healthy People 2010* (Vol. 2, pp. 145–154). College Station, TX: Texas A & M University System Health Science Center, School of Rural Public Health, Southwest Rural Health Research Center.
- Jemmott, J. B., III. (1992). Reductions in HIV risk-associated sexual behaviors among black male adolescents: Effects of an AIDS prevention intervention. *American Journal of Public Health*, 82, 372–377. doi: 10.2105/AJPH.82.3.372
- Jemmott, J. B., III, Jemmott, L. S., & Fong, G. T. (1998). Abstinence and safer sex HIV risk-reduction interventions for African American

- adolescents: A randomized controlled trial. *Journal of the American Medical Association*, 279, 1529–1536. doi: 10.1001/jama.279.19.1529
- Jemmott, J. B., Jemmott, L. S., Fong, G. T., & McCaffree, K. (1999). Reducing HIV risk-associated sexual behavior among African American adolescents: Testing the generality of intervention effects. *American Journal of Community Psychology*, 27, 161–187. doi: 10.1007/BF02503158
- Jemmott, J. B., III, Jemmott, L. S., & Fong, G. T. (2010). Efficacy of a theory-based abstinence-only intervention over 24 months: A randomized controlled trial with young adolescents. *Archives of Pediatrics & Adolescent Medicine*, 164, 152–159. doi: 10.1001/archpediatrics.2009.267
- Jemmott J. B., III, Jemmott, L. S., Fong, G. T., & Morales, K. H. (2010). Effectiveness of an HIV/STD risk-reduction intervention for adolescents when implemented by community-based organizations: A cluster-randomized controlled trial. *American Journal of Public Health*, 100, 720–726. doi: 10.2105/AJPH.2008.140657
- Jensen, M., Wong, J. J., Gonzales, N. A., Millsap, R., Dumka, L. E. (2014). Long term effects of a universal family intervention: Mediation through parent adolescent conflict. *Journal of Clinical Child and Adolescent Psychology*, 43(3), 415–427.
- Johnston, L. D., Bachman, J. G., O'Malley, P. M., & Schulenberg, J. (2008). *Monitoring the Future: A Continuing Study of American Youth (8th- and 10th-Grade Surveys)*, 2000. Ann Arbor, MI: Inter-University Consortium for Political and Social Research.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2009). *Monitoring the future national results on adolescent drug use: Overview of key findings, 2008 (NIH Publication No. 09-7401)*. Bethesda, MD: National Institute on Drug Abuse.
- Kameoka, V. A. (1996). *The effects of a family-focused intervention on reducing risk for substance abuse among Asian and Pacific Island youths and families: Evaluation of the Strengthening Hawaii's Families Project*. Honolulu, HI: University of Hawaii, Social Welfare Evaluation and Research Unit.
- Kanekar, A., & Sharma, M. (2009). Meta-analysis of school-based childhood obesity interventions in the UK and US. *International Quarterly of Community Health Education*, 29, 241–256.
- Karno, M., Golding, J. M., Burnam, M. A., Hough, R. L., Escobar, J. T., Wells, K. M., & Boyer, R. (1989). Anxiety disorders among Mexican Americans and non-Hispanic Whites in Los Angeles. *Journal of Nervous and Mental Disease*, 177, 202–209. doi: 10.1097/00005053-198904000-00003
- Kazdin A. E. (2008). Evidence-based treatment and practice: new opportunities to bridge clinical research and practice, enhance the knowledge base, and improve patient care. *American Psychologist*, 63, 146–59. doi: 10.1037/0003-066X.63.3.146
- Kazdin, A. E., & Whitley, M. K. (2003). Treatment of parental stress to enhance therapeutic change among children referred for aggressive and antisocial behavior. *Journal of Consulting and Clinical Psychology*, 71, 504–515. doi: 10.1037/0022-006X.71.3.504
- Kessler, R. C., Avenevoli, S., Costello, J., Green, J. G., Gruber, M. J., McLaughlin, K. A., ... Merikangas, K. R. (2012). Severity of 12-month DSM-IV disorders in the National Comorbidity Survey replication adolescent supplement. *Archives of General Psychiatry*, 69, 381–389. doi: 10.1001/archgenpsychiatry.2011.1603
- Kessler, R. C., Avenevoli, S., & Ries Merikangas, K. (2001). Mood disorders in children and adolescents: An epidemiologic perspective. *Biological Psychiatry*, 49, 1002–1014. doi: 10.1016/S0006-3223(01)01129-5
- Klesges, R. C., Obarzanek, E., Kumanyika, S., Murray, D. M., Klesges, L. M., Relyea, G. E., ... Slawson, D. L. (2010). The Memphis Girls' Health Enrichment Multi-site Studies (GEMS): An evaluation of the efficacy of a 2-year obesity prevention in African American girls. *Archives of Pediatric and Adolescent Medicine*, 164, 1007–1014. doi: 10.1001/archpediatrics.2010.196
- Knerr, W., Gardner, F., & Cluver, L. (2013). Improving positive parenting skills and reducing harsh and abusive parenting in low- and middle-income countries: A systematic review. *Prevention Science*, 14, 352–363. doi: 10.1007/s11121-012-0314-1
- Koniak-Griffin, D., Lesser, J., Nyamathi, A., Uman, G., Stein, J. A., & Cumberland, W. G. (2003). Project CHARM: An HIV prevention program for adolescent mothers. *Family and Community Health*, 26, 94–107. doi: 10.1097/00003727-200304000-00003
- Kovacs, M., Gatsonis, C., Paulauskas, S. L., & Richards, C. (1989). Depressive disorders in childhood: III. A longitudinal study of comorbidity with and risk for anxiety disorders. *Archives of General Psychiatry*, 46, 776–782. doi: 10.1001/archpsyc.1989.01810090018003
- Kulis, S., Marsiglia, F. F., Elek, E., Dustman, P., Wagstaff, D. A. & Hecht, M. L. (2005). Mexican/Mexican American adolescents and 'leepin' It R.E.A.L.: An evidence-based substance use prevention program. *Children & Schools*, 27, 133–145. doi: 10.1093/cs/27.3.133
- Kumpfer, K. L., & Alvarado, R. (1995). Strengthening families to prevent drug use in multiethnic youth. In G. Botvin, S. Schinke, & M. Orlandi (Eds.), *Drug abuse prevention with multiethnic youth* (pp. 253–292). Newbury Park, CA: Sage.
- Kumpfer, K. L., Alvarado, R., Smith, P., & Bellamy, N. (2002). Cultural sensitivity and adaptation in family-based prevention interventions. *Prevention Science*, 3, 241–246. doi: 10.1023/A:1019902902119
- Kumpfer, K. L., Alvarado, R., Tait, C., & Turner, C. (2002). Effectiveness of school-based family and children's skills training for substance abuse prevention among 6–8-year-old rural children. *Psychology of Addictive Behaviors*, 16, Suppl.4). S65–S71.
- Kumpfer, K. L., Pinyuchon, M., de Melo, A. T., & Whiteside, H. O. (2008). Cultural adaptation process for international dissemination of the strengthening families program. *Evaluation & the Health Professions*, 31, 226–239. doi: 10.1177/0163278708315926
- Kumanyika, S. K., Story, M., Beech, B. M., Sherwood, N. E., Baranowski, J. C., Powell, T. M., ... Owens, A. S. (2003). Collaborative planning for formative assessment and cultural appropriateness in the Girls Health Enrichment Multi-site Studies (GEMS): A retrospective. *Ethnicity and Disease*, 13, S1–15–S1–29.
- Lansford, J. E., Chang, L., Dodge, K. A., Malone, P. S., Oburu, P., Palmérus, K., ... Quinn, N. (2005). Physical discipline and children's adjustment: Cultural normativeness as a moderator. *Child Development*, 76, 1234–1246. doi: 10.1111/j.1467-8624.2005.00847.x
- Lau, A. S. (2006). Making the case for selective and directed cultural adaptations of evidence-based treatments: Examples from parent training. *Clinical Psychology: Science and Practice*, 13, 295–310. doi: 10.1111/j.1467-8624.2005.00847.x
- Lau, A. S. (2010). Physical discipline in Chinese American immigrant families: An adaptive culture perspective. *Cultural Diversity and Ethnic Minority Psychology*, 16, 313–322. doi: 10.1037/a0018667
- Lau, A. S. (2012). Reflections on adapting parent training for Chinese immigrants: Blind alleys, thoroughfares, and test drives. In G. Bernal & M. M. Domenech Rodríguez (Eds.), *Cultural adaptations: Tools for evidence-based practice with diverse populations* (pp. 133–156). Washington, DC: American Psychological Association.
- Lau, A. S., Chang, D. F., Okazaki, S., & Bernal, G. (in press). Psychotherapy research with ethnic minorities: What is the research agenda? In N. W. S. Zane, G. Bernal, & F. T. L. Leong (Eds.), *Cultural-informed evidence-based practices for ethnic minorities: Challenges and solutions*. Washington, DC: American Psychological Association.

- Lau, A. S., Fung, J. J., Ho, L. Y., Liu, L. L., & Gudiño, O. G. (2011). Parent training with high-risk immigrant Chinese families: A pilot group randomized trial yielding practice-based evidence. *Behavior Therapy, 42*, 413–426. doi: 10.1016/j.beth.2010.11.001
- Lau, A. S., Fung, J. J., & Yung, V. (2010). Group parent training with immigrant Chinese families: Enhancing engagement and augmenting skills training. *Journal of Clinical Psychology: In Session, 66*, 880–894. doi: 10.1002/jclp.20711
- Lesser, J., Koniak-Griffin, D., Huang, R., Takayanagi, S., & Cumberland, W. G. (2009). Parental protectiveness and unprotected sexual activity among Latino adolescent mothers and fathers. *AIDS Education & Prevention, 21*, 88–102. doi: 10.1521/aeap.2009.21.5_suppl.88
- Lieh-Mak, F., Lee, P. W. H., & Luk, S. L. (1984). Problems encountered in teaching Chinese parents to be behavior therapists. *Psychologia, 27*, 56–64.
- Littlefield, L., Story, K., Woolcock, C., Trinder, M., Burke, S., & Reid, K. (2005). *Exploring Together preschool program: Program manual*. Melbourne, Australia: Victorian Parenting Centre.
- Lock, S., & Barrett, P. M. (2003). A longitudinal study of developmental differences in universal preventive intervention for child anxiety. *Behaviour Change, 20*, 183–199. doi: 10.1375/behc.20.4.183.29383
- Lowry-Webster, H. M., Barrett, P. M., & Dadds, M. R. (2001). A universal prevention trial of anxiety and depressive symptomatology in childhood: Preliminary data from an Australian study. *Behaviour Change, 18*, 36–50. doi: 10.1375/behc.18.1.36
- Luepker, R. V., Perry, C. L., McKinlay, S. M., Nader, P. R., Parcel, G. S., Stone, E. J., . . . Smisson, J. (1996). Outcomes of a field trial to improve children's dietary patterns and physical activity. *Journal of the American Medical Association, 275*, 768–776. doi: 10.1001/jama.1996.03530340032026
- MacKinnon, D. P., Fairchild, A. J., & Fritz, M. S. (2007). Mediation analysis. *Annual Review of Psychology, 58*, 593–614. doi: 10.1146/annurev.psych.58.110405.085542
- MacLachlan, M. (2006). *Culture and health: A critical perspective towards global health (2nd ed.)*. Hoboken, NJ: Wiley.
- Magarey, A. M., Daniels, L. A., Boulton, T. J., & Cockington, R. A. (2003). Predicting obesity in early adulthood from childhood and parental obesity. *International Journal of Obesity, 27*, 505–513. doi: 10.1038/sj.ijo.0802251
- Mansfield, C. J., Conroy, M. E., Emans, S. J., & Woods, E. R. (1993). A pilot study of AIDS education and counseling of high-risk adolescents in an office setting. *Journal of Adolescent Health, 14*, 115–119. doi: 10.1016/1054-139X(93)90095-7
- Markus, H. R., & Kitayama, S. (1991). Culture and the self: Implications for cognition, emotion, and motivation. *Psychological Review, 98*, 224–253. doi: 10.1037/0033-295X.98.2.224
- Martinez, C. R. (2006). Effects of differential family acculturation on Latino adolescent substance use. *Family Relations: An Interdisciplinary Journal of Applied Family Studies, 55*, 306–317. doi: 10.1111/j.1741-3729.2006.00404.x
- Martinez, C. R., & Eddy, J. M. (2005). Effects of culturally adapted parent management training on Latino youth behavioral health outcomes. *Journal of Consulting and Clinical Psychology, 73*, 841–851. doi: 10.1037/0022-006X.73.5.841
- Matos, M., Bauermeister, J. J., & Bernal, G. (2009). Parent-child interaction therapy for Puerto Rican preschool children with ADHD and behavior problems: A pilot efficacy study. *Family Process, 48*, 232–52. doi: 10.1111/j.1545-5300.2009.01279.x
- Matos, M., Torres, R., Santiago, R., Jurado, M., & Rodríguez, I. (2006). Adaptation of parent-child interaction therapy for Puerto Rican families: A preliminary study. *Family Process, 45*, 205–22. doi: 10.1111/j.1545-5300.2006.00091.x
- McCabe, C. K., Baptiste, D., Traube, D., Paikoff, R. L., Madison-Boyd, S., Coleman, D., & McKay, M. M. (2007). Family-based HIV preventive intervention: Child level results from the CHAMP family program. *Social Work in Mental Health, 5*, 203–220. doi: 10.1300/J200v05n01_10
- McCabe, K., & Yeh, M. (2009). Parent-Child Interaction Therapy for Mexican Americans: A randomized clinical trial. *Journal of Clinical Child and Adolescent Psychology, 38*, 753–759. doi: 10.1080/15374410903103544
- McCabe, K. M., Yeh, M., Garland, A. F., Lau, A. S., & Chavez, G. (2005). The GANA program: A tailoring approach to adapting Parent Child Interaction Therapy for Mexican Americans. *Education & Treatment of Children, 28*, 111–129.
- McCabe, K., Yeh, M., Lau, A., & Argote, C. B. (2012). Parent-Child Interaction Therapy for Mexican Americans: Results of a pilot randomized clinical trial at follow-up. *Behavior Therapy, 43*, 606–618. doi: 10.1016/j.beth.2011.11.001
- McGoldrick, M., & Giordano, J. (1996). *Ethnicity and family therapy (2nd ed.)*. New York, NY: Guilford Press.
- McKay, M., Block, M., Mellins, C., Traube, D. E., Brackis-Cott, E., Minott, D., . . . Abrams, E. J. (2007). Adapting a family-based HIV prevention program for HIV-infected preadolescents and their families: Youth, families and health care providers coming together to address complex needs. *Social work in mental health, 5*(3–4), 355–378. doi: 10.1300/J200v05n03_06
- McKay, M. M., Chasse, K. T., Paikoff, R., McKinney, L. D., Baptiste, D., Coleman, D., & Bell, C. C. (2004). Family-level impact of the CHAMP Family Program: A community collaborative effort to support urban families and reduce youth HIV risk exposure. *Family Process, 43*, 79–93. doi: 10.1111/j.1545-5300.2004.04301007.x
- McKleroy, V. S., Galbraith, J. S., Cummings, B., Jones, P., Harshbarger, C., Collins, C., . . . Adapt Team. (2006). Adapting evidence-based behavioral interventions for new settings and target populations. *AIDS Education and Prevention, 18*, 59–73. doi: 10.1521/aeap.2006.18.supp.59
- Miller, G. E., & Prinz, R. J. (1990). Enhancement of social learning family interventions for childhood conduct disorder. *Psychological Bulletin, 108*, 291–307. doi: 10.1037/0033-2909.108.2.291
- Miller, L. D., Laye-Gindhu, A., Bennett, J. L., Liu, Y., Gold, S., March, J. S., . . . Waechter, V. E. (2011). An effectiveness study of a culturally enriched school-based CBT anxiety prevention program. *Journal of Clinical Child and Adolescent Psychology, 40*, 618–629. doi: 10.1080/15374416.2011.581619
- Minkler, M., & Wallerstein, N. (Eds.). (2003). *Community-based participatory research for health*. San Francisco, CA: Jossey-Bass.
- Morawska, A., Sanders, M., Goadby, E., Headley, C., Hodge, L., McAuliffe, C., . . . Anderson, E. (2011). Is the Triple P-Positive Parenting Program acceptable to parents from culturally diverse backgrounds? *Journal of Child and Family Studies, 20*, 614–622. doi: 10.1007/s10826-010-9436-x
- Mrazek, P. J., & Haggerty, R. J. (Eds.). (1994). *Reducing risks for mental disorders: Frontiers for preventive intervention research*. Washington, DC: National Academy Press.
- Muñoz, R. F., & Mendelson, T. (2005). Toward evidence-based interventions for diverse populations. *Journal of Consulting and Clinical Psychology, 73*, 790–799. doi: 10.1037/0022-006X.73.5.790
- Murry, V. M., Berkel, C., Brody, G. H., Gibbons, M., & Gibbons, F. X. (2007). The Strong African American Families program: Longitudinal pathways to sexual risk reduction. *Journal of Adolescent Health, 41*, 333–342. doi: 10.1016/j.jadohealth.2007.04.003
- Murry, V. M., Berkel, C., Brody, G. H., Miller, S. J., & Chen, Y. F. (2009). Linking parental socialization to interpersonal protective processes,

- academic self-presentation, and expectations among rural African American youth. *Cultural Diversity and Ethnic Minority Psychology*, 15(1), 1–10. doi: 10.1037/a0013180
- Murry, V. M., Berkel, C., Pantin, H., & Guillermo, P. (2011). Family-based HIV prevention with African American and Hispanic youth. In W. Pequegnat & C. Bell (Eds.), *Families and HIV/AIDS* (pp. 229–248). New York, NY: Springer.
- Murry, V. M., McNair, L. D., Myers, S. S., Chen, F., & Brody, G. H. (2014). Intervention induced changes in perceptions of parenting and risk opportunities among rural African American youth. *Journal of Child and Family Studies*, 23, 422–436. Advanced online publication. doi: 10.1007/s10826-013-9714-5
- Murry, V. M., Smith, E. P., & Hill, N. E. (2001). Race, ethnicity, and culture in studies of families in context. *Journal of Marriage and the Family*, 63, 911–914. doi: 10.1111/j.1741-3737.2001.00911.x
- Must, A., Jacques, P. F., Dallal, G. E., Bajema, C. J., & Dietz, W. H. (1992). Long-term morbidity and mortality of overweight adolescents: A follow-up of the Harvard Growth Study of 1922–1935. *New England Journal of Medicine*, 327, 1350–1355. doi: 10.1056/NEJM19921053271904
- Myers, H. F., Alvy, K. T., Arlington, A., Richardson, M. A., Marigna, M., Huff, R., . . . Newcomb, M. D. (1992). The impact of a parent training program on inner-city African-American families. *Journal of Community Psychology*, 20, 132–147. doi: 10.1002/1520-6629
- Nader, P. R., Stone, E. J., Lytle, L. A., Perry, C. L., Osganian, S. K., Kelder, S., . . . Luepker, R. V. (1999). Three-year maintenance of improved diet and physical activity: The CATCH cohort. *Child and Adolescent Trial for Cardiovascular Health. Archives of Pediatric and Adolescent Medicine*, 153, 695–704. doi: 10.1001/archpedi.153.7.695
- Narayan, K. M. V., Boyle, J. P., Thompson, T. J., Sorensen, S. W., & Williamson, D. F. (2003). Lifetime risk for diabetes mellitus in the United States. *Journal of the American Medical Association*, 290, 1884–1890. doi: 10.1001/jama.290.14.1884
- National Institute of Drug Abuse. (1997). *Prevention of drug use among children and adolescents: A research-based guide*. Retrieved from: <http://www.drugabuse.gov/publications/preventing-drug-abuse-among-children-adolescents/prevention-principles>
- National Institutes of Health. (1994). *NIH guidelines for the inclusion of women and minorities as subjects in clinical research*. Retrieved from <http://grants.nih.gov/grants/guide/noticefilesnot94-100.html>
- National Research Council and Institute of Medicine. (2009). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities*. Committee on the Prevention of Mental Disorders and Substance Abuse among Children, Youth, and Young Adults: Research Advances and Promising Interventions. M. E. O'Connell, T. Boat, & K. E. Warner (Eds.). Washington, DC: National Academies Press.
- Nelson, K., & Nazbah, T. (2011). Evaluation of a substance abuse, HIV and hepatitis prevention initiative for urban Native Americans: The Native Voices program. *Journal of Psychoactive Drugs*, 43, 349–354. doi: 10.1080/02791072.2011.629158
- Ogden, T., Amlund-Hagen, K., Askeland, E., & Christensen, B. (2009). Implementing and evaluating evidence-based treatments of conduct problems in children and youth in Norway. *Research on Social Work Practice*, 19, 582–591. doi: 10.1177/1049731509335530
- Ogden, T., & Hagen, K. A. (2008). Treatment effectiveness of parent management training in Norway: A randomized controlled trial of children with conduct problems. *Journal of Consulting and Clinical Psychology*, 76, 607–621. doi: 10.1037/0022-006X.76.4.607
- Olds, D. L., Eckenrode, J., Henderson, C. R., Jr., Kitzman, H., Powers, J., Cole, R. S., . . . Luckey, D. (1997). Long-term effects of home visitation on maternal life course and child abuse and neglect: Fifteen-year follow-up of a randomized trial. *Journal of the American Medical Association*, 278, 637–643. doi: 10.1001/jama.1997.03550080047038
- Ortiz, C., & Del Vecchio, T. (2013). Cultural diversity: Do we need a new wake-up call for parent training? *Behavior Therapy*, 44, 443–458. doi: 10.1016/j.beth.2013.03.009
- Orrell-Valente, J., Pinderhughes, E. E., Valente, E., & Laird, R. D. (1999). If it's offered, will they come? Influences on parents' participation in a community-based conduct problems prevention program. *American Journal of Community Psychology*, 27, 753–783. doi: 10.1023/A:1022258525075
- Oyserman, D., & Markus, H. R. (1990). Possible selves and delinquency. *Journal of Personality and Social Psychology*, 59, 112–125. doi: 10.1037/0022-3514.59.1.112
- Pantin, H., Coatsworth, J. D., Feaster, D., Newman, F., Briones, E., Prado, G., . . . Szapocznik, J. (2003). Familias Unidas: The efficacy of an intervention to promote parental investment in Hispanic immigrant families. *Prevention Science*, 4, 189–201. doi: 10.1023/a:1024601906942
- Pantin, H., Schwartz, S. J., Sullivan, S., Prado, G., & Szapocznik, J. (2004). Ecodevelopmental HIV prevention programs for Hispanic adolescents. *American Journal of Orthopsychiatry*, 74, 545–558. doi: 10.1037/0002-9432.74.4.545
- Patterson, G. R., Chamberlain, P., & Reid, J. B. (1982). A comparative evaluation of a parent-training program. *Behavior Therapy*, 13, 638–650. doi: 10.1016/S0005-7894(82)80021-X
- Patterson, G. R., Reid, J. B., & Dishion, T. J. (1992). *A social interactional approach: Vol. 4: Antisocial boys*. Eugene, OR: Castalia.
- Pemberton, J. R., & Borrego, J., Jr. (2007). Increasing acceptance of behavioral child management techniques: What do parents say? *Child & Family Behavior Therapy*, 29, 27–45. doi: 10.1300/J019v29n02_03
- Pentz, P. A., Dwyer, J. H., MacKinnon, D. P., Flay, B. R., Phil, D., Hansen, W. B., . . . Johnson, C. A. (1989). A multicommunity trial for primary prevention of adolescent drug use. *Journal of the American Medical Association*, 261, 3259–3266. doi: 10.1001/jama.261.22.3259
- Pina, A. A., Villalta, I. K., & Zerr, A. A. (2009). Exposure-based cognitive behavioral treatment of anxiety in youth: An emerging culturally-prescriptive framework. *Behavioral Psychology*, 17, 111–135.
- Pina, A. A., Zerr, A. A., Gonzales, N. A., & Ortiz, C. D. (2009). Psychosocial interventions for school refusal behavior in children and adolescents. *Child Development Perspectives*, 3, 11–20. doi: 10.1111/j.1750-8606.2008.00070.x
- Pina, A. A., Zerr, A. A., Villalta, I. K., & Gonzales, N. A. (2012). Indicated prevention and early intervention for childhood anxiety: A randomized trial with Caucasian and Hispanic/Latino youth. *Journal of Consulting and Clinical Psychology*, 80, 940–946. doi: 10.1037/a0029460
- Prado, G., Pantin, H., Briones, E., Schwartz, S. J., Feaster, D., Huang, S., . . . Szapocznik, J. (2007). A randomized controlled trial of a parent-centered intervention in preventing substance use and HIV risk behaviors in Hispanic adolescents. *Journal of Consulting and Clinical Psychology*, 75, 914–926. doi: 10.1037/0022-006X.75.6.914
- Prochaska, J. O., & DiClemente, C. C. (1992). Stages of change in the modification of problem behaviors. In M. Hersen, R. M. Eisler, & P. M. Miller (Eds.), *Progress in behavior modification* (pp. 184–214). Sycamore, IL: Sycamore Press.
- Pruitt, L. R. (2009). The forgotten fifth: Rural youth and substance abuse. *Stanford Law and Policy Review*, 20, 259–304.
- Rapee, R. M., Kennedy, S., Ingram, M., Edwards, S., & Sweeney, L. (2005). Prevention and early intervention of anxiety disorders in inhibited preschool children. *Journal of Consulting and Clinical Psychology*, 73, 488–497. doi: 10.1037/0022-006X.73.3.488

- Reid, M. J., Webster-Stratton, C., & Baydar, N. (2004). Halting the development of conduct problems in Head Start children: The effects of parenting training. *Journal of Clinical Child and Adolescent Psychology, 33*, 279–291. doi: 10.1207/s15374424jccp3302_10
- Reid, M. J., Webster-Stratton, C., & Beauchaine, T. P. (2001). Parent training in Head Start: A comparison of program response among African American, Asian American, Caucasian, and Hispanic mothers. *Prevention Science, 2*, 209–227. doi: 10.1023/A:1013618309070
- Renzaho, A. M. N. (2002). *Addressing the needs of refugees and humanitarian entrants in Victoria: An evaluation of health & community services*. Melbourne, Australia: Centre for Culture Ethnicity and Health.
- Renzaho, A. M. N., Swinburn, B., & McCabe, M. (2008). Sociocultural aspect of obesity among African migrants in Australia: Examining the intergenerational conflict. In *Population Health Congress 2008: A Global World Practical Action for Health and Wellbeing* (p. 165).
- Renzaho, A. M. N., & Vignjevic, S. (2011). The impact of a parenting intervention in Australia among migrants and refugees from Liberia, Sierra Leone, Congo, and Burundi: Results from the African Migrant Parenting Program. *Journal of Family Studies, 17*(1), 71–79. doi: 10.5172/jfs.2011.17.1.71
- Resnicow, K., Baranowski, T., Ahluwalia, J. S., & Braithwaite, R. L. (1999). Cultural sensitivity in public health: Defined and demystified. *Ethnicity and Disease, 9*, 10–21.
- Resnicow, K., Soler, R., Braithwaite, R. L., Ahluwalia, J. S., & Butler, J. (2000). Cultural sensitivity in substance use prevention. *Journal of Community Psychology, 28*, 271–290. doi: 10.1002/(SICI)1520-6629(200005)
- Resnicow, K., Taylor, R., Baskin, M., & McCarty, F. (2005). Results of Go Girls: A weight control program for overweight African-American adolescent females. *Obesity Research, 13*, 1739–1748. doi: 10.1038/oby.2005.212
- Reyno, S. M., & McGrath, P. J. (2006). Predictors of parent training efficacy for child externalizing behavior problems—a meta-analytic review. *Journal of Child Psychology & Psychiatry, 47*, 99–111. doi: 10.1111/j.1469-7610.2005.01544.x
- Riolo, S. A., Nguyen, T. A., Greden, J. F., & King, C. A. (2005). Prevalence of depression by race/ethnicity: Findings from the National Health and Nutrition Examination Survey III. *American Journal of Public Health, 95*, 998–1000. doi: 10.2105/AJPH.2004.047225
- Robinson, G., Tyler, W., Jones, Y., Silburn, S., & Zubrick, S. R. (2012). Context, diversity and engagement: Early intervention with Australian Aboriginal families in urban and remote contexts. *Children & Society, 26*, 343–355. doi: 10.1111/j.1099-0860.2010.00353.x
- Robinson, T. N., Killen, J. D., Kraemer, H. C., Wilson, D. M., Matheson, D. M., Haskell, W. L., ... Varady, A. (2003). Dance and reducing television viewing to prevent weight gain in African-American girls: The Stanford GEMS pilot study. *Ethnicity and Disease, 13*(1, suppl.), S65–S77.
- Robinson, T. N., Kraemer, H. C., Matheson, D. M., Obarzanek, E., Wilson, D. M., Haskell, W. L., ... Killen, J. D. (2008). Stanford GEMS phase 2 obesity prevention trial for low-income African-American girls: Design and sample baseline characteristics. *Contemporary Clinical Trials, 29*, 56–69. doi: 10.1016/j.cct.2007.04.007
- Robinson, T. N., Matheson, D. M., Kraemer, H. C., Wilson, D. M., ... & Killen, J. D. (2010). A randomized controlled trial of culturally tailored dance and reducing screen time to prevent weight gain in low-income African American girls: Stanford GEMS. *Archives of Pediatric and Adolescent Medicine, 164*, 995–1004. doi: 10.1001/archpediatrics.2010.197
- Rogler, L. H., Malgady, R. G., Constantino, G., & Blumenthal, R. (1987). What do culturally sensitive mental health services mean? The case of Hispanics. *American Psychologist, 42*, 565–570. doi: 10.1037/0003-066X.42.6.565
- Rotheram-Borus, M. J., Swendeman, D., & Chorpita, B. F. (2012). Disruptive innovations for designing and diffusing evidence-based interventions. *American Psychologist, 67*, 463–476. doi: 10.1037/a0028180
- Sanders, M. R. (1999). Triple P-Positive Parenting Program: Towards an empirically validated multilevel parenting and family support strategy for the prevention of behavior and emotional problems in children. *Clinical Child and Family Psychology Review, 2*, 71–90. doi: 10.1023/A:1021843613840
- Sanders, M. R. (2008). Triple P-Positive Parenting Program as a public health approach to strengthening parenting. *Journal of Family Psychology, 22*, 506. doi: 10.1037/0893-3200.22.3.506
- Sanders, M. R. (2012). Development, evaluation, and multinational dissemination of the triple P-positive parenting program. *Annual Review of Clinical Psychology, 8*, 345–379. doi: 10.1146/annurev-clinpsy-032511-143104
- Sandler, I. N., Ayers, T. S., Wolchik, S. A., Tein, J., Kwok, O., Haine, R. A., ... Griffin, W. A. (2003). The family bereavement program: Efficacy evaluation of a theory-based prevention program for parentally bereaved children and adolescents. *Journal of Consulting and Clinical Psychology, 71*(3), 587–600. doi:10.1037/0022-006X.71.3.587.
- Santelli, J. S., Lindberg, L. D., Abma, J., McNeely, C. S. & Resnick, M. (2000). Adolescent sexual behavior: Estimates and trends from four national surveys. *Family Planning Perspectives, 32*, 156–165. doi: 10.2307/12648232
- Schinke, S., Gilchrist, L. D., Schilling, R. F., II, Walker, R. D., Locklear, V. S., & Kitajima, E. (1986). Smokeless tobacco use among Native American adolescents. *New England Journal of Medicine, 314*, 1051–1052. doi: 10.1056/NEJM198604173141614
- Schinke, S. P., Moncher, M. S., Holden, G. W., Botvin, G. J., & Orlandi, M. A. (1989). American Indian youth and substance abuse: Tobacco use problems, risk factors and preventive interventions. *Health Education Research, 4*, 137–144. doi: 10.1093/her/4.1.137
- Schinke, S. P., Tepavac, L., & Cole, K. C. (2000). Preventing substance use among Native American youth: Three-year results. *Addictive Behaviors, 25*, 387–397. doi: 10.1016/S0306-4603(99)00071-4
- Schoenwald, S. K. (2008). Toward evidence-based transport of evidence-based treatments: MST as an example. *Journal of Child and Adolescent Substance Abuse, 17*, 69–91. doi: 10.1080/15470650802071671
- Schroeder, B. A., Messina, A., Schroeder, D., Good, K., Barto, S., Saylor, J., & Masiello, M. (2012). The implementation of a statewide bullying prevention program: Preliminary findings from the field and the importance of coalitions. *Health Promotion and Practice, 13*, 489–95. doi: 10.1177/1524839910386887
- Serdula, M. K., Ivery, D., Coates, R. J., Freedman, D. S., Williamson, D. F., & Byers, T. (1992). Do obese children become obese adults? A review of the literature. *Preventive Medicine, 22*, 167–177. doi: 10.1006/pmed.1993.1014
- Sheldrick, R., Kendall, P. C., & Heimberg, R. (2001). The clinical significance of treatments: A comparison of three treatments for conduct disordered children. *Clinical Psychology: Science and Practice, 8*, 418–430. doi: 10.1093/clipsy.8.4.418
- Shrier, L. A., Angheta, R., Goodman, E., Chiou, V. M., Lyden, M. R., & Emans, S. J. (2001). Randomized controlled trial of a safer sex intervention for high-risk adolescent girls. *Archives of Pediatrics & Adolescent Medicine, 155*, 73–79. doi: 10.1001/archpedi.155.1.73
- Sikkema K. J., Anderson, E. S., Kelly, J. A., Winett, R. A., Gore-Felton, C., Roffman, R. A., ... & Brondino, M. J. (2005). Outcomes of a randomized, controlled community-level HIV prevention intervention for adolescents in low-income housing developments. *AIDS, 19*, 1509–1516. doi: 10.1097/01.aids.0000183128.39701.34

- Silverman, W. K., Pina, A. A., & Viswesvaran, C. (2008). Evidence-based psychosocial treatments for phobic and anxiety disorders in children and adolescents. *Journal of Clinical Child and Adolescent Psychology, 37*, 105–130. doi: 10.1080/15374410701817907
- Simon, E., Bögels, S. M., & Voncken, J. M. (2011). Efficacy of child-focused and parent-focused interventions in a child anxiety prevention study. *Journal of Clinical Child and Adolescent Psychology, 40*(2), 204–219.
- Simons-Morton, B. G., Parcel, G. S., Baranowski, T., Forthofer, R., & O'Hara, N. M. (1991). Promoting physical activity and a healthful diet among children: Results of a school-based intervention study. *American Journal of Public Health, 81*, 986–991. doi: 10.2105/AJPH.81.8.986
- Sloboda, Z., & Bukoski, W. J. (2003). *Handbook of drug abuse prevention*. New York, NY: Springer.
- Smedley, B. D., Stith, A. Y., & Nelson, A. R., (2002) *Unequal treatment: Confronting racial and ethnic disparities in health care*. Washington, DC: Institute of Medicine.
- Smith, T. B., Domenech Rodríguez, M. M., & Bernal, G. (2011). Culture. *Journal of Clinical Psychology: In Session, 67*, 166–175. doi: org/10.1002/jclp.20757
- Spoth, R. (1997). Challenges in defining and developing the field of rural mental disorder prevention intervention research. *American Journal of Community Psychology, 25*, 425–448. doi: 10.1023/A:1024603504781
- Spoth, R., & Molgaard, V. (1999). Project Family: A partnership integrating research with the practice of promoting family and youth competencies. In T. R. Chibucos & R. Lerner (Eds.), *Serving children and families through community-university partnerships: success stories* (pp. 127–137). Boston, MA: Kluwer Academic.
- Spoth, R., Redmond, C., & Shin, C. (2001). Randomized trial of brief family interventions for general populations: Adolescent substance use outcomes four years following baseline. *Journal of Consulting and Clinical Psychology, 69*, 627–642. doi: 10.1037/0022-006X.69.4.627
- Spoth, R., Rohrbach, L. A., Greenberg, M., Leaf, P., Brown, C. H., Fagan, A. C., . . . Hawkins, J. D. (2013). Addressing the core challenges for the next generation of Type 2 translation research and systems: The Translation Science to Population Impact (TSci Impact) Framework. *Prevention Science, 14*, 319–351. doi: 10.1007/s11121-012-0362-6
- Stanton, B., Li, X., Black M., Ricardo, B., Balbriath, J., Kaljee, L., & Feigelman, S. (1994). Sexual practices and intentions among pre-adolescent and early adolescent low income urban African Americans. *Pediatrics, 93*, 966–973.
- Stice, E., Shaw, H., Bohon, C., Marti, C. N., & Rohde, P. (2009). A meta-analytic review of depression prevention programs for children and adolescents: Factors that predict magnitude of intervention effects. *Journal of Consulting and Clinical Psychology, 77*, 486–503. doi: 10.1037/0022-006X.70.4.1010
- Szapocznik, J., & Coatsworth, J. D. (1999). An ecodevelopmental framework for organizing the influences on drug abuse: A developmental model of risk and protection. In M. D. Glantz & C. R. Hartel (Eds.), *Drug abuse: Origins & interventions* (pp. 331–366). Washington, DC: American Psychological Association.
- Suarez-Orozco, C., & Suarez-Orozco, M. (1995). *Transformations: Immigration, family life, and achievement motivation among Latino adolescents*. Stanford, CA: Stanford University Press.
- Sue, S. (1998). In search of cultural competence in psychotherapy and counseling. *American Psychologist, 53*, 440–448. doi: 10.1037/0003-066X.53.4.440
- Sue, S. (2006). Cultural competency: From philosophy to research and practice. *Journal of Community Psychology, 34*, 237–245. doi: 10.1002/jcop.20095
- Sue, S., & Morishima, J. K. (1982). *The mental health of Asian Americans*. San Francisco, CA: Jossey-Bass.
- Sundell, K., Ferrer-Wreder, L. F., & Fraser, M. W. (2014). Going global: A model for evaluating empirically supported family-based interventions in new contexts. *Evaluation & the Health Professions, 37*, 203–230. doi: 10.1177/0163278712469813.
- Swedish National Board of Health and Welfare. (2011). *Svensk och internationell forskning om sociala interventioners effekter* [Swedish and international research on the effects of social interventions]. Stockholm, Sweden: Author.
- Szapocznik, J., & Kurtines, W. M. (1993). Family psychology and cultural diversity: Opportunities for theory, research and application. *American Psychologist, 48*, 400–407. doi: 10.1037/0003-066X.48.4.400
- Tein, J. Y., Sandler, I. N., MacKinnon, D. P., & Wolchik, S. A. (2004). How did it work? Who did it work for? Mediation in the context of a moderated prevention effect for children of divorce. *Journal of Consulting and Clinical Psychology, 72*, 617–624. doi: 10.1037/0022-006X.72.4.617
- Teubert, D., & Pinquart, M. (2011). A meta-analytic review on the prevention of symptoms of anxiety in children and adolescents. *Journal of Anxiety Disorders, 25*, 1046–1059. doi: 10.1016/j.janxdis.2011.07.001
- Teufel, N. I., Perry, C. L., Story, M., Flint-Wagner, H. G., Levin, S., Clay, T. E., . . . Pablo, J. L. (1999). Pathways family intervention for third-grade American Indian children. *American Journal of Clinical Nutrition, 69*(Suppl.), 803S–809S.
- Treviño, R. P., Hernandez, A. E., Yin, Z., Garcia, O. A., & Hernandez, I. (2005). Effect of the Bienestar health program on physical fitness in low-income Mexican American children. *Hispanic Journal of Behavioral Sciences, 27*, 120–132. doi: 10.1177/0739986304272359
- Treviño, R. P., Pugh, J. A., Hernandez, A. E., Menchaca, V. D., Ramirez, R. R., & Mendoza, M. (1998). Bienestar: A diabetes risk-factor prevention program. *Journal of School Health, 68*, 62–67. doi: 10.1111/j.1746-1561.1998.tb07192.x
- Treviño, R. P., Yin, Z., Hernandez, A., Hale, D. E., Garcia, O. A., & Mobley, C. (2004). Impact of the Bienestar school-based diabetes mellitus prevention program on fasting capillary glucose levels. *Archives of Pediatric and Adolescent Medicine, 158*, 911–917. doi: 10.1001/archpedi.158.9.911
- Trimble, J. E. 1995. Toward and understanding of ethnicity and ethnic identity, and their relationship with drug use research. In G. J. Botvin, S. Schinke, & M. A. Orlandi (Eds.), *Drug abuse prevention with multi-ethnic youth* (pp. 3–27). Thousand Oaks, CA: Sage
- Turner, K. M., Richards, M., & Sanders, M. R. (2007). Randomised clinical trial of a group parent education programme for Australian Indigenous families. *Journal of Paediatrics and Child Health, 43*, 429–437. doi: 10.1111/j.1440-1754.2007.01053.x
- United States Census Bureau. (2010). *Overview of race and hispanic origin: 2010*. Retrieved from <http://www.census.gov/prod/cen2010/briefs/c2010br-02.pdf>
- United States Department of Health and Human Services. (2001). *Eliminating health disparities in the United States*. HRSA Archives. Retrieved from: <http://archive.hrsa.gov/NewsBrief/2001/eliminating>
- United States Department of Health and Human Services. (2010). *Affordable Care Act: Prevention and wellness*. Retrieved from: <http://www.hhs.gov/healthcare/prevention/index.html>
- Villarruel, A. M., Cherry, C. L., Cabriaes, E. G., Ronis, D. L., & Zhou, Y. (2008). A parent-adolescent intervention to increase sexual risk communication: Results of a randomized control trial. *AIDS Education and Prevention, 20*, 371–383. doi: 10.1521/aeap.2008.20.5.371
- Wandersman, A. (2003). Community science: Bridging the gap between science and practice with community-centered models. *American Journal of Community Psychology, 31*, 227–242. doi: 10.1023/A:1023954503247

- Wang, Y., & Beydoun, M. A. (2007). The obesity epidemic in the United States—gender, age, socioeconomic, racial/ethnic, geographic characteristics: A systematic review and meta-regression analysis. *Epidemiologic Reviews*, *29*, 6–28. doi: 10.1093/epirev/mxm007
- Warner, V., Weissman, M. M., Fendrich, M., Wickramaratne, P., & Moreau, D. (1992). The course of major depression in the offspring of depressed parents: Incidence, recurrence, and recovery. *Archives of General Psychiatry*, *49*(10), 795–801. Retrieved from <http://login.ezproxy1.lib.asu.edu/login?url=http://search.proquest.com/docview/618251989?accountid=4485>.
- Weisner, T. S. (2002). Ecocultural understanding of children's developmental pathways. *Human Development*, *45*, 275–281. doi: 10.1159/000064989
- Weisz, J. R., Sandler, I. N., Durlak, J. A., & Anton, B. S. (2005). Promoting and protecting youth mental health through evidence-based prevention and treatment. *American Psychologist*, *60*, 628–648. doi: 10.1037/0003-066X.60.6.628
- Wendler, D., Kington, R., Madans, J., Van Wye, G., Christ-Schmidt, H., Pratt, L. A., . . . Emanuel, E. (2006). Are racial and ethnic minorities less willing to participate in health research? *PLoS Medicine*, *3*, 201–210. doi: 10.1371/journal.pmed.0030201
- Wills, T. A., Gibbons, F. X., Gerrard, M., & Brody, G. H. (2000). Protection and vulnerability processes relevant for early onset of substance use: A test among African American children. *Health Psychology*, *19*, 253–263. doi: 10.1037/0278-6133.19.3.253
- Wills, T. A., Gibbons, F. X., Gerrard, M., Murry, V. M., & Brody, G. H. (2003). Family communication and religiosity related to substance use and sexual behavior in early adolescence: A test for pathways through self-control and prototype perceptions. *Psychology of Addictive Behaviors*, *17*, 312–323. doi: 10.1037/0893-164X.17.4.312
- Wilson, B. D. M., & Miller, R. L. (2003). Examining strategies for culturally grounded HIV prevention: A review. *AIDS Education and Prevention*, *15*, 184–202. doi: 10.1521/aeap.15.3.184.23838
- Wilson, D. K., Van Horn, M. L., Kitzman-Ulrich, H., Saunders, R., Pate, R., Lawman, H. G., . . . Brown, P. V. (2011). Results of the “Active by Choice Today” (ACT) randomized trial for increasing physical activity in low-income and minority adolescents. *Health Psychology*, *30*, 463–471. doi: 10.1037/a0023390
- Wingood, G. M., & DiClemente, R. J. (2008). The ADAPT-ITT model: A model for adapting evidence based HIV interventions. *Journal of Acquired Immunodeficiency Syndrome*, *47*(Suppl. 1), S40–S46. doi: 10.1097/QAI.0b013e3181605df1
- Wolchik, S. A., Sandler, I. N., Jones, S., Gonzales, N., Doyle, K., Winslow, E., . . . Braver, S. L. (2009). The New Beginnings Program for divorcing and separating families: Moving from efficacy to effectiveness. *Family Court Review*, *47*, 416–435. doi: 10.1111/j.1744-1617.2009.01265.x
- Wong, M. D., Tagawa, T., Hsieh, H.-J., Shapiro, M. F., Boscardin, W. J., & Ettner, S. L. (2005). Differences in cause-specific mortality between Latino and White adults. *Medical Care*, *43*, 1058–1062. doi: 10.1097/01.mlr.0000178196.14532.40
- Young, T. (1988). Substance use and abuse among Native Americans. *Clinical Psychology Review*, *8*, 125–138. doi: 10.1016/0272-7358(88)90055-4
- Yu, D. L., & Seligman, M. E. P. (2002). Preventing depressive symptoms in Chinese children. *Prevention & Treatment*, *5*(1), Article 9. doi: 10.1037/1522-3736.5.1.59a
- Zayas, L. H., Bellamy, J. L., & Proctor, E. K. (2012). Considering the multiple service contexts in cultural adaptations of evidence-based practice. In R. C. Brownson, G. A. Colditz, & E. K. Proctor (Eds.), *Dissemination and Implementation Research in Health* (pp. 483–497). New York, NY: Oxford University Press.
- Zollinger, T. W., Saywell, R. M., Overgaard, A. D., Przybylski, M. J., & Dutta-Bergman, M. (2006). Anti-tobacco media awareness of rural youth compared to suburban and urban youth in Indiana. *Journal of Rural Health*, *22*, 119–123. doi: 10.1111/j.1748-0361.2006.00019.x
- Zhou, Q., Sandler, I. N., Millsap, R. E., Wolchik, S. A., & Dawson-McClure, S. R. (2008). Mother-child relationship quality and effective discipline as mediators of the 6-year effects of the New Beginnings Program for children from divorced families. *Journal of Consulting and Clinical Psychology*, *76*, 579–594. doi: 10.1037/0022-006X.76.4.579